

## 2E2 Tutorial Sheet 7<sup>1</sup>

27 November 2005

### Useful facts:

- Z-transform:

$$\mathcal{Z}[(x_n)_{n=0}^{\infty}] = \sum_{n=0}^{\infty} \frac{x_n}{z^n} \quad (1)$$

- Linearity:

$$\mathcal{Z}[a(x_n) + b(y_n)] = aX(z) + bY(z) \quad (2)$$

where  $\mathcal{Z}[(x_n)] = X(z)$  and  $\mathcal{Z}[(y_n)] = Y(z)$

- The Z-transform of a geometric sequence:

$$\mathcal{Z}[(r^n)_{n=0}^{\infty}] = \frac{z}{z - r} \quad (3)$$

- Another useful Z-transform:

$$\mathcal{Z}[(nr^{n-1})_{n=0}^{\infty}] = \frac{z}{(z - r)^2} \quad (4)$$

- Advancing:

$$\mathcal{Z}[(x_{n+1})_{n=0}^{\infty}] = zX(z) - zx_0 \quad (5)$$

where  $\mathcal{Z}[(x_n)] = X(z)$ .

### Questions

1. (2) Find the Z-transform of the sequence  $(2, 0, 1, 0, -3, 0, 0, \dots)$  where all the other entries are zero.
2. (2) Find the Z-transform of the geometric sequence  $(3, 15, 75, 375, \dots)$ .
3. (2) Find the Z-transform of the sequence  $(6, 12, 24, \dots)$  both by considering it the advance of the sequence  $(3, 6, 12, 24, \dots)$  and by applying the formula for geometrical sequences directly. Do you get the same answer?
4. (2) For the difference equation

$$x_{k+1} = -3x_k \quad (6)$$

with  $x_0 = 1$  work out  $X(z) = \mathcal{Z}[(x_n)]$  by taking the Z-transform of both sides of the equation. Use this to solve the equation.

---

<sup>1</sup>Conor Houghton, [houghton@maths.tcd.ie](mailto:houghton@maths.tcd.ie), see also <http://www.maths.tcd.ie/~houghton/2E2.html>