

2E2 Tutorial Sheet 1¹

16 October 2005

Useful formulae:

- The Laplace transform of $f(t)$:

$$\mathcal{L}(f) = \int_0^\infty f(t)e^{-st}dt \quad (1)$$

- Linearity:

$$\mathcal{L}(af + bg) = a\mathcal{L}(f) + b\mathcal{L}(g) \quad (2)$$

where a and b are constants.

- Integration by parts:

$$\int_a^b u dv = uv]_a^b - \int_a^b u dv \quad (3)$$

- Table of Laplace transforms: $\mathcal{L}(1) = 1/s$,

$$\mathcal{L}(t^n) = \frac{n!}{s^{n+1}} \quad (4)$$

and

$$\mathcal{L}(e^{at}) = \frac{1}{s-a} \quad (5)$$

- The first shift theorem: if $\mathcal{L}[f(t)] = F(s)$ then

$$\mathcal{L}[f(t)e^{at}] = F(s-a) \quad (6)$$

- Laplace transform and differentiation: if $\mathcal{L}[f(t)] = F(s)$ then

$$\mathcal{L}(f') = sF - f(0) \quad (7)$$

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/2E2.html>

Questions

1. (1) Using the linearity of the Laplace transform, calculate the Laplace transform of

$$f(t) = 2 - \frac{t}{2} \quad (8)$$

2. (1) Using the linearity of the Laplace transform, calculate the Laplace transform of

$$f(t) = 2e^{2t} + 3t + 4e^{-4t} \quad (9)$$

3. (2) The hyperbolic sine is defined as

$$\sinh x = \frac{e^x - e^{-x}}{2} \quad (10)$$

using the linearity of the Laplace transform, show that

$$\mathcal{L}(\sinh at) = \frac{a}{s^2 - a^2} \quad (11)$$

4. (2) Using the shift theorem find the Laplace transform of

$$f(t) = e^{2t}t^2 \quad (12)$$

5. (2) Using the formula for the Laplace transform of the differential find $\mathcal{L}(f')$ where $f = t^2$, check your answer by differentiating f directly and then working out its Laplace transform.