

Vegetation Classification for Orange County
Final Report



Prepared by:

AECOM

Prepared for:

Nature Reserve of Orange County

January 2013

Table of Contents

| | |
|--|----|
| INTRODUCTION | 1 |
| METHODS | 1 |
| Sampling Strategy | 1 |
| Rapid Assessment Data Collection | 2 |
| Analysis | 4 |
| RESULTS | 5 |
| DISCUSSION | 6 |
| REFERENCES | 6 |
| VEGETATION DOMINATED BY TREES | 7 |
| Acer macrophyllum Alliance | 7 |
| Alnus rhombifolia Alliance | 8 |
| Hesperocyparis forbesii Alliance | 12 |
| Juglans californica Alliance | 13 |
| Pinus attenuata Alliance | 15 |
| Pinus coulteri Alliance | 17 |
| Platanus racemosa Alliance | 20 |
| Pseudotsuga macrocarpa Alliance | 25 |
| Quercus agrifolia Alliance | 27 |
| Quercus chrysolepis Alliance | 34 |
| Quercus wislizeni Alliance | 36 |
| Salix gooddingii Alliance | 37 |
| Salix laevigata Alliance | 38 |
| VEGETATION DOMINATED BY SHRUBS | 41 |
| Acmispon glaber Alliance | 41 |
| Adenostoma fasciculatum Alliance | 42 |
| Adenostoma fasciculatum-Arctostaphylos glandulosa Alliance | 44 |
| Adenostoma fasciculatum-Arctostaphylos glandulosa-Ceanothus crassifolius Association | 45 |
| Adenostoma fasciculatum-Arctostaphylos glauca Alliance | 46 |
| Adenostoma fasciculatum-Salvia mellifera Alliance | 47 |
| Arctostaphylos glandulosa Alliance | 49 |
| Artemisia californica Alliance | 50 |
| Artemisia californica-Eriogonum fasciculatum Alliance | 56 |
| Artemisia californica-Salvia mellifera Alliance | 57 |
| Atriplex amnicola Alliance | 58 |
| Atriplex lentiformis Alliance | 59 |
| Baccharis pilularis Alliance | 60 |
| Baccharis salicifolia Alliance | 61 |
| Ceanothus crassifolius Alliance | 62 |
| Ceanothus megacarpus Alliance | 63 |

| | |
|--|-----|
| Ceanothus tomentosus Alliance..... | 64 |
| Cercocarpus betuloides Alliance | 66 |
| Encelia californica Alliance | 67 |
| Ericameria palmeri Alliance..... | 69 |
| Eriodictyon crassifolium Alliance..... | 70 |
| Eriogonum fasciculatum Alliance | 71 |
| Isocoma menziesii Alliance..... | 74 |
| Keckiella antirrhinoides Alliance | 75 |
| Lepidospartum squamatum Alliance..... | 76 |
| Lycium californicum Alliance | 81 |
| Malacothamnus fasciculatus Alliance | 82 |
| Malosma laurina Alliance | 83 |
| Opuntia littoralis Alliance | 87 |
| Quercus berberidifolia Alliance | 88 |
| Quercus berberidifolia-Adenostoma fasciculatum Alliance..... | 92 |
| Rhus integrifolia Alliance..... | 94 |
| Salix lasiolepis Alliance | 96 |
| Salvia apiana Alliance | 98 |
| Salvia leucophylla Alliance..... | 99 |
| Salvia mellifera Alliance..... | 100 |
| Sambucus nigra Alliance..... | 104 |
| Toxicodendron diversilobum Alliance | 105 |
| VEGETATION DOMINATED BY HERBS | 107 |
| Avena (barbata, fatua) Alliance | 107 |
| Brassica nigra Alliance | 108 |
| Bromus diandrus Alliance..... | 109 |
| Cynara cardunculus Provisional Alliance..... | 110 |
| Festuca perennis Alliance..... | 111 |
| Frankenia salina Alliance | 113 |
| Mediterranean California Naturalized Annual and Perennial Grassland Alliance | 114 |
| Muhlenbergia rigens Alliance..... | 115 |
| Phalaris aquatica Alliance..... | 117 |
| Schoenoplectus acutus Alliance | 118 |
| Spartina foliosa Alliance | 119 |
| Stipa lepida Provisional Alliance..... | 120 |
| Stipa pulchra Alliance | 122 |
| Typha (angustifolia, domingensis, latifolia) Alliance | 123 |

| | |
|---|---|
| Figure 1 Rapid Assessment Sampling Locations | 3 |
|---|---|



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 1

Introduction

The ultimate goal of this project is to create an updated fine-scale vegetation map for about 58,000 acres of Orange County, consisting of the 37,000-acre Orange County Central and Coastal Subregions Natural Community Conservation Plan (NCCP)/Habitat Conservation Plan (HCP) Habitat Reserve System; approximately 9,500 acres of associated NCCP/HCP Special Linkages, Existing Use Areas, and Non-Reserve Open Space; and approximately 11,000 acres of adjoining conserved open space (study area). To update vegetation mapping, the Natural Reserve of Orange County (NROC) proposes to use Manual of California Vegetation (MCV) methods (2009), which will be implemented in two phases:

Phase 1 – Development of a vegetation classification system for the Central and Coastal Subregions of Orange County that is consistent with the MCV.

Phase 2 – Application of the vegetation classification system to create a vegetation map through photo-interpretation of available aerial imagery and ground reconnaissance.

The MCV methods were developed by the California Department of Fish and Game (CDFG) Vegetation Classification and Mapping Program in collaboration with the California Native Plant Society (CNPS). This approach relies on the collection of quantifiable environmental data to identify and classify biological associations that repeat across the landscape. For areas where documentation is lacking to effectively define all of the vegetation patterns found in California, CDFG and CNPS developed the Vegetation Rapid Assessment Protocol. This protocol guides data collection and analysis to refine vegetation classifications that are consistent with CDFG and MCV standards. Based on an earlier classification by Gray and Bramlet (1992), Orange County is expected to have vegetation types not yet described in the MCV. Using the MCV approach, Rapid Assessment (RA) data was collected throughout the study area and analyzed to characterize these new vegetation types or show concurrence with existing MCV types.

Methods

This project is currently at the end of Phase 1, which consisted of three tasks:

1. Define the sampling strategy.
2. Collect field data in the form of Rapid Assessments.
3. Analyze data, resulting in a vegetation classification system (VCS).

Task 1 was completed in February 2012 with collaboration from NROC, CDFG, and the U.S. Fish and Wildlife Service (USFWS). Task 2 was initiated in March and completed in October 2012. Ed Kentner (AECOM) led the data collection effort with the assistance of Kyle Harper and Jonathan Dunn (AECOM) and Mark Bibbo (Garcia and Associates). Property access and permitting were coordinated by Milan Mitrovich (NROC).

Sampling Strategy

Through discussions with CDFG staff (Todd Keeler-Wolf, Diana Hickson), AECOM achieved concurrence that approximately 330 RAs from the study area would be sufficient to prepare the VCS. This number was determined through consideration of the size of the study area and the number of vegetation types described by Gray and Bramlet (1992), and the expectation that existing datasets collected for similar efforts in San Diego and Riverside Counties would be suitable for inclusion in this current classification effort.

A set of draft 330 sampling points were distributed across the Central and Coastal, and Southern Reserve areas of Orange County using a stratified random sampling method. Sampling locations within the Southern Reserves were included to ensure the inclusion of vegetation types that are unmapped within the Central and Coastal Reserves.

but potentially present therein, provide greater geographical continuity for the inclusion of San Diego and Riverside County data, and yield a VCS of greater utility for the region. As a first step, AECOM created new habitat categories based on the existing Habitat/Land Cover Classification System for Orange County, as derived from Gray and Bramlet (1992). Some existing categories were combined and categorized at higher levels, and others were left at the most descriptive level. For example, black sage scrub, white sage scrub, and buckwheat scrub (among others) were combined into the habitat type "Scrub." However, purple sage scrub and southern cactus scrub were left as individual habitat types. The decision about which habitat types to combine and which habitat types to leave as a separate type was based on habitat types that are unique to Orange County versus habitat types for which data already exists in San Diego and/or Riverside Counties. Habitats unique to Orange County were left as individual types to ensure that they were represented in the sampling. Table 1 details the final categories that AECOM used for stratifying the 330 sampling points.

In addition to habitat type, acreage of each habitat type was considered when stratifying the 330 sampling points. Habitat types with greater acreages received a proportionally larger number of sampling points. Additionally, since the Central and Coastal Reserve lands are the primary study area, preference was given to these lands when a habitat type occurred in both the Central and Coastal Reserves and Southern Reserve. After the number of sampling points was stratified based on habitat type, acreage, and reserve, geographic information systems (GIS) software was used to randomly place the sampling points.

This sampling system was designed to capture adequate vegetation data, in the form of Rapid Assessments, to prepare a vegetation classification for the study area. The stratification method is purposefully objective to ensure that known vegetation types are sampled; however, it would be entirely appropriate to reallocate a portion of the proposed sampling locations or identify additional sampling based on the knowledge obtained from land managers and other local experts.

On February 2, 2012, a meeting was held at the San Diego offices of CDFG to review the sampling strategy and provide comments regarding the draft sampling locations. In attendance were Kristine Preston and Milan Mitrovich (NROC), Will Miller (USFWS), Cara Allen (CDFG), and Jonathan Dunn (AECOM). The overall strategy was accepted as sound, and several suggestions were made for the reallocation of a portion of the sampling locations. Based on these suggestions, adjustments were made to the draft sampling locations. These adjustments included the allocation of addition points within the unchannelized portions of the Santa Ana River, additional points within grassland habitats, additional points in Upper Newport Bay, additional points in the coastal region, representation of the limited natural resource areas of the northern part of the county, and several other minor adjustments. Approximately 150 "alternate" sampling locations were also selected using the same stratified random sampling method described above. These alternate locations were used in cases where the original sampling point could not be accessed or the habitat type had changed and that location no longer represented the habitat type desired. Therefore, a total of 480 primary and alternate sampling locations were identified.

As a practical matter, field ecologists who collect the RA data assessed conditions within the vicinity of a proposed sampling location and had discretion to adjust the sampling location or collect multiple RAs within the vicinity. Therefore, throughout the data collection stage, sampling was tracked to ensure that all ecological types were represented. Table 1 details the number of sampling units in each habitat type and in each reserve.

Rapid Assessment Data Collection

Field data collection was conducted between March 6 and October 10, 2012. A total of 346 RAs were collected throughout the study area (Figure 1). Of this total, 216 RAs were collected in the Central and Coastal Reserves, 127 RAs were collected in the Southern Reserves, and three RAs were collected in the Northern Reserves.




Figure 1
Rapid Assessment Sampling Locations

NROC Vegetation Classification

Path: P:\2012\60242012\06GIS\6.3_Layout\NROC_classification_report_15DEC2012.mxd, 31 October 2012, dunnj

Table 1. Sampling Points Stratified by Habitat Type (1992) and Reserve Area

| Habitat Type | Central & Coastal Reserve | Southern Reserve | Northern Reserve | Total |
|---|---------------------------|------------------|------------------|------------|
| Grassland | 37 | 12 | 2 | 51 |
| Generalized Annual Grassland | 33 | 7 | 2 | 42 |
| Deergrass | | 3 | | 3 |
| Needlegrass | 3 | 2 | | 5 |
| Wild Rye | 1 | | | 1 |
| Scrub | 79 | 25 | 3 | 107 |
| Generalized Scrub | 61 | 22 | 1 | 84 |
| Box Springs Goldenbush-Grassland | 3 | | | 3 |
| Chenopod Scrub or Quail Brush | 2 | | | 2 |
| Purple Sage | 2 | | | 2 |
| Southern Cactus | 5 | 2 | 2 | 9 |
| White Sage | 2 | | | 2 |
| Floodplain Sage Scrub | 2 | 1 | | 3 |
| Maritime Succulent Scrub | 1 | | | 1 |
| Riversidian Coastal Sage Scrub | 1 | | | 1 |
| Chaparral | 57 | 49 | 3 | 109 |
| Generalized Chaparral | 43 | 47 | 3 | 93 |
| Bigpod Ceanothus | 2 | | | 2 |
| Nolina | 4 | | | 4 |
| Snowball Ceanothus | 4 | 2 | | 6 |
| Maritime Chaparral | 4 | | | 4 |
| Woodland | 8 | 1 | 4 | 13 |
| California Walnut Woodland | | | 1 | 1 |
| Coast Live Oak Woodland | 8 | 1 | 3 | 12 |
| Forest | 7 | 4 | | 11 |
| Bigcone Spruce-Canyon Live Oak Forest | 1 | 2 | | 3 |
| Canyon Live Oak Forest | 1 | 1 | | 2 |
| Coast Live Oak Forest | 1 | 1 | | 2 |
| Interior Live Oak Forest | 1 | | | 1 |
| Knobcone Pine Forest | 1 | | | 1 |
| Southern Interior Cypress Forest or Tecate Cypress Forest | 2 | | | 2 |
| Riparian | 14 | 11 | 5 | 30 |
| Generalized Riparian | 5 | 5 | 5 | 15 |
| Canyon Live Oak Ravine Forest | 1 | | | 1 |
| Coast Live Oak Riparian Forest | 6 | 6 | | 12 |
| White Alder Riparian Forest | 2 | | | 2 |
| Wetlands | 3 | | 1 | 4 |
| Generalized Wetlands | 3 | | 1 | 4 |
| Disturbed and Other Types | 1 | | 4 | 5 |
| Uncategorized | 1 | | 4 | 5 |
| Total | 206 | 102 | 22 | 330 |

Analysis

As discussed above, recent classification efforts in San Diego and Riverside Counties produced an available RA dataset consistent with the methods employed for this current effort. For the types of analyses employed in producing a vegetation classification, a larger dataset is generally expected to produce more robust results by

reducing artificial grouping. Therefore, the data from the 346 RAs collected for this project were combined with data from 1,494 RAs collected for the Western Riverside classification effort and the 1,309 RAs collected for the Western San Diego classification effort. These data were prepared for analysis through the following steps:

1. Generate “pseudo-species” to combine similar or ecologically equivalent taxa.
2. Filter out nonnative species in stands dominated by native species.
3. Remove uncommon taxa occurring in less than three stands.
4. Divide the data into “Tree,” “Shrub,” or “Herb” subsets.
5. Normalize data through Arcsine Square Root transformation of cover values.

The resulting datasets were subjected to two stages of an agglomerative hierarchical cluster analysis within a biostatistical software application (PC-ORD). When conducting cluster analyses on large heterogeneous datasets, it is often useful to conduct these analyses in stages that divide the dataset into more manageable groups. In the first stage of analysis, cluster analyses were performed for each of the Tree, Shrub, and Herb subsets. This stage produced subclusters of a manageable size. The RA datasets for each of these subclusters was then reanalyzed to produce the preliminary classification. All stages of the cluster analyses were performed using a Bray-Curtis distance matrix and a flexible linkage of -0.25.

This preliminary classification based strictly on the cluster groupings was then compared to published membership rules. The results of these analyses defined the number of distinct vegetation types sampled, and, when compared to published membership rules, allowed the assignment of each stand to a known alliance and association from the MCV, or, in cases where no known vegetation types agreed with the current analyses, allowed the definition of new or provisional alliances and associations.

Results

The above analysis resulted in the classification of 66 Alliances with 68 Associations and six Semi-Natural types within the study area. Of particular note, these results include two Provisional Alliances and five Provisional Associations defined by native species, and two Semi-Natural Stand types defined by nonnative species that have not been described in previous studies (Table 2). The term “Provisional” is used here to describe a vegetation type that appears valid based on the current analyses but for which a greater number of samples may be required for acceptance in the MCV.

Table 2. Previously Undescribed Vegetation Types Found within the Study Area

| Alliance | Association |
|--|--|
| <i>Artemisia californica</i> | <i>Artemisia californica-Eriogonum fasciculatum-Opuntia littoralis</i> * |
| | <i>Artemisia californica-Mimulus aurantiacus-Elymus condensatus</i> * |
| | <i>Artemisia californica-Nolina cismontana</i> * |
| <i>Atriplex amnicola</i> Semi-Natural Stands | (Semi-Natural Stand Types) |
| <i>Cynara cardunculus</i> Semi-Natural Stands | (Semi-Natural Stand Types) |
| <i>Ericameria palmeri</i> * | <i>Ericameria palmeri/Stipa pulchra</i> * |
| <i>Quercus wislizeni</i> | <i>Quercus wislizeni-Arctostaphylos glandulosa</i> * |
| <i>Rhus integrifolia/Dudleya edulis-Carpobrotus</i> spp. | (Alliance level only) |

* Provisional

The classification results were subdivided into the dominant life form(s) and are presented in the following sections of this report. Data that describe the species composition and environmental conditions recorded within the stands for each vegetation type are summarized in tabular form. Data from the current study (OC stands) are presented alongside the aggregate dataset for Western San Diego (WSD) and Western Riverside (WR). Measured parameters presented in the Local Vegetation Description and Local Environmental Description, such as “cover” and “aspect,” are defined according to the common ecological usage of these terms.

The Stand Tables present calculated values of species frequency and cover under abbreviated headings that require some explanation. “CON” represents the frequency (or constancy) at which a species was detected within all of the stands of the vegetation type represented by the table. The values for AVE, MIN, and MAX represent the average, minimum, and maximum cover values for each species detected within all of the stands for the vegetation type represented by the table.

Discussion

The next phase of this project involves the creation of a new vegetation map for the study area. A mapping classification scheme will need to be defined that will include this vegetation classification and classifications for features not defined by vegetation. CDFG Vegetation Classification and Mapping Program staff provided an initial review of these results. The program currently has limited staff availability to review this vegetation classification in detail for consistency with the MCV, but a more detailed review is projected for completion by March 2013. Therefore, in the course of preparing the mapping classification, it may be determined that minor modifications to the vegetation classification may be required.

The second phase of this project will result in the following products:

- A vegetation map for the approximately 60,000-acre study area.
- A field manual of vegetation types for the Central and Coastal Reserves (including a key to distinguish types).
- A crosswalk and discussion on how to compare the current classification scheme to prior classifications.
- A discussion of mapping methods.

References

Sawyer, J., T. Keeler-Wolf and J. Evens. 2009. *A Manual of California Vegetation, 2nd Edition*. California Native Plant Society Press. Sacramento, California.

Gray, J. and D. Bramlet. 1992. Habitat Classification System Natural Resources Geographic Information Systems (GIS) Project. Report prepared for Orange County Environmental Management Agency, Santa Ana, California.

VEGETATION DOMINATED BY TREES

Acer macrophyllum Alliance

Acer macrophyllum Association

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1230JD01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 62.0 (62.0) | 62-62 (62-62) | - |
| Herb cover (%) | 1.0 (-) | 1-1 (-) | 0-0.5 (-) |
| Shrub cover (%) | 1.0 (-) | 1-1 (-) | 1-2 (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 55.0 (-) | 55-55 (-) | 10-15 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: E (1,1)

Macrotopography: middle 1/3 of slope (1,1)

Microtopography: concave (1,1)

Parent material:

Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 4308 (4308) | 4308-4308 (4308-4308) |
| Slope (degrees) | 15 (15.0) | 15-15 (15-15) |
| Large rock cover (%) | 5.0 (5.0) | 5-5 (5-5) |
| Small rock cover (%) | 41.0 (41.0) | 41-41 (41-41) |
| Bare ground cover (%) | 18.0 (18.0) | 18-18 (18-18) |
| Litter cover (%) | 34.0 (34.0) | 34-34 (34-34) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-------------------------------|-----------|-----------|-----------|-----------|
| Tree | <i>Acer macrophyllum</i> | 100 (100) | 34 (34) | 34 (34) | 34 (34) |
| | <i>Platanus racemosa</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Pseudotsuga macrocarpa</i> | 100 (100) | 6 (6) | 6 (6) | 6 (6) |
| | <i>Quercus chrysolepis</i> | 100 (100) | 18 (18) | 18 (18) | 18 (18) |
| Shrub | <i>Ribes</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | <i>Dryopteris arguta</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Polystichum imbricans</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Alnus rhombifolia* Alliance**
***Alnus rhombifolia* (Alliance Only)**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: IDYL.056/BPU 379-33, IDYL.061, LAFU.029/1330

Rapid Assessment(s), OC stands (n=1): 1195EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|-------------|
| Total vegetation cover (%) | 70.0 (56.0) | 56-94 (56-56) | - |
| Herb cover (%) | 27.8 (26.0) | 15-50 (26-26) | 0-1 (0-0.5) |
| Shrub cover (%) | 8.3 (0.2) | 0.2-15 (0.2-0.2) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0-5 (0-0.5) |
| Overstory tree cover (%) | 48.3 (39.0) | 39-65 (39-39) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (2,0), NW (1,1), SE (1,0)

Macrotopography: lower 1/3 of slope (1,0), bottom (3,1)

Microtopography: concave (4,1)

Parent material: mixed alluvium (2,1), sandy alluvium (2,0)

Soil texture: sandy loam (1,0), silt (1,0), sand (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 4096 (1512) | 1512-5271 (1512-1512) |
| Slope (degrees) | 3 (2.0) | 2-5 (2-2) |
| Large rock cover (%) | 10.5 (1.0) | 1-20 (1-1) |
| Small rock cover (%) | 11.5 (24.0) | 5-24 (24-24) |
| Bare ground cover (%) | 9.8 (16.0) | 2-16 (16-16) |
| Litter cover (%) | 57.3 (54.0) | 40-75 (54-54) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---------------------------------|-----------|-----------|-----------|-----------|
| Tree | <i>Alnus rhombifolia</i> | 100 (100) | 39 (38) | 20 (38) | 60 (38) |
| | <i>Calocedrus decurrens</i> | 50 (-) | 12 (-) | 7 (-) | 17 (-) |
| | <i>Pinus coulteri</i> | 50 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| | <i>Pinus jeffreyi</i> | 50 (-) | 2.5 (-) | 1 (-) | 4 (-) |
| | <i>Pinus lambertiana</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Platanus racemosa</i> | 25 (100) | 1.2 (1.2) | 1.2 (1.2) | 1.2 (1.2) |
| | <i>Quercus chrysolepis</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| | <i>Quercus kelloggii</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| | <i>Quercus wislizeni</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| Shrub | <i>Rosa californica</i> | 50 (-) | 8.5 (-) | 2 (-) | 15 (-) |
| | <i>Euonymus occidentalis</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | * <i>Ficus carica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Lonicera subspicata</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Rhamnus californica</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Rhamnus ilicifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Rhododendron occidentale</i> | 25 (-) | 12 (-) | 12 (-) | 12 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 9

| Species Name | Con | Ave | Min | Max |
|---|----------|-----------|-----------|-----------|
| <i>Ribes nevadense</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Salix lasiolepis</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Toxicodendron diversilobum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>Urtica dioica</i> | 75 (-) | 9.7 (-) | 1 (-) | 25 (-) |
| <i>Carex</i> | 50 (-) | 3 (-) | 1 (-) | 5 (-) |
| <i>Claytonia parviflora</i> | 50 (-) | 2 (-) | 1 (-) | 3 (-) |
| <i>Fragaria vesca</i> | 50 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Pteridium aquilinum</i> var. <i>pubescens</i> | 50 (-) | 2.5 (-) | 1 (-) | 4 (-) |
| <i>Achillea millefolium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Agrostis exarata</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Artemisia douglasiana</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| * <i>Arundo donax</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Athyrium filix-femina</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Bromus diandrus</i> | 25 (-) | 12 (-) | 12 (-) | 12 (-) |
| * <i>Bromus tectorum</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| * <i>Epilobium</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Equisetum arvense</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Juncus effusus</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Juncus</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| <i>Lilium humboldtii</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Nasturtium officinale</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| * <i>Plantago major</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Potentilla glandulosa</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Rumex crispus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Scirpus microcarpus</i> | 25 (-) | 4 (-) | 4 (-) | 4 (-) |
| <i>Stachys</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stachys albens</i> | 25 (-) | 7 (-) | 7 (-) | 7 (-) |
| * <i>Stipa miliacea</i> var. <i>miliacea</i> | 25 (100) | 10 (10) | 10 (10) | 10 (10) |
| * <i>Veronica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Vinca major</i> | 25 (100) | 14 (14) | 14 (14) | 14 (14) |



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 10

***Alnus rhombifolia* Alliance**

***Alnus rhombifolia*-*Acer macrophyllum* Association**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1283EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 50.0 (50.0) | 50-50 (50-50) | - |
| Herb cover (%) | 4.0 (4.0) | 4-4 (4-4) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 15.0 (15.0) | 15-15 (15-15) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | 4.0 (4.0) | 4-4 (4-4) | 1-2 (1-2) |
| Overstory tree cover (%) | 40.0 (40.0) | 40-40 (40-40) | 10-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1)

Macrotopography: middle 1/3 of slope (1,1)

Microtopography: concave (1,1)

Parent material: mixed alluvium (1,1)

Soil texture: sand (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3009 (3009) | 3009-3009 (3009-3009) |
| Slope (degrees) | 6 (6.0) | 6-6 (6-6) |
| Large rock cover (%) | 10.0 (10.0) | 10-10 (10-10) |
| Small rock cover (%) | 34.0 (34.0) | 34-34 (34-34) |
| Bare ground cover (%) | 16.0 (16.0) | 16-16 (16-16) |
| Litter cover (%) | 35.0 (35.0) | 35-35 (35-35) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | |
| <i>Acer macrophyllum</i> | 100 (100) | 8 (8) | 8 (8) | 8 (8) |
| <i>Alnus rhombifolia</i> | 100 (100) | 17 (17) | 17 (17) | 17 (17) |
| <i>Pseudotsuga macrocarpa</i> | 100 (100) | 5 (5) | 5 (5) | 5 (5) |
| <i>Umbellularia californica</i> | 100 (100) | 14 (14) | 14 (14) | 14 (14) |
| Shrub | | | | |
| <i>Frangula californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Ribes californicum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Rubus ursinus</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| <i>Symphoricarpos mollis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Toxicodendron diversilobum</i> | 100 (100) | 12 (12) | 12 (12) | 12 (12) |
| Herb | | | | |
| <i>Artemisia douglasiana</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Dryopteris arguta</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Elymus triticoides</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Galium aparine</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lilium humboldtii</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Melica imperfecta</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Osmorhiza brachypoda</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|----------------------------|-----------|-----------|-----------|-----------|
| <i>Stachys rigida</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Thalictrum fendleri</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Alnus rhombifolia* Alliance**

***Alnus rhombifolia*-*Platanus racemosa* Association**

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: BLACK.031\1382-BPU1216-293

Rapid Assessment(s), OC stands (n=2): 1212EK01, 1526EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|---------------|
| Total vegetation cover (%) | 70.0 (67.5) | 67-75 (67-68) | - |
| Herb cover (%) | 10.0 (12.0) | 4-20 (4-20) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 20.1 (24.1) | 0.2-48 (0.2-48) | 0-2 (0-2) |
| Tall shrub / low tree cover (%) | 14.0 (14.0) | 14-14 (14-14) | 2-5 (2-5) |
| Overstory tree cover (%) | 54.3 (49.0) | 49-65 (49-49) | 10-20 (10-20) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), SW (1,0), W (1,1)

Macrotopography: bottom (2,2), lower 1/3 of slope (1,0)

Microtopography: concave (3,2)

Parent material: granitic (1,0), mixed alluvium (2,2)

Soil texture: sand (2,1)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1969 (1450) | 823-3007 (823-2077) |
| Slope (degrees) | 3 (2.5) | 1-4 (1-4) |
| Large rock cover (%) | 12.0 (0.5) | 0-35 (0-1) |
| Small rock cover (%) | 18.0 (23.5) | 7-25 (22-25) |
| Bare ground cover (%) | 3.3 (5.0) | 0-8 (2-8) |
| Litter cover (%) | 56.7 (62.5) | 45-65 (60-65) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | |
| <i>Alnus rhombifolia</i> | 100 (100) | 31 (36.5) | 20 (28) | 45 (45) |
| <i>Platanus racemosa</i> | 100 (100) | 12 (6.5) | 4 (4) | 23 (9) |
| <i>Salix laevigata</i> | 67 (50) | 19.5 (26) | 13 (26) | 26 (26) |
| <i>Acer macrophyllum</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Populus fremontii</i> | 33 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Quercus agrifolia</i> | 33 (-) | 10 (-) | 10 (-) | 10 (-) |
| <i>Umbellularia californica</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | |
| <i>Toxicodendron diversilobum</i> | 67 (50) | 4 (2) | 2 (2) | 6 (2) |
| <i>Acmispon glaber</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Baccharis salicifolia</i> | 33 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Frangula californica</i> | 33 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>Heteromeles arbutifolia</i> | 33 (-) | 5 (-) | 5 (-) | 5 (-) |

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| <i>Keckiella cordifolia</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Phoradendron</i> | 33 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Prunus ilicifolia</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Rhamnus ilicifolia</i> | 33 (-) | 2 (-) | 2 (-) | 2 (-) |
| * <i>Ricinus communis</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Rubus ursinus</i> | 33 (50) | 45 (45) | 45 (45) | 45 (45) |
| Herb | | | | |
| <i>Artemisia douglasiana</i> | 100 (100) | 2.4 (3.5) | 0.2 (3) | 4 (4) |
| <i>Stachys</i> | 67 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Urtica dioica</i> | 67 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus diandrus</i> | 33 (-) | 3 (-) | 3 (-) | 3 (-) |
| * <i>Cyperus</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Galium aparine</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Juncus oxymeris</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lupinus latifolius</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Mimulus cardinalis</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Mimulus guttatus</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Nasturtium officinale</i> | 33 (50) | 5 (5) | 5 (5) | 5 (5) |
| * <i>Oxalis pes-caprae</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Scrophularia californica</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Stipa miliacea</i> var. <i>miliacea</i> | 33 (50) | 11 (11) | 11 (11) | 11 (11) |
| <i>Xanthium strumarium</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Hesperocyparis forbesii* Alliance**

***Hesperocyparis forbesii* Provisional Association**

Samples Used To Describe Association (n= 9)

Rapid Assessment(s), WR and WSD stands: 1834MB02, 1600LS01, 1834MB01, 1361LS02, 1834MB04, 1401MB01, 1839MB01

Rapid Assessment(s), OC stands (n=2): 1275JD01, 1506EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 63.0 (50.5) | 41-87 (41-60) | - |
| Herb cover (%) | 7.8 (2.0) | 1-22 (2-2) | 0-1 (0-0.5) |
| Shrub cover (%) | 41.0 (41.0) | 41-41 (41-41) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 36.0 (-) | 5-60 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NE (2,1), Flat (2,0), NW (4,0)

Macrotopography: bottom (2,0), lower 1/3 of slope (2,0), ridge top (1,0), upper 1/3 of slope (4,2)

Microtopography: convex (2,0), flat (5,1), concave (2,1)

Parent material: metamorphic (5,0), sandstone (1,1)

Soil texture: clay loam (6,0), sand (1,1), sandy loam (1,1), unknown (1,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 13

| | Average | Range |
|-----------------------|-------------|----------------------|
| Elevation (ft) | 1727 (3066) | 249-4491 (1640-4491) |
| Slope (degrees) | 9 (13.5) | 0-25 (12-15) |
| Large rock cover (%) | 5.0 (-) | 0-25 (-) |
| Small rock cover (%) | 32.8 (10.0) | 10-75 (10-10) |
| Bare ground cover (%) | 22.1 (63.0) | 7-63 (63-63) |
| Litter cover (%) | 34.8 (25.0) | 0-67 (25-25) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|-----------|------------|-----------|----------|
| Tree | <i>Hesperocyparis forbesii</i> | 100 (100) | 30.4 (2.6) | .2 (.2) | 60 (5) |
| | | | | | |
| Shrub | <i>Malosma laurina</i> | 78 (50) | 3.4 (0.2) | 0.2 (0.2) | 12 (0.2) |
| | <i>Adenostoma fasciculatum</i> | 67 (50) | 9.2 (0.2) | 0.2 (0.2) | 20 (0.2) |
| | <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 56 (-) | 1.5 (-) | 0.2 (-) | 4 (-) |
| | <i>Ceanothus tomentosus</i> | 44 (50) | 18.3 (24) | 15 (24) | 24 (24) |
| | <i>Arctostaphylos glandulosa</i> subsp. <i>glandulosa</i> | 33 (-) | 8.7 (-) | 7 (-) | 12 (-) |
| | <i>Chamaebatia australis</i> | 33 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Dendromecon rigida</i> | 33 (50) | 1.1 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 33 (50) | 1.4 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Mimulus aurantiacus</i> | 33 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| | | | | | |
| | | | | | |
| Herb | <i>*Festuca myuros</i> | 78 (50) | 1.7 (0.2) | 0.2 (0.2) | 4 (0.2) |
| | <i>*Bromus rubens</i> | 67 (-) | 3.2 (-) | 0.2 (-) | 12 (-) |
| | <i>*Centaurea melitensis</i> | 56 (50) | 1.9 (0.2) | 0.2 (0.2) | 6 (0.2) |
| | <i>*Hirschfeldia incana</i> | 44 (50) | 0.8 (1) | 0.2 (1) | 1 (1) |
| | <i>*Avena barbata</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Galium porrigens</i> var. <i>porrigens</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Juglans californica* Alliance**

***Juglans californica*/Bromus diandrus Association**

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=3): 1444EK01, 1444EK02, 1490EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 58.7 (58.7) | 33-80 (33-80) | - |
| Herb cover (%) | 50.7 (50.7) | 25-75 (25-75) | 0-2 (0-2) |
| Shrub cover (%) | 2.7 (2.7) | 1-5 (1-5) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 7.3 (7.3) | 7-8 (7-8) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 14

Local Environmental Description (All stands, OC stands)

Aspect: E (1,1), N (1,1), SE (1,1)

Macrotopography: middle 1/3 of slope (2,2), middle to upper 1/3 of slope (1,1)

Microtopography: undulating (3,3)

Parent material: sandstone (1,1)

Soil texture: clay (1,1), clay loam (2,2)

| | Average | Range |
|------------------------------|-------------|-------------------|
| Elevation (ft) | 595 (595) | 456-751 (456-751) |
| Slope (degrees) | 23 (22.7) | 15-35 (15-35) |
| Large rock cover (%) | 0.3 (0.3) | 0-1 (0-1) |
| Small rock cover (%) | 1.0 (1.0) | 0-3 (0-3) |
| Bare ground cover (%) | 42.3 (42.3) | 32-52 (32-52) |
| Litter cover (%) | 53.3 (53.3) | 45-65 (45-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--|-----------|-------------|-----------|-----------|
| Tree | <i>Juglans californica</i> | 100 (100) | 6.7 (6.7) | 6 (6) | 7 (7) |
| | * <i>Schinus molle</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Nicotiana glauca</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Quercus agrifolia</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | <i>Heteromeles arbutifolia</i> | 67 (67) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>Malosma laurina</i> | 67 (67) | 2.5 (2.5) | 1 (1) | 4 (4) |
| | <i>Rhus integrifolia</i> | 67 (67) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>Sambucus nigra</i> subsp. <i>caerulea</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Artemisia californica</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Marah</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Opuntia littoralis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Ricinus communis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Salvia apiana</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Salvia leucophylla</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | <i>Toxicodendron diversilobum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Bromus diandrus</i> | 100 (100) | 35 (35) | 25 (25) | 40 (40) |
| | * <i>Centaurea melitensis</i> | 100 (100) | 1.8 (1.8) | 0.2 (0.2) | 5 (5) |
| | <i>Acourtia microcephala</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Brassica nigra</i> | 67 (67) | 22.5 (22.5) | 15 (15) | 30 (30) |
| | <i>Elymus condensatus</i> | 67 (67) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | * <i>Silybum marianum</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Calystegia macrostegia</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Chaenactis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Croton setiger</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Foeniculum vulgare</i> | 33 (33) | 5 (5) | 5 (5) | 5 (5) |
| | * <i>Hirschfeldia incana</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Malacothrix saxatilis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Phacelia</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|--|---------|-----------|-----------|-----------|
| <i>Pseudognaphalium californicum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Solanum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Sonchus asper</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Stephanomeria</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Stipa miliacea</i> var. <i>miliacea</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Pinus attenuata* Alliance**

***Pinus attenuata* (Alliance Only)**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1321EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 62.0 (62.0) | 62-62 (62-62) | - |
| Herb cover (%) | 26.0 (26.0) | 26-26 (26-26) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 2.0 (2.0) | 2-2 (2-2) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 40.0 (40.0) | 40-40 (40-40) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1)

Macrotopography:

Microtopography: undulating (1,1)

Parent material:

Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3645 (3645) | 3645-3645 (3645-3645) |
| Slope (degrees) | 4 (4.0) | 4-4 (4-4) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 5.0 (5.0) | 5-5 (5-5) |
| Bare ground cover (%) | 27.0 (27.0) | 27-27 (27-27) |
| Litter cover (%) | 65.0 (65.0) | 65-65 (65-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------|-----------|-----------|-----------|-----------|
| Tree | <i>Pinus attenuata</i> | 100 (100) | 35 (35) | 35 (35) | 35 (35) |
| | <i>Pinus coulteri</i> | 100 (100) | 5 (5) | 5 (5) | 5 (5) |
| | | | | | |
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Helianthemum scoparium</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Quercus berberidifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | | | | | |
| Herb | * <i>Bromus diandrus</i> | 100 (100) | 10 (10) | 10 (10) | 10 (10) |
| | * <i>Centaurea melitensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-----------|-----------|-----------|
| <i>Corethrogyne filaginifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Festuca myuros</i> | 100 (100) | 15 (15) | 15 (15) | 15 (15) |
| * <i>Marrubium vulgare</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Pinus attenuata* Alliance**

***Pinus attenuata*/Arctostaphylos glandulosa Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: CLNF.125/310, CLNF.110, CLNF.114

Rapid Assessment(s), OC stands (n=1): 1342EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 61.5 (45.0) | 45-69 (45-45) | - |
| Herb cover (%) | 11.3 (7.0) | 0-38 (7-7) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 28.3 (9.0) | 4-55 (9-9) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-2 (-) |
| Overstory tree cover (%) | 26.5 (29.0) | 20-35 (29-29) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NE (2,0), Variable (1,0)

Macrotopography: middle 1/3 of slope (2,0), ridge top (1,0), upper 1/3 of slope (1,1)

Microtopography: undulating (4,1)

Parent material: sedimentary (3,0)

Soil texture: clay loam (1,0), loam (2,0), sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3176 (3661) | 1657-3744 (3661-3661) |
| Slope (degrees) | 19 (4.0) | 4-35 (4-4) |
| Large rock cover (%) | 0.3 (-) | 0-1 (-) |
| Small rock cover (%) | 12.8 (19.0) | 0-20 (19-19) |
| Bare ground cover (%) | 15.0 (26.0) | 3-26 (26-26) |
| Litter cover (%) | 57.5 (55.0) | 20-85 (55-55) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-----------|-----------|-----------|
| Tree | <i>Pinus attenuata</i> | 100 (100) | 21 (25) | 17 (25) | 25 (25) |
| | <i>Pinus coulteri</i> | 50 (100) | 11.5 (4) | 4 (4) | 19 (4) |
| Shrub | <i>Arctostaphylos glandulosa</i> | 100 (100) | 23 (6) | 2 (6) | 54 (6) |
| | <i>Adenostoma fasciculatum</i> | 75 (100) | 5.1 (2) | 0.2 (2) | 13 (2) |
| | <i>Dendromecon rigida</i> | 75 (100) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Quercus berberidifolia</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Ceanothus papillosus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Ceanothus tomentosus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Garrya flavescens</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 17

| Species Name | Con | Ave | Min | Max |
|--------------------------------|----------|-----------|-----------|-----------|
| <i>Helianthemum scoparium</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Heteromeles arbutifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Mimulus aurantiacus</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Rhamnus ilicifolia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Brassica nigra</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Bromus hordeaceus</i> | 25 (-) | 15 (-) | 15 (-) | 15 (-) |
| * <i>Bromus madritensis</i> | 25 (100) | 2 (2) | 2 (2) | 2 (2) |
| * <i>Bromus tectorum</i> | 25 (100) | 5 (5) | 5 (5) | 5 (5) |
| * <i>Festuca myuros</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Penstemon spectabilis</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Pinus coulteri* Alliance**

***Pinus coulteri* (Alliance Only)**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1172EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 35.0 (35.0) | 35-35 (35-35) | - |
| Herb cover (%) | 6.0 (6.0) | 6-6 (6-6) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 1.0 (1.0) | 1-1 (1-1) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 34.0 (34.0) | 34-34 (34-34) | 10-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1)

Macrotopography: upper 1/3 of slope (1,1)

Microtopography: convex (1,1)

Parent material: granitic (1,1)

Soil texture: clay loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3451 (3451) | 3451-3451 (3451-3451) |
| Slope (degrees) | 5 (5.0) | 5-5 (5-5) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 24.0 (24.0) | 24-24 (24-24) |
| Litter cover (%) | 74.0 (74.0) | 74-74 (74-74) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--------------------------|-----------|---------|---------|---------|
| Tree | | | | |
| <i>Pinus coulteri</i> | 100 (100) | 33 (33) | 33 (33) | 33 (33) |
| <i>Quercus agrifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |

| | Species Name | Con | Ave | Min | Max |
|--------------|--|-----------|-----------|-----------|-----------|
| Shrub | <i>Cercocarpus betuloides</i> var. <i>betuloides</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhamnus ilicifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | | | | | |
| Herb | <i>Drymocallis lactea</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Dryopteris arguta</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Elymus glaucus</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Galium porrigens</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Osmorhiza brachypoda</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Paeonia californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Sidalcea malviflora</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Viola purpurea</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |

***Pinus coulteri* Alliance**

***Pinus coulteri*/Arctostaphylos glandulosa-Quercus wislizeni Association**

Samples Used To Describe Association (n= 7)

Rapid Assessment(s), WR and WSD stands: BLACK.035/65, CLNF.081, CLNF.094, CLNF.032, CLNF.063

Rapid Assessment(s), OC stands (n=2): 1210EK02, 1213JD01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 72.7 (57.0) | 48-93 (48-66) | - |
| Herb cover (%) | 8.4 (-) | 0-40 (-) | 0-0.5 (-) |
| Shrub cover (%) | 46.2 (-) | 38-62 (-) | 0.5-2 (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 37.2 (-) | 13-47 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (4,1), SW (1,0), Variable (2,1)

Macrotopography: middle 1/3 of slope (1,0), upper 1/3 of slope (4,0), upper 1/3 of slope to ridgetop (2,2)

Microtopography: concave (1,1), Convex (3,0), Undulating (3,1)

Parent material: granitic (5,0)

Soil texture: sand (1,1), sandy loam (5,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3821 (4011) | 3500-4488 (3914-4108) |
| Slope (degrees) | 25 (20.5) | 10-42 (15-26) |
| Large rock cover (%) | 1.7 (-) | 0-8 (-) |
| Small rock cover (%) | 7.1 (3.5) | 0-40 (2-5) |
| Bare ground cover (%) | 8.9 (19.5) | 0-31 (8-31) |
| Litter cover (%) | 63.9 (75.0) | 0-96 (65-85) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-------------|-----------|---------|
| Tree | <i>Pinus coulteri</i> | 100 (100) | 28.6 (13.5) | 10 (12) | 45 (15) |
| | <i>Quercus wislizeni</i> | 86 (100) | 6.8 (6) | 4 (4) | 10 (8) |
| | <i>Quercus chrysolepis</i> | 43 (-) | 4 (-) | 1 (-) | 8 (-) |
| | | | | | |
| Shrub | <i>Arctostaphylos glandulosa</i> | 100 (100) | 29.1 (35) | 10 (32) | 40 (38) |
| | <i>Ceanothus leucodermis</i> | 71 (100) | 5 (2) | 1 (1) | 15 (3) |
| | <i>Adenostoma fasciculatum</i> | 57 (100) | 0.9 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Dendromecon rigida</i> | 43 (50) | 1.1 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 43 (-) | 5 (-) | 1 (-) | 11 (-) |
| | <i>Ceanothus crassifolius</i> | 29 (50) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>Cercocarpus betuloides</i> | 29 (-) | 9 (-) | 1 (-) | 17 (-) |
| | <i>Eriodictyon crassifolium</i> | 29 (50) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 29 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Rhamnus californica</i> | 29 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Rhus ovata</i> | 29 (-) | 1 (-) | 1 (-) | 1 (-) |
| | | | | | |
| Herb | <i>*Bromus tectorum</i> | 29 (-) | 10.1 (-) | 0.2 (-) | 20 (-) |
| | | | | | |

Pinus coulteri Alliance

Pinus coulteri/*Quercus wislizeni* Association

Samples Used To Describe Association (n= 5)

Rapid Assessment(s), WR and WSD stands: BUTT.008/128, PALM.008/553, IDYL.007, BUTT.004/138

Rapid Assessment(s), OC stands (n=1): 1184EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 48.4 (53.0) | 14-60 (53-53) | - |
| Herb cover (%) | 11.2 (1.0) | 1-30 (1-1) | 0-1 (0-0.5) |
| Shrub cover (%) | 26.0 (41.0) | 9-41 (41-41) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 19.4 (27.0) | 4-36 (27-27) | 10-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), SW (3,0), Variable (1,0)

Macrotopography: upper 1/3 of slope (1,1), lower 1/3 of slope (2,0), middle 1/3 of slope (2,0)

Microtopography: Concave (3,1), Undulating (2,0)

Parent material: granitic (4,0), metamorphic (1,1)

Soil texture: clay loam (1,1), sandy loam (4,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 20

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 4922 (4157) | 4157-5476 (4157-4157) |
| Slope (degrees) | 7 (7.0) | 3-14 (7-7) |
| Large rock cover (%) | 27.0 (-) | 0-93 (-) |
| Small rock cover (%) | 9.8 (-) | 0-23 (-) |
| Bare ground cover (%) | 19.0 (17.0) | 1-50 (17-17) |
| Litter cover (%) | 39.4 (80.0) | 2-80 (80-80) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------|-----------|-----------|-----------|---------|
| Tree | <i>Pinus coulteri</i> | 100 (100) | 12.6 (25) | 4 (25) | 25 (25) |
| | <i>Quercus wislizeni</i> | 100 (100) | 13 (8) | 3 (8) | 26 (8) |
| | <i>Calocedrus decurrens</i> | 40 (100) | 1.1 (2) | 0.2 (2) | 2 (2) |
| | <i>Quercus chrysolepis</i> | 40 (-) | 3 (-) | 2 (-) | 4 (-) |
| | | | | | |
| Shrub | <i>Adenostoma sparsifolium</i> | 60 (-) | 2 (-) | 1 (-) | 3 (-) |
| | <i>Cercocarpus betuloides</i> | 60 (-) | 0.7 (-) | 0.2 (-) | 1 (-) |
| | <i>Rhamnus ilicifolia</i> | 60 (100) | 0.7 (1) | 0.2 (1) | 1 (1) |
| | <i>Artemisia tridentata</i> | 40 (-) | 6 (-) | 4 (-) | 8 (-) |
| | <i>Eriogonum fasciculatum</i> | 40 (-) | 2.6 (-) | 0.2 (-) | 5 (-) |
| | <i>Lonicera subspicata</i> | 40 (100) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Salix lasiolepis</i> | 40 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| | | | | | |
| Herb | <i>*Bromus tectorum</i> | 40 (-) | 10.5 (-) | 2 (-) | 19 (-) |
| | | | | | |

Platanus racemosa Alliance

Platanus racemosa Association

Samples Used To Describe Association (n= 10)

Rapid Assessment(s), WR and WSD stands: SUNN.003/482, NOPE.007/1111, BOSP.004/727, LAEL.006, BLACK.052, SUNN.004, LAMA.036, RIVW.004/400, WILD.009/392

Rapid Assessment(s), OC stands (n=1): 1108MA01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 71.5 (65.0) | 50-85 (65-65) | - |
| Herb cover (%) | 23.1 (35.0) | 0-65 (35-35) | 0-1 (0-0.5) |
| Shrub cover (%) | 23.1 (1.0) | 1-75 (1-1) | 0.5-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-5 (-) |
| Overstory tree cover (%) | 36.7 (55.0) | 14-60 (55-55) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (2,1), NW (3,0), SW (5,0)

Macrotopography: bottom (4,1), lower 1/3 of slope (4,0), middle 1/3 of slope (1,0), upper 1/3 of slope (1,0)

Microtopography: Concave (8,0), Flat (2,1)

Parent material: granitic (2,0), mixed alluvium (3,0), mixed igneous (1,0), sandy alluvium (2,0), sedimentary (1,1)

Soil texture: clay loam (2,0), sand (3,0), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 1604 (367) | 367-3166 (367-367) |
| Slope (degrees) | 5 (0.0) | 0-15 (0-0) |
| Large rock cover (%) | 7.5 (-) | 0-20 (-) |
| Small rock cover (%) | 13.4 (-) | 0-47 (-) |
| Bare ground cover (%) | 10.9 (2.0) | 0-30 (2-2) |
| Litter cover (%) | 49.3 (98.0) | 0-100 (98-98) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---------------------------------------|-----------|-----------|---------|---------|
| Tree | | | | | |
| | <i>Platanus racemosa</i> | 100 (100) | 32.9 (55) | 13 (55) | 58 (55) |
| Shrub | | | | | |
| | <i>Salix lasiolepis</i> | 60 (-) | 4.2 (-) | 0.2 (-) | 10 (-) |
| | <i>Toxicodendron diversilobum</i> | 60 (-) | 10.8 (-) | 1 (-) | 25 (-) |
| | <i>Baccharis salicifolia</i> | 50 (-) | 4.4 (-) | 1 (-) | 7 (-) |
| | <i>Sambucus mexicana</i> | 40 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Sambucus nigra subsp. caerulea</i> | 40 (100) | 1.1 (1) | 0.2 (1) | 2 (1) |
| | <i>Artemisia californica</i> | 30 (-) | 1.7 (-) | 1 (-) | 2 (-) |
| | <i>Eriogonum fasciculatum</i> | 30 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| Herb | | | | | |
| | * <i>Bromus diandrus</i> | 60 (100) | 14.8 (35) | 4 (35) | 35 (35) |
| | * <i>Bromus madritensis</i> | 40 (-) | 11.5 (-) | 7 (-) | 23 (-) |

Platanus racemosa Alliance

Platanus racemosa/Baccharis salicifolia Association

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: SDRP0199, 0610TK02, 0506RB05

Rapid Assessment(s), OC stands (n=1): 1293EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 50.8 (29.0) | 29-65 (29-29) | - |
| Herb cover (%) | 25.3 (3.0) | 3-50 (3-3) | 0-2 (0-0.5) |
| Shrub cover (%) | 16.0 (16.0) | 16-16 (16-16) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 21.0 (10.0) | 9-50 (10-10) | 5-15 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,0), SE (1,1), SW (2,0)

Macrotopography: bottom (3,0), middle 1/3 of slope (1,1)

Microtopography: convex (1,0), flat (2,0), concave (1,1)

Parent material: metamorphic (1,1), sandy alluvium (2,0)

Soil texture: sandy loam (1,0), sand (3,1)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 22

| | Average | Range |
|-----------------------|-------------|----------------------|
| Elevation (ft) | 1267 (2172) | 325-2172 (2172-2172) |
| Slope (degrees) | 2 (5.0) | 1-5 (5-5) |
| Large rock cover (%) | 3.0 (-) | 0-12 (-) |
| Small rock cover (%) | 19.0 (65.0) | 0-65 (65-65) |
| Bare ground cover (%) | 8.8 (13.0) | 0-14 (13-13) |
| Litter cover (%) | 41.0 (20.0) | 0-75 (20-20) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-----------------------------------|-----------|------------|-----------|-----------|
| Tree | <i>Platanus racemosa</i> | 100 (100) | 11.5 (10) | 9 (10) | 15 (10) |
| | <i>Salix lasiolepis</i> | 50 (100) | 5.5 (1) | 1 (1) | 10 (1) |
| | * <i>Eucalyptus globulus</i> | 25 (-) | 40 (-) | 40 (-) | 40 (-) |
| | | | | | |
| Shrub | <i>Baccharis salicifolia</i> | 100 (100) | 6.5 (2) | 2 (2) | 12 (2) |
| | <i>Frangula californica</i> | 50 (100) | 5.5 (5) | 5 (5) | 6 (5) |
| | <i>Heteromeles arbutifolia</i> | 50 (100) | 2.5 (1) | 1 (1) | 4 (1) |
| | <i>Keckiella cordifolia</i> | 50 (100) | 5.1 (0.2) | 0.2 (0.2) | 10 (0.2) |
| | <i>Mimulus aurantiacus</i> | 50 (100) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Prunus ilicifolia</i> | 50 (100) | 1.1 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Rosa californica</i> | 50 (-) | 2 (-) | 1 (-) | 3 (-) |
| | <i>Acmispon glaber</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Artemisia californica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Brickellia californica</i> | 25 (100) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Ceanothus spinosus</i> | 25 (-) | 8 (-) | 8 (-) | 8 (-) |
| | <i>Eriophyllum confertiflorum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Keckiella antirrhinoides</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Mimulus guttatus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Rhamnus ilicifolia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rubus ursinus</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| | * <i>Tamarix</i> | 25 (-) | 9 (-) | 9 (-) | 9 (-) |
| | <i>Toxicodendron diversilobum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | | | | | |
| | | | | | |
| Herb | <i>Artemisia douglasiana</i> | 75 (100) | 15.7 (0.2) | 0.2 (0.2) | 40 (0.2) |
| | * <i>Bromus diandrus</i> | 50 (-) | 15 (-) | 5 (-) | 25 (-) |
| | <i>Ambrosia psilostachya</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| | <i>Asclepias fascicularis</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | * <i>Avena barbata</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Bromus hordeaceus</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| | * <i>Bromus madritensis</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Calystegia macrostegia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Dactylis glomerata</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| | <i>Datura</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Drymocallis lactea</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Elymus condensatus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | | | | | |
| | | | | | |
| | | | | | |

| Species Name | Con | Ave | Min | Max |
|---------------------------------|----------|-----------|-----------|-----------|
| <i>Elymus glaucus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Elymus triticoides</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Erigeron foliosus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Juncus rugulosus</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Juncus textilis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Lupinus latifolius</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Rafinesquia californica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Scrophularia californica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stachys</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Stachys rigida</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Thalictrum fendleri</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| * <i>Vicia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Platanus racemosa* Alliance**

***Platanus racemosa-Quercus agrifolia* Association**

Samples Used To Describe Association (n= 36)

Rapid Assessment(s), WR and WSD stands: 1550MB02, SDRP0196, 1708JD02, 1067MB01, 0505MG02, 0504TK24, 1350EB02, 0507TK04, 1708JD01, 0309AK21, 1136MB01, 0505AK03, SDRP0323, 1576LS01, 0310DH26, 0610DH04, 1126MB01, 1068MB02, 1710JD08, 0603JS02, 1710KH04, 1123MB01, 0611RB01, 1041MB01, 0311TK23, SDRP0433, 0324TKF3, SDRP0194, 1800FS01, SDRP0476, 0610DH01, SDRP0191, OGPSMB03
Rapid Assessment(s), OC stands (n=3): 1123EK01, 1229EK01, 1450EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-----------------|-------------|
| Total vegetation cover (%) | 67.3 (56.3) | 24-97 (48-66) | - |
| Herb cover (%) | 19.4 (24.3) | 2-60 (7-54) | 0-2 (0-1) |
| Shrub cover (%) | 12.7 (12.7) | 0.2-32 (0.2-32) | 0-10 (1-5) |
| Tall shrub / low tree cover (%) | 4.8 (4.8) | 0.2-14 (0.2-14) | 0-15 (0-5) |
| Overstory tree cover (%) | 33.5 (23.0) | 8-75 (14-38) | 5-20 (5-15) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (13,0), Flat (7,1), NE (5,0), NW (5,0), S (1,1), SE (5,1)
Macrotopography: lower 1/3 of slope (1,1), lower to middle 1/3 of slope (1,0), bottom to lower 1/3 of slope (6,0), bottom (26,1), middle 1/3 of slope (2,1)
Microtopography: convex (2,0), flat (12,1), undulating (3,0), concave (19,2)
Parent material: granitic (9,0), gravelly alluvium (1,0), metamorphic (1,0), mixed alluvium (3,1), sandstone (1,1), sandy alluvium (8,0), silty alluvium (1,0)
Soil texture: unknown (1,0), silty loam (1,0), silt (1,0), sandy loam (20,0), sand (4,0), clay loam (5,1), loam (3,1)

| | Average | Range |
|------------------------------|-------------|--------------------|
| Elevation (ft) | 797 (795) | 49-2060 (640-1076) |
| Slope (degrees) | 3 (2.0) | 0-28 (0-4) |
| Large rock cover (%) | 3.8 (0.3) | 0-69 (0-1) |
| Small rock cover (%) | 10.0 (1.0) | 0-73 (0-3) |
| Bare ground cover (%) | 14.2 (22.3) | 0-86 (15-37) |
| Litter cover (%) | 56.3 (72.0) | 0-94 (54-82) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|-----------|-------------|-----------|-----------|
| Tree | <i>Platanus racemosa</i> | 100 (100) | 11.3 (14.7) | 1 (1) | 63 (35) |
| | <i>Quercus agrifolia</i> | 81 (100) | 20.9 (9.1) | 0.2 (6) | 58 (12.2) |
| | <i>Salix lasiolepis</i> | 58 (67) | 7.2 (5.1) | 0.2 (0.2) | 25 (10) |
| | <i>Salix laevigata</i> | 36 (33) | 9.4 (2) | 1 (2) | 35 (2) |
| Shrub | <i>Toxicodendron diversilobum</i> | 86 (67) | 15.5 (8.1) | 0.2 (0.2) | 55 (16) |
| | <i>Baccharis salicifolia</i> | 42 (67) | 2.7 (1.1) | 0.2 (0.2) | 10 (2) |
| | <i>Rubus ursinus</i> | 33 (33) | 5.6 (14) | 0.2 (14) | 14 (14) |
| | <i>Sambucus nigra subsp. caerulea</i> | 25 (33) | 1.8 (0.2) | 0.2 (0.2) | 7 (0.2) |
| Herb | <i>Artemisia douglasiana</i> | 56 (67) | 3.8 (3.1) | 0.2 (0.2) | 12 (6) |
| | * <i>Bromus diandrus</i> | 50 (100) | 14.6 (18.1) | 0.2 (0.2) | 52 (52) |
| | <i>Vitis girdiana</i> | 28 (-) | 14.9 (-) | 0.2 (-) | 55 (-) |
| | <i>Ambrosia psilostachya</i> | 25 (-) | 2.6 (-) | 0.2 (-) | 7 (-) |
| | <i>Carex spissa</i> | 25 (-) | 2.4 (-) | 0.2 (-) | 10 (-) |
| | <i>Nasturtium officinale</i> | 25 (33) | 0.8 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | * <i>Stipa miliacea</i> var. <i>miliacea</i> | 25 (33) | 1.7 (1) | 0.2 (1) | 4 (1) |

Platanus racemosa Alliance

Platanus racemosa-*Salix laevigata*/*Salix lasiolepis*-*Baccharis salicifolia* Association

Samples Used To Describe Association (n= 9)

Rapid Assessment(s), WR and WSD stands: SCWP.004, RECA.016, SITT.038, CLNF.026, SITT.040/BPU 122139-354, SITT.010, SMER.065/River.07, SAAN.020

Rapid Assessment(s), OC stands (n=1): 1513EK05

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 49.4 (45.0) | 20-78 (45-45) | - |
| Herb cover (%) | 10.4 (9.0) | 0-25 (9-9) | 0-2 (0-0.5) |
| Shrub cover (%) | 22.8 (10.0) | 5-42 (10-10) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | 8.0 (8.0) | 8-8 (8-8) | 1-5 (2-5) |
| Overstory tree cover (%) | 20.7 (31.0) | 9-38 (31-31) | 10-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: W (1,1), SW (6,0), Flat (1,0), SE (1,0)

Macrotopography: bottom (7,1), lower 1/3 of slope (2,0)

Microtopography: Concave (8,1), Flat (1,0)

Parent material: granitic (1,0), gravelly alluvium (1,0), mixed alluvium (2,0), sandstone (1,1), sandy alluvium (3,0), silty alluvium (1,0)

Soil texture: silty loam (1,0), sand (5,1), sandy loam (2,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 25

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 1095 (804) | 700-2146 (804-804) |
| Slope (degrees) | 2 (2.0) | 1-4 (2-2) |
| Large rock cover (%) | 23.0 (-) | 0-85 (-) |
| Small rock cover (%) | 18.1 (-) | 0-60 (-) |
| Bare ground cover (%) | 8.3 (10.0) | 0-20 (10-10) |
| Litter cover (%) | 31.2 (82.0) | 0-85 (82-82) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-----------------------------------|-----------|-----------|---------|---------|
| Tree | <i>Platanus racemosa</i> | 100 (100) | 12.7 (32) | 1 (32) | 32 (32) |
| | <i>Salix laevigata</i> | 89 (100) | 9 (6) | 4 (6) | 14 (6) |
| | <i>Quercus agrifolia</i> | 56 (100) | 2.6 (2) | 1 (2) | 5 (2) |
| | <i>Fraxinus velutina</i> | 33 (-) | 7.1 (-) | 0.2 (-) | 21 (-) |
| Shrub | <i>Baccharis salicifolia</i> | 100 (100) | 7.2 (2) | 2 (2) | 14 (2) |
| | <i>Salix lasiolepis</i> | 78 (-) | 10.9 (-) | 2 (-) | 23 (-) |
| | <i>Toxicodendron diversilobum</i> | 56 (100) | 2 (5) | 1 (5) | 5 (5) |
| | <i>Amorpha fruticosa</i> | 33 (-) | 1.3 (-) | 1 (-) | 2 (-) |
| Herb | <i>Artemisia douglasiana</i> | 33 (-) | 1.7 (-) | 1 (-) | 3 (-) |
| | <i>Typha domingensis</i> | 33 (-) | 3.7 (-) | 1 (-) | 5 (-) |

***Pseudotsuga macrocarpa* Alliance**

***Pseudotsuga macrocarpa*-*Quercus agrifolia* Association**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1205EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 60.0 (60.0) | 60-60 (60-60) | - |
| Herb cover (%) | - (-) | - (-) | - (-) |
| Shrub cover (%) | 23.0 (23.0) | 23-23 (23-23) | 2-5 (2-5) |
| Tall shrub / low tree cover (%) | 22.0 (22.0) | 22-22 (22-22) | 5-10 (5-10) |
| Overstory tree cover (%) | 27.0 (27.0) | 27-27 (27-27) | 10-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,1)

Macrotopography: lower to middle 1/3 of slope (1,1)

Microtopography: undulating (1,1)

Parent material: metamorphic (1,1)

Soil texture:

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 26

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 2110 (2110) | 2110-2110 (2110-2110) |
| Slope (degrees) | 40 (40.0) | 40-40 (40-40) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | - (-) | - (-) |
| Litter cover (%) | - (-) | - (-) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|-----------|-----------|-----------|-----------|
| Tree | <i>Acer macrophyllum</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Pseudotsuga macrocarpa</i> | 100 (100) | 25 (25) | 25 (25) | 25 (25) |
| | <i>Quercus agrifolia</i> | 100 (100) | 20 (20) | 20 (20) | 20 (20) |
| | | | | | |
| Shrub | <i>Cercocarpus betuloides</i> var. <i>betuloides</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Fraxinus dipetala</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Heteromeles arbutifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Keckiella cordifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Lonicera subspicata</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Rhamnus ilicifolia</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| | <i>Sambucus nigra</i> subsp. <i>caerulea</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Toxicodendron diversilobum</i> | 100 (100) | 12 (12) | 12 (12) | 12 (12) |
| | | | | | |
| | | | | | |
| Herb | <i>Dryopteris arguta</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | | | | | |

***Pseudotsuga macrocarpa* Alliance**

***Pseudotsuga macrocarpa*-*Quercus chrysolepis* Association**

Samples Used To Describe Association (n= 5)

Rapid Assessment(s), WR and WSD stands: CLNF.100, CLNF.088, CLNF.087, CLNF.080
Rapid Assessment(s), OC stands (n=1): 1210EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 69.0 (72.0) | 45-82 (72-72) | - |
| Herb cover (%) | 7.5 (13.0) | 0-15 (13-13) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 22.2 (9.0) | 2-50 (9-9) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 54.8 (-) | 35-68 (-) | 15-20 (15-20) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NE (4,0)
Macrotopography: middle 1/3 of slope (2,0), upper 1/3 of slope (3,1)
Microtopography: Concave (5,1)
Parent material: granitic (4,0), metamorphic (1,1)
Soil texture: loam (1,0), sandy loam (3,0)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3821 (4052) | 3009-4126 (4052-4052) |
| Slope (degrees) | 41 (40.0) | 38-50 (40-40) |
| Large rock cover (%) | 9.8 (-) | 0-25 (-) |
| Small rock cover (%) | 9.0 (-) | 0-25 (-) |
| Bare ground cover (%) | 6.0 (8.0) | 0-15 (8-8) |
| Litter cover (%) | 50.0 (90.0) | 0-90 (90-90) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-----------|-----------|---------|
| Tree | <i>Pseudotsuga macrocarpa</i> | 100 (100) | 31.4 (44) | 20 (44) | 44 (44) |
| | <i>Quercus chrysolepis</i> | 100 (100) | 35.8 (23) | 20.2 (23) | 52 (23) |
| | <i>Pinus coulteri</i> | 40 (-) | 2 (-) | 1 (-) | 3 (-) |
| | <i>Quercus wislizeni</i> | 40 (-) | 9 (-) | 8 (-) | 10 (-) |
| | | | | | |
| Shrub | <i>Heteromeles arbutifolia</i> | 80 (100) | 1.1 (1) | 0.2 (1) | 2 (1) |
| | <i>Arctostaphylos glandulosa</i> | 60 (-) | 3.7 (-) | 1 (-) | 5 (-) |
| | <i>Keckiella cordifolia</i> | 60 (100) | 3.1 (8) | 0.2 (8) | 8 (8) |
| | <i>Dendromecon rigida</i> | 40 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Eriogonum fasciculatum</i> | 40 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | | | | | |
| Herb | <i>*Bromus diandrus</i> | 40 (100) | 2.6 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | | | | | |

Quercus agrifolia Alliance

Quercus agrifolia Association

Samples Used To Describe Association (n= 9)

Rapid Assessment(s), WR and WSD stands: SAGE.023, SAAN.015, SAGE.044, BUCK.021, BLACK.034/97
Rapid Assessment(s), OC stands (n=4): 1066EK03, 1133EK01, 1173EK02, 1195EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 60.6 (48.8) | 40-82 (40-60) | - |
| Herb cover (%) | 18.8 (21.0) | 1-50 (1-35) | 0-1 (0-0.5) |
| Shrub cover (%) | 10.9 (5.0) | 1-26 (1-10) | 0-2 (0-2) |
| Tall shrub / low tree cover (%) | 0.6 (0.6) | 0.2-1 (0.2-1) | 0-5 (0-2) |
| Overstory tree cover (%) | 44.8 (32.7) | 9-81 (9-48) | 5-15 (5-15) |

Local Environmental Description (All stands, OC stands)

Aspect: W (1,1), Variable (1,0), SW (2,2), NW (2,0), NE (1,0), Flat (1,0), N (1,1)
Macrotopography: bottom to lower 1/3 of slope (1,1), lower 1/3 of slope (6,2), lower to middle 1/3 of slope (1,1), upper 1/3 of slope (1,0)
Microtopography: Undulating (3,1), Concave (2,1), Flat (3,1)
Parent material: granitic (3,0), metamorphic (1,1), mix of two or more rock types (1,0), mixed alluvium (1,1), sandy alluvium (1,0)
Soil texture: sandy loam (5,1), clay loam (2,2), loam (1,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 28

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1671 (965) | 636-3810 (636-1657) |
| Slope (degrees) | 12 (8.8) | 0-40 (1-15) |
| Large rock cover (%) | 1.2 (0.3) | 0-10 (0-1) |
| Small rock cover (%) | 4.2 (0.5) | 0-25 (0-2) |
| Bare ground cover (%) | 19.8 (31.0) | 0-52 (19-52) |
| Litter cover (%) | 52.0 (66.3) | 0-90 (45-80) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---------------------------------------|-----------|-------------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 100 (100) | 40 (30.3) | 7 (7) | 80 (48.2) |
| Shrub | | | | | |
| | <i>Sambucus nigra subsp. caerulea</i> | 33 (75) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| Herb | | | | | |
| | * <i>Bromus diandrus</i> | 33 (75) | 13.4 (13.4) | 0.2 (0.2) | 25 (25) |
| | * <i>Carduus pycnocephalus</i> | 33 (75) | 2.1 (2.1) | 0.2 (0.2) | 4 (4) |

***Quercus agrifolia* Alliance**

***Quercus agrifolia*/*Artemisia californica* Association**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: SDRP0268, SDRP0085, 1550MB01, SDRP0189, SDRP0482

Rapid Assessment(s), OC stands (n=1): 1499KH03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 65.0 (23.0) | 23-85 (23-23) | - |
| Herb cover (%) | 27.2 (1.0) | 1-50 (1-1) | 0-1 (0-0.5) |
| Shrub cover (%) | 15.0 (15.0) | 15-15 (15-15) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-1 (-) |
| Overstory tree cover (%) | 27.2 (12.0) | 10-45 (12-12) | 2-20 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (4,0), SE (1,0), SW (1,1)

Macrotopography: lower 1/3 of slope (2,0), lower to middle 1/3 of slope (1,0), bottom to lower 1/3 of slope (1,0), bottom (1,0), upper 1/3 of slope to ridgetop (1,1)

Microtopography: undulating (2,0), concave (2,0), flat (2,1)

Parent material: granitic (3,0), igneous (1,0)

Soil texture: unknown (1,1), clay loam (1,0), sandy loam (4,0)

| | Average | Range |
|-----------------------|-------------|----------------------|
| Elevation (ft) | 571 (1401) | 299-1401 (1401-1401) |
| Slope (degrees) | 19 (26.0) | 1-33 (26-26) |
| Large rock cover (%) | 1.7 (-) | 0-10 (-) |
| Small rock cover (%) | 0.8 (-) | 0-5 (-) |
| Bare ground cover (%) | 18.3 (89.0) | 0-89 (89-89) |
| Litter cover (%) | 29.2 (10.0) | 0-85 (10-10) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-----------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 100 (100) | 20.7 (12) | 4 (12) | 46.2 (12) |
| Shrub | | | | | |
| | <i>Artemisia californica</i> | 100 (100) | 5.7 (0.2) | 0.2 (0.2) | 20 (0.2) |
| | <i>Malosma laurina</i> | 83 (100) | 3 (1) | 0.2 (1) | 5 (1) |
| | <i>Eriogonum fasciculatum</i> | 67 (100) | 3.8 (1) | 1 (1) | 7 (1) |
| | <i>Mimulus aurantiacus</i> | 67 (100) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Salvia apiana</i> | 67 (100) | 2.8 (0.2) | 0.2 (0.2) | 7 (0.2) |
| | <i>Rhamnus crocea</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Acemison glaber</i> | 33 (100) | 4.5 (7) | 2 (7) | 7 (7) |
| | <i>Keckiella antirrhinoides</i> | 33 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| | <i>Rhamnus ilicifolia</i> | 33 (-) | 19.5 (-) | 1 (-) | 38 (-) |
| | <i>Toxicodendron diversilobum</i> | 33 (-) | 5.1 (-) | 0.2 (-) | 10 (-) |
| Herb | | | | | |
| | * <i>Bromus diandrus</i> | 83 (-) | 9.4 (-) | 0.2 (-) | 25 (-) |
| | * <i>Bromus madritensis</i> | 83 (-) | 10.6 (-) | 2 (-) | 23 (-) |
| | * <i>Bromus hordeaceus</i> | 50 (-) | 6 (-) | 1 (-) | 10 (-) |
| | * <i>Hirschfeldia incana</i> | 50 (-) | 2.1 (-) | 0.2 (-) | 3 (-) |
| | * <i>Centaurea melitensis</i> | 33 (-) | 3 (-) | 2 (-) | 4 (-) |

Quercus agrifolia Alliance

Quercus agrifolia/Quercus (berberidifolia, xacutidens) Association

Samples Used To Describe Association (n= 8)

Rapid Assessment(s), WR and WSD stands: 0506RB24, 1362JD02, 0520DH04, 1693FS02, 0609AK01, 0609JJ01, 0423RB01

Rapid Assessment(s), OC stands (n=1): 1057EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|-------------|
| Total vegetation cover (%) | 45.5 (40.0) | 38-66 (40-40) | - |
| Herb cover (%) | 20.0 (10.0) | 0.2-50 (10-10) | 0-1 (0-0.5) |
| Shrub cover (%) | 10.0 (10.0) | 10-10 (10-10) | 1-5 (1-2) |
| Tall shrub / low tree cover (%) | 6.0 (6.0) | 6-6 (6-6) | 1-5 (2-5) |
| Overstory tree cover (%) | 24.6 (15.0) | 8-45 (15-15) | 2-15 (2-5) |

Local Environmental Description All stands, OC stands)

Aspect: N (1,1), SE (2,0), SW (1,0), Flat (1,0), NW (2,0), NE (1,0)

Macrotopography: bottom to lower 1/3 of slope (1,0), lower 1/3 of slope (1,0), lower to middle 1/3 of slope (2,1), middle 1/3 of slope (2,0), middle to upper 1/3 of slope (1,0), upper 1/3 of slope (1,0)

Microtopography: concave (3,0), convex (1,1), flat (1,0), undulating (3,0)

Parent material: granitic (3,0), metamorphic (1,0), mixed alluvium (1,0), sandstone (1,1)

Soil texture: clay (1,0), clay loam (2,0), sandy loam (5,1)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 30

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 1468 (180) | 180-2536 (180-180) |
| Slope (degrees) | 19 (30.0) | 2-34 (30-30) |
| Large rock cover (%) | 12.2 (1.0) | 0-53 (1-1) |
| Small rock cover (%) | 8.5 (-) | 0-22 (-) |
| Bare ground cover (%) | 17.8 (37.0) | 1-49 (37-37) |
| Litter cover (%) | 59.8 (60.0) | 10-90 (60-60) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---|-----------|-----------|-----------|-----------|
| Tree | <i>Quercus agrifolia</i> | 100 (100) | 24.1 (15) | 8 (15) | 46 (15) |
| | | | | | |
| Shrub | <i>Toxicodendron diversilobum</i> | 75 (100) | 6.2 (3) | 2 (3) | 12 (3) |
| | <i>Quercus berberidifolia</i> | 63 (100) | 6.6 (7) | 1 (7) | 9 (7) |
| | <i>Heteromeles arbutifolia</i> | 50 (100) | 2.6 (1) | 0.2 (1) | 5 (1) |
| | <i>Lonicera subspicata</i> | 50 (-) | 2.6 (-) | 0.2 (-) | 8 (-) |
| | <i>Eriogonum fasciculatum</i> | 38 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Malosma laurina</i> | 38 (-) | 2 (-) | 1 (-) | 4 (-) |
| | <i>Mimulus aurantiacus</i> | 38 (100) | 1.4 (2) | 0.2 (2) | 2 (2) |
| | <i>Quercus acutidens</i> | 38 (-) | 7 (-) | 3 (-) | 13 (-) |
| | <i>Sambucus nigra subsp. caerulea</i> | 38 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Acmispon glaber</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Adenostoma fasciculatum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Arctostaphylos glandulosa</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Artemisia californica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Ceanothus tomentosus</i> | 25 (-) | 7.6 (-) | 0.2 (-) | 15 (-) |
| | <i>Eriophyllum confertiflorum var. confertiflorum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Frangula californica</i> | 25 (-) | 4.5 (-) | 4 (-) | 5 (-) |
| | <i>Keckiella cordifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Polygala cornuta var. fishiae</i> | 25 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Rhus ovata</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| Herb | <i>*Bromus diandrus</i> | 63 (100) | 17 (0.2) | 0.2 (0.2) | 38 (0.2) |
| | <i>*Bromus hordeaceus</i> | 25 (-) | 3 (-) | 2 (-) | 4 (-) |
| | <i>*Bromus madritensis</i> | 25 (-) | 7.6 (-) | 0.2 (-) | 15 (-) |
| | <i>*Carduus pycnocephalus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Erodium botrys</i> | 25 (-) | 14 (-) | 3 (-) | 25 (-) |
| | <i>Galium angustifolium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Galium aparine</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Marah macrocarpus var. macrocarpus</i> | 25 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Pseudognaphalium californicum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | | | | | |

Quercus agrifolia Alliance

Quercus agrifolia/*Salix lasiolepis* Association

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: 1355MB02, 0308TK21, 1710JD02

Rapid Assessment(s), OC stands (n=1): 1415EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 65.0 (60.0) | 48-77 (60-60) | - |
| Herb cover (%) | 11.8 (19.0) | 5-19 (19-19) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 11.0 (11.0) | 11-11 (11-11) | 0.5-10 (1-2) |
| Tall shrub / low tree cover (%) | 26.0 (26.0) | 26-26 (26-26) | 0-10 (5-10) |
| Overstory tree cover (%) | 40.5 (23.0) | 19-65 (23-23) | 5-15 (10-15) |

Local Environmental Description All stands, OC stands)

Aspect: Flat (1,0), NE (1,0), SW (2,1)

Macrotopography: bottom (4,1)

Microtopography: concave (1,1), flat (3,0)

Parent material: sandstone (1,1), sandy alluvium (1,0)

Soil texture: clay (1,0), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|-------------------|
| Elevation (ft) | 369 (440) | 108-810 (440-440) |
| Slope (degrees) | 2 (4.0) | 0-4 (4-4) |
| Large rock cover (%) | 0.3 (1.0) | 0-1 (1-1) |
| Small rock cover (%) | 2.8 (2.0) | 0-9 (2-2) |
| Bare ground cover (%) | 21.8 (35.0) | 4-35 (35-35) |
| Litter cover (%) | 73.8 (60.0) | 60-95 (60-60) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| Tree | | | | |
| <i>Salix lasiolepis</i> | 100 (100) | 14.8 (24) | 7 (24) | 24 (24) |
| <i>Quercus agrifolia</i> | 75 (100) | 30.7 (24) | 16 (24) | 52 (24) |
| * <i>Phoenix canariensis</i> | 50 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Salix gooddingii</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Nicotiana glauca</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Platanus racemosa</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Quercus agrifolia</i> var. <i>agrifolia</i> | 25 (-) | 60 (-) | 60 (-) | 60 (-) |
| <i>Salix laevigata</i> | 25 (-) | 17 (-) | 17 (-) | 17 (-) |
| * <i>Washingtonia robusta</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| Shrub | | | | |
| <i>Toxicodendron diversilobum</i> | 100 (100) | 18.3 (10) | 10 (10) | 30 (10) |
| <i>Baccharis pilularis</i> | 50 (100) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |
| <i>Heteromeles arbutifolia</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Sambucus nigra</i> subsp. <i>caerulea</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| <i>Baccharis salicifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Cercocarpus minutiflorus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Clematis ligusticifolia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 32

| Species Name | Con | Ave | Min | Max |
|---|----------|-----------|-----------|-----------|
| <i>Eriogonum fasciculatum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Malosma laurina</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Rubus ursinus</i> | 25 (100) | 12 (12) | 12 (12) | 12 (12) |
| Herb | | | | |
| <i>*Brassica nigra</i> | 75 (100) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>*Bromus diandrus</i> | 75 (100) | 2.7 (1) | 1 (1) | 5 (1) |
| <i>Ambrosia psilostachya</i> | 50 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Artemisia palmeri</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Bromus hordeaceus</i> | 50 (-) | 2.5 (-) | 2 (-) | 3 (-) |
| <i>*Carduus pycnocephalus</i> | 50 (100) | 4.1 (0.2) | 0.2 (0.2) | 8 (0.2) |
| <i>*Hypochaeris glabra</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| <i>*Marrubium vulgare</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Rumex crispus</i> | 50 (100) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>*Anagallis arvensis</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Anemopsis californica</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Artemisia douglasiana</i> | 25 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>*Avena fatua</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>*Brachypodium distachyon</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Carex triquetra</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Carpobrotus edulis</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>*Cortaderia selloana</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Dipsacus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Distichlis spicata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Elymus condensatus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Epilobium brachycarpum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Foeniculum vulgare</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Galium aparine</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Geranium dissectum</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Hirschfeldia incana</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Marah macrocarpus var. macrocarpus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Medicago polymorpha</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Melica imperfecta</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Melilotus indicus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Nasturtium officinale</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Phacelia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Plantago major</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Polypodium californicum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pteridium aquilinum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Sonchus oleraceus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Stipa miliacea var. miliacea</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Urtica dioica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Vitis girdiana</i> | 25 (-) | 10 (-) | 10 (-) | 10 (-) |

***Quercus agrifolia* Alliance**

***Quercus agrifolia*/Toxicodendron diversilobum/Grass Association**

Samples Used To Describe Association (n= 52)

Rapid Assessment(s), WR and WSD stands: SMER.022/Scrub trail #1, 1811TK01, 0423AK03, 1350JD01, SRPL.003/422, 0324DHF6, 0421RB02, 0421TK02, 1202KH02, 0423TK01, 1202KH01, 1782MB02, 0422TK03, HEME.009, 1030MB01, 1708MM02, 0507DH03, OGPSMB02, SDG0110, 0504DH22, 1075FS01, 0422DH03, 0603RB03, SDRP0069, 0505TK01, SDRP0071, 0625JJ02, 0521DH02, 0504TK01, SDRP0134, SITT.022/450/BPU 3255-350, 0604JJ01

Rapid Assessment(s), OC stands (n=20): 1015EK06, 1019EK02, 1021EK01, 1048EK01, 1056EK03, 1069EK02, 1073EK01, 1110EK02, 1133EK03, 1172MA01, 1214KH01, 1237EK01, 1253MA01, 1289EK01, 1429EK01, 1449EK01, 1456EK01, 1463EK04, 1520MA01, ek3.8.3

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|-----------------|-------------------|
| Total vegetation cover (%) | 58.0 (55.9) | 29-88 (37-75) | - |
| Herb cover (%) | 21.7 (16.0) | 0.2-85 (0.2-50) | 0-1 (0-1) |
| Shrub cover (%) | 14.8 (13.8) | 1-42 (1-42) | 0-5 (0-5) |
| Tall shrub / low tree cover (%) | 4.1 (4.1) | 0.2-20 (0.2-20) | 0-5 (0-5) |
| Overstory tree cover (%) | 35.5 (37.8) | 7-75 (7-65) | 2-20 (2-15) |

Local Environmental Description (All stands, OC stands)

Aspect: S (2,2), W (2,1), Variable (3,1), SE (9,3), NW (8,0), NE (9,1), N (4,4), Flat (5,2), E (2,2), SW (8,4)

Macrotopography: lower to middle 1/3 of slope (3,0), upper 1/3 of slope (1,0), middle to upper 1/3 of slope (3,2), lower 1/3 of slope (10,5), bottom to mid 1/3 of slope (1,0), bottom to lower 1/3 of slope (11,8), bottom (13,4), middle 1/3 of slope (9,1)

Microtopography: concave (24,10), flat (11,5), undulating (16,5)

Parent material: diorite (1,0), gabbro (1,0), granitic (12,1), gravelly alluvium (2,0), igneous (1,0), metamorphic (2,0), mixed alluvium (3,1), sandstone (15,15), sandy alluvium (6,0), sedimentary (1,0)

Soil texture: clay (1,1), silty loam (1,0), sandy loam (27,8), clay loam (8,3), loam (9,5), sand (1,0)

| | Average | Range |
|------------------------------|----------------|---------------------|
| Elevation (ft) | 1011 (916) | 135-3330 (459-3330) |
| Slope (degrees) | 10 (7.8) | 0-36 (0-35) |
| Large rock cover (%) | 4.4 (-) | 0-50 (-) |
| Small rock cover (%) | 6.4 (1.7) | 0-30 (0-15) |
| Bare ground cover (%) | 16.3 (20.6) | 0-46 (4-42) |
| Litter cover (%) | 62.2 (75.4) | 0-95 (55-94) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|------------|-------------|------------|------------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 79 (100) | 33.1 (39.6) | 4 (4) | 71 (70) |
| Shrub | | | | | |
| | <i>Toxicodendron diversilobum</i> | 98 (100) | 8 (6.1) | 0.2 (0.2) | 35 (27) |
| | <i>Rhamnus ilicifolia</i> | 42 (45) | 3 (4) | 0.2 (0.2) | 10 (10) |
| | <i>Mimulus aurantiacus</i> | 38 (20) | 1.2 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 31 (45) | 4.9 (5) | 0.2 (0.2) | 18 (18) |
| | <i>Malosma laurina</i> | 31 (25) | 3.2 (0.9) | 0.2 (0.2) | 18 (2) |
| | <i>Rhus integrifolia</i> | 25 (55) | 3.2 (3.7) | 0.2 (0.2) | 20 (20) |
| Herb | | | | | |

| Species Name | Con | Ave | Min | Max |
|---|---------|-------------|-----------|---------|
| <i>*Bromus diandrus</i> | 77 (85) | 16.8 (11.8) | 0.2 (0.2) | 60 (35) |
| <i>*Carduus pycnocephalus</i> | 38 (60) | 2.5 (2.5) | 0.2 (0.2) | 15 (15) |
| <i>Galium aparine</i> | 29 (45) | 0.7 (0.3) | 0.2 (0.2) | 5 (1) |
| <i>Elymus condensatus</i> | 27 (55) | 0.5 (0.4) | 0.2 (0.2) | 2 (2) |
| <i>Melica imperfecta</i> | 27 (30) | 0.8 (0.3) | 0.2 (0.2) | 5 (1) |
| <i>Marah macrocarpus</i> var. <i>macrocarpus</i> | 25 (-) | 0.5 (-) | 0.2 (-) | 2 (-) |

***Quercus chrysolepis* Alliance** ***Quercus chrysolepis* Association**

Samples Used To Describe Association (n= 15)

Rapid Assessment(s), WR and WSD stands: IDYL.030/1427, CAHUI.006/689, ANZA.123, PALM.010, CLNF.083, IDYL.064/BPU 5456-333, ANZA.132, IDYL.032/1166, ANZA.138, IDYL.066/81, LAFU.021/274

Rapid Assessment(s), OC stands (n=4): 1184JD01, 1209EK03, 1215JD01, 1240EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|---------------|
| Total vegetation cover (%) | 70.1 (71.5) | 53-86 (56-86) | - |
| Herb cover (%) | 11.8 (1.5) | 0-48 (1-2) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 18.1 (17.5) | 2-74 (8-27) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-5 (-) |
| Overstory tree cover (%) | 57.9 (59.0) | 37-75 (48-70) | 10-20 (10-20) |

Local Environmental Description (All stands, OC stands)

Aspect: N (3,3), NE (8,1), NW (1,0), SW (3,0)

Macrotopography: bottom (1,0), middle 1/3 of slope (4,0), middle to upper 1/3 of slope (1,1), upper 1/3 of slope (9,3)

Microtopography: Flat (1,0), Undulating (9,3), convex (1,1), Concave (4,0)

Parent material: granitic (9,0), gravelly alluvium (1,1), igneous (1,0), metamorphic (2,1)

Soil texture: clay loam (1,1), sandy loam (14,3)

| | Average | Range |
|------------------------------|-------------|-----------------------|
| Elevation (ft) | 5080 (4485) | 4026-6374 (4026-5295) |
| Slope (degrees) | 24 (28.3) | 3-40 (15-40) |
| Large rock cover (%) | 7.0 (-) | 0-25 (-) |
| Small rock cover (%) | 10.1 (1.3) | 0-30 (0-5) |
| Bare ground cover (%) | 12.1 (14.0) | 2-38 (2-38) |
| Litter cover (%) | 67.1 (83.3) | 35-98 (60-98) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-------------------------------|-----------|-----------|-----------|---------|
| Tree | <i>Quercus chrysolepis</i> | 100 (100) | 53.5 (57) | 7 (48) | 82 (70) |
| | <i>Quercus wislizeni</i> | 67 (75) | 4.7 (5.3) | 0.2 (2) | 10 (10) |
| | <i>Pinus coulteri</i> | 40 (25) | 2.5 (2) | 2 (2) | 4 (2) |
| | | | | | |
| Shrub | <i>Ceanothus integerrimus</i> | 27 (25) | 5.1 (8) | 0.2 (8) | 9 (8) |
| | <i>Clematis lasiantha</i> | 27 (100) | 2.1 (2.1) | 0.2 (0.2) | 5 (5) |
| | <i>Lonicera subspicata</i> | 27 (25) | 0.8 (1) | 0.2 (1) | 1 (1) |
| | | | | | |

| | Species Name | Con | Ave | Min | Max |
|------|------------------------------|---------|-----------|-----------|---------|
| Herb | | | | | |
| | <i>*Bromus tectorum</i> | 47 (-) | 7.6 (-) | 0.2 (-) | 18 (-) |
| | <i>Galium angustifolium</i> | 40 (50) | 0.3 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Cryptantha intermedia</i> | 33 (-) | 2.4 (-) | 0.2 (-) | 7 (-) |

***Quercus chrysolepis* Alliance**

***Quercus chrysolepis*-*Pseudotsuga macrocarpa* Association**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: BLACK.016/858, CLNF.033, VAIL.011, IDYL.001
Rapid Assessment(s), OC stands (n=2): 1209EK01, 1230EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 70.2 (63.0) | 54-93 (54-72) | - |
| Herb cover (%) | 23.0 (4.0) | 1-85 (4-4) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 14.0 (19.0) | 1-32 (6-32) | 0-5 (1-5) |
| Tall shrub / low tree cover (%) | 4.0 (4.0) | 4-4 (4-4) | 1-5 (2-5) |
| Overstory tree cover (%) | 60.5 (54.5) | 48-75 (51-58) | 15-50 (15-50) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,0), SE (3,1), NE (1,0), N (1,1)

Macrotopography: middle 1/3 of slope (3,0), middle to upper 1/3 of slope (1,1), upper 1/3 of slope (2,1)

Microtopography: Convex (1,0), Undulating (2,1), concave (3,1)

Parent material: granitic (4,1), metamorphic (1,1)

Soil texture: loam (1,0), sand (1,1), sandy loam (1,0)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 4243 (3980) | 3825-4854 (3980-3980) |
| Slope (degrees) | 26 (35.0) | 11-36 (35-35) |
| Large rock cover (%) | 9.3 (10.0) | 0-30 (0-20) |
| Small rock cover (%) | 8.5 (10.0) | 0-20 (0-20) |
| Bare ground cover (%) | 8.5 (10.5) | 0-25 (9-12) |
| Litter cover (%) | 54.5 (67.5) | 0-88 (50-85) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-----------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>Pseudotsuga macrocarpa</i> | 100 (100) | 5 (5) | 2 (2) | 10 (8) |
| | <i>Quercus chrysolepis</i> | 100 (100) | 54.2 (46) | 42 (42) | 66 (50) |
| | <i>Acer macrophyllum</i> | 50 (100) | 3.4 (2.1) | 0.2 (0.2) | 6 (4) |
| | <i>Alnus rhombifolia</i> | 33 (50) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Platanus racemosa</i> | 33 (50) | 2.1 (0.2) | 0.2 (0.2) | 4 (0.2) |
| | <i>Quercus wislizeni</i> | 33 (50) | 4 (4) | 4 (4) | 4 (4) |
| | <i>Salix laevigata</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Toxicodendron diversilobum</i> | 50 (100) | 3.4 (2.1) | 0.2 (0.2) | 6 (4) |
| | <i>Heteromeles arbutifolia</i> | 33 (50) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>Yucca whipplei</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

| Species Name | Con | Ave | Min | Max |
|--------------------------|----------|-----------|-----------|---------|
| Herb | | | | |
| <i>Dryopteris arguta</i> | 50 (100) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Melica imperfecta</i> | 50 (50) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Galium aparine</i> | 33 (50) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |

Quercus wislizeni Alliance

Quercus wislizeni/*Arctostaphylos glandulosa* Provisional Association

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1275JD02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 66.0 (66.0) | 66-66 (66-66) | - |
| Herb cover (%) | - (-) | - (-) | - (-) |
| Shrub cover (%) | - (-) | - (-) | - (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1)

Macrotopography: middle to upper 1/3 of slope (1,1)

Microtopography:

Parent material:

Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 4459 (4459) | 4459-4459 (4459-4459) |
| Slope (degrees) | 15 (15.0) | 15-15 (15-15) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 17.0 (17.0) | 17-17 (17-17) |
| Bare ground cover (%) | 36.0 (36.0) | 36-36 (36-36) |
| Litter cover (%) | 45.0 (45.0) | 45-45 (45-45) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---------------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | |
| <i>Quercus wislizeni</i> | 100 (100) | 28 (28) | 28 (28) | 28 (28) |
| Shrub | | | | |
| <i>Amorpha californica</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>Arctostaphylos glandulosa</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>Clematis lasiantha</i> | 100 (100) | 12 (12) | 12 (12) | 12 (12) |
| <i>Eriophyllum confertiflorum</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>Fraxinus dipetala</i> | 100 (100) | 5 (5) | 5 (5) | 5 (5) |
| <i>Keckiella cordifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Quercus berberidifolia</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Sambucus nigra subsp. caerulea</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| Herb | | | | |
| * <i>Bromus tectorum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Salix gooddingii* Alliance**
***Salix gooddingii* Association**

Samples Used To Describe Association (n= 22)

Rapid Assessment(s), WR and WSD stands: 1186KH01, PRAD.002/8/BPU 1168-31, SDRP0240, SDRP0322, PRAD.003/43/BPU 275531-40, 0623DH02, SDRP0317, SDRP0086, LAMA.020, 0506MJ21, SDRP0077, CORN.002/450, 1633FS01, 0708TK02, 1531EB02, 0709MG01, SDRP0239, 1744MM01, SDRP0238, 0503RB22, SDRP0245
Rapid Assessment(s), OC stands (n=1): 1041EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|----------------|-------------------|
| Total vegetation cover (%) | 63.8 (65.0) | 32-98 (65-65) | - |
| Herb cover (%) | 21.0 (17.0) | 0.2-70 (17-17) | 0-5 (0-0.5) |
| Shrub cover (%) | 9.8 (1.0) | 1-17 (1-1) | 0-10 (0.5-1) |
| Tall shrub / low tree cover (%) | 25.0 (25.0) | 25-25 (25-25) | 1-5 (2-5) |
| Overstory tree cover (%) | 31.2 (20.0) | 7-80 (20-20) | 2-35 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,0), NW (4,0), SW (6,1), (0,0), Flat (7,0), SE (3,0)
Macrotopography: middle 1/3 of slope (1,0), bottom (16,1), bottom to lower 1/3 of slope (1,0), lower 1/3 of slope (3,0)
Microtopography: concave (9,1), Flat (9,0), undulating (3,0)
Parent material: calcareous (1,0), clayey alluvium (1,0), granitic (1,0), gravelly alluvium (1,0), mixed alluvium (5,1), mixed sedimentary (1,0), sandy alluvium (4,0), silty alluvium (6,0)
Soil texture: sandy loam (8,0), sand (2,1), silt (1,0), loam (2,0), silty loam (1,0), clay loam (3,0), clay (2,0), muck (1,0)

| | Average | Range |
|------------------------------|----------------|------------------|
| Elevation (ft) | 366 (167) | 0-1371 (167-167) |
| Slope (degrees) | 1 (1.0) | 0-5 (1-1) |
| Large rock cover (%) | 0.4 (-) | 0-8 (-) |
| Small rock cover (%) | 0.7 (6.0) | 0-7 (6-6) |
| Bare ground cover (%) | 18.1 (42.0) | 0-85 (42-42) |
| Litter cover (%) | 35.5 (45.0) | 0-95 (45-45) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|------------------------------|------------|------------|------------|------------|
| Tree | <i>Salix gooddingii</i> | 100 (100) | 28.3 (20) | 7 (20) | 85 (20) |
| | <i>Salix lasiolepis</i> | 32 (100) | 31.7 (25) | 7 (25) | 65 (25) |
| | <i>Salix laevigata</i> | 27 (-) | 9.7 (-) | 0.2 (-) | 35 (-) |
| | | | | | |
| Shrub | <i>Baccharis salicifolia</i> | 68 (100) | 4.4 (1) | 0.2 (1) | 18 (1) |
| | * <i>Tamarix</i> | 36 (-) | 6.7 (-) | 0.2 (-) | 30 (-) |
| | | | | | |
| Herb | * <i>Hirschfeldia incana</i> | 45 (-) | 1.3 (-) | 0.2 (-) | 5 (-) |
| | * <i>Arundo donax</i> | 27 (100) | 0.9 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Distichlis spicata</i> | 27 (-) | 4.6 (-) | 0.2 (-) | 20 (-) |
| | | | | | |

***Salix laevigata* Alliance**
***Salix laevigata* Association**

Samples Used To Describe Association (n= 16)

Rapid Assessment(s), WR and WSD stands: 1339EB01, SDG0072, 1737KH04, 1187KH01, 1744MB01, 1705KH01, 1782MB01, 1710KH02, 0603JJ01, 1177KH01, 1705MM01, 1737KH03

Rapid Assessment(s), OC stands (n=4): 1069EK03, 1128MA03, 1521EK01, 3.8jd06

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 57.8 (58.0) | 23-85 (47-85) | - |
| Herb cover (%) | 15.3 (20.3) | 3-45 (14-33) | 0-5 (0-1) |
| Shrub cover (%) | 12.0 (12.0) | 1-20 (1-20) | 0-10 (1-5) |
| Tall shrub / low tree cover (%) | 55.0 (55.0) | 55-55 (55-55) | 2-10 (5-10) |
| Overstory tree cover (%) | 31.8 (50.0) | 5-77 (23-77) | 2-15 (2-15) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (7,2), NE (2,0), NW (2,1), S (1,1), SE (1,0), SW (2,0), W (1,0)

Macrotopography: bottom (13,4), bottom to lower 1/3 of slope (3,0)

Microtopography: concave (9,2), flat (5,2), undulating (2,0)

Parent material: granitic (2,0), gravelly alluvium (1,0), mixed alluvium (1,0), sandstone (2,2), sandy alluvium (6,0), sedimentary (1,1)

Soil texture: clay loam (2,1), sand (10,2), sandy loam (4,1)

| | Average | Range |
|------------------------------|----------------|------------------|
| Elevation (ft) | 304 (465) | 56-607 (384-607) |
| Slope (degrees) | 1 (0.5) | 0-3 (0-1) |
| Large rock cover (%) | 0.4 (0.3) | 0-6 (0-1) |
| Small rock cover (%) | 13.0 (1.0) | 0-70 (0-4) |
| Bare ground cover (%) | 27.0 (19.3) | 0-95 (1-32) |
| Litter cover (%) | 53.9 (77.3) | 2-99 (60-99) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|------------|-------------|------------|------------|
| Tree | <i>Salix laevigata</i> | 100 (100) | 26.9 (38.3) | 5 (12) | 77 (77) |
| | <i>Salix lasiolepis</i> | 44 (-) | 12.3 (-) | 2 (-) | 20 (-) |
| | <i>Salix gooddingii</i> | 31 (-) | 1.4 (-) | 0.2 (-) | 4 (-) |
| | | | | | |
| Shrub | <i>Baccharis salicifolia</i> | 63 (100) | 9 (2.8) | 0.2 (0.2) | 40 (7) |
| | <i>Baccharis pilularis</i> | 31 (-) | 2.4 (-) | 0.2 (-) | 6 (-) |
| | <i>Toxicodendron diversilobum</i> | 31 (50) | 3.6 (4) | 1 (1) | 7 (7) |
| | <i>Rhus integrifolia</i> | 25 (-) | 6.1 (-) | 0.2 (-) | 12 (-) |
| | | | | | |
| Herb | <i>Artemisia douglasiana</i> | 38 (50) | 3.4 (6) | 0.2 (2) | 10 (10) |
| | * <i>Helminthotheca echioides</i> | 38 (25) | 1.3 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Ambrosia psilostachya</i> | 31 (25) | 0.9 (2) | 0.2 (2) | 2 (2) |
| | <i>Artemisia palmeri</i> | 31 (-) | 3 (-) | 0.2 (-) | 7 (-) |
| | * <i>Raphanus sativus</i> | 31 (25) | 1.9 (3) | 0.2 (3) | 4 (3) |
| | * <i>Bromus diandrus</i> | 25 (50) | 14.3 (7.6) | 0.2 (0.2) | 38 (15) |
| | * <i>Conium maculatum</i> | 25 (25) | 1.4 (0.2) | 0.2 (0.2) | 4 (0.2) |
| | | | | | |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 39

| Species Name | Con | Ave | Min | Max |
|-----------------------------|--------|---------|---------|-------|
| <i>*Hirschfeldia incana</i> | 25 (-) | 2.6 (-) | 0.2 (-) | 5 (-) |

***Salix laevigata* Alliance**

***Salix laevigata*/*Salix lasiolepis*/*Artemisia douglasiana* Association**

Samples Used To Describe Association (n= 13)

Rapid Assessment(s), WR and WSD stands: SAGE.067, ANZA.120/211, ANZA.008, SMER.048, CLNF.019, SAGE.064, SAGE.034, PALM.001, PALM.003, ANZA.037, IDYL.048/641/BPU#112664/41

Rapid Assessment(s), OC stands (n=2): 1300EK01, 1513EK04

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 64.3 (44.5) | 32-90 (44-45) | - |
| Herb cover (%) | 27.4 (10.5) | 5-55 (10-11) | 0-1 (0-0.5) |
| Shrub cover (%) | 29.2 (5.0) | 4-60 (4-6) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | 15.5 (15.5) | 8-23 (8-23) | 2-10 (2-5) |
| Overstory tree cover (%) | 23.5 (19.5) | 4-55 (10-29) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (3,0), SE (2,0), NE (1,0), N (1,1), SW (6,1)

Macrotopography: bottom (9,1), bottom to lower 1/3 of slope (1,1), lower 1/3 of slope (2,0), middle 1/3 of slope (1,0)

Microtopography: concave (12,2), Convex (1,0)

Parent material: granitic (4,0), mixed alluvium (6,1), sandstone (1,1), sandy alluvium (2,0)

Soil texture: sand (2,2), sandy loam (9,0), loam (1,0), clay loam (1,0)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 2837 (840) | 791-5027 (791-889) |
| Slope (degrees) | 2 (2.0) | 1-5 (2-2) |
| Large rock cover (%) | 6.3 (-) | 0-20 (-) |
| Small rock cover (%) | 5.9 (11.0) | 0-30 (4-18) |
| Bare ground cover (%) | 13.2 (43.5) | 0-45 (42-45) |
| Litter cover (%) | 38.0 (41.5) | 0-90 (35-48) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-------------|-----------|---------|
| Tree | | | | |
| <i>Salix laevigata</i> | 100 (100) | 20.7 (12) | 4 (8) | 50 (16) |
| <i>Salix lasiolepis</i> | 100 (100) | 21.4 (15.5) | 4 (8) | 60 (23) |
| <i>Platanus racemosa</i> | 31 (100) | 3 (2.5) | 1 (2) | 6 (3) |
| <i>Quercus agrifolia</i> | 31 (50) | 3 (10) | 0.2 (10) | 10 (10) |
| Shrub | | | | |
| <i>Salix lasiolepis</i> | 100 (100) | 21.4 (15.5) | 4 (8) | 60 (23) |
| <i>Baccharis salicifolia</i> | 46 (50) | 7 (4) | 0.2 (4) | 15 (4) |
| <i>Artemisia tridentata</i> | 38 (-) | 4.8 (-) | 1 (-) | 18 (-) |
| <i>Toxicodendron diversilobum</i> | 31 (100) | 6 (2.5) | 1 (1) | 15 (4) |
| Herb | | | | |
| <i>Artemisia douglasiana</i> | 62 (100) | 8.6 (2.5) | 1 (1) | 40 (4) |
| <i>Ambrosia psilostachya</i> | 46 (50) | 4.9 (3) | 0.2 (3) | 18 (3) |
| <i>*Hirschfeldia incana</i> | 46 (100) | 0.9 (0.2) | 0.2 (0.2) | 2 (0.2) |



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 40

| Species Name | Con | Ave | Min | Max |
|---------------------------|----------|-----------|-----------|----------|
| <i>*Bromus diandrus</i> | 38 (100) | 3 (3) | 1 (1) | 5 (5) |
| <i>*Melilotus indicus</i> | 31 (100) | 3.6 (0.2) | 0.2 (0.2) | 10 (0.2) |
| <i>Unknown</i> | 31 (-) | 11.1 (-) | 0.2 (-) | 23 (-) |
| <i>Urtica dioica</i> | 31 (50) | 1.3 (1) | 0.2 (1) | 2 (1) |

VEGETATION DOMINATED BY SHRUBS

Acmispon glaber Alliance

Acmispon glaber Association

Samples Used To Describe Association (n= 13)

Rapid Assessment(s), WR and WSD stands: SDRP0193, SDRP0192, SDG0104, SDG0084, SDG0086, 0519DH05, 0423RB02

Rapid Assessment(s), OC stands (n=6): 1197EK03, 1218EK02, 1237MA01, 1289MA01, 1425EK01, khunk01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|---------------|
| Total vegetation cover (%) | 45.2 (53.7) | 8-80 (40-80) | - |
| Herb cover (%) | 12.9 (6.7) | 0.2-49 (0.2-30) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 47.5 (47.5) | 35-80 (35-80) | 0-5 (0.5-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,0), N (1,1), NE (2,1), NW (3,1), SE (1,1), SW (4,1), W (1,1)

Macrotopography: lower to middle 1/3 of slope (2,0), middle 1/3 of slope (5,2), middle 1/3 of slope to ridgetop (1,1), middle to upper 1/3 of slope (3,3), upper 1/3 of slope (1,0), upper 1/3 of slope to ridgetop (1,0)

Microtopography: convex (5,2), flat (1,1), undulating (7,3)

Parent material: granitic (4,0), igneous (2,0), metamorphic (1,0), sandstone (2,2)

Soil texture: clay (1,0), clay loam (2,2), loam (1,0), sandy loam (8,3), silty loam (1,1)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1061 (1016) | 459-1680 (656-1450) |
| Slope (degrees) | 23 (29.2) | 0-40 (5-40) |
| Large rock cover (%) | 0.3 (-) | 0-3 (-) |
| Small rock cover (%) | 2.9 (1.8) | 0-14 (0-5) |
| Bare ground cover (%) | 52.3 (29.6) | 5-88 (5-50) |
| Litter cover (%) | 31.4 (66.0) | 0-90 (47-90) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-------------|-----------|---------|
| Shrub | | | | |
| <i>Acmispon glaber</i> | 100 (100) | 24.6 (35.8) | 4 (25) | 72 (72) |
| <i>Artemisia californica</i> | 77 (100) | 3.1 (4.9) | 0.2 (0.2) | 12 (12) |
| <i>Eriogonum fasciculatum</i> | 69 (50) | 1.7 (2) | 0.2 (1) | 5 (4) |
| <i>Malacothamnus fasciculatus</i> | 46 (67) | 0.9 (0.8) | 0.2 (0.2) | 2 (1) |
| <i>Hazardia squarrosa</i> | 38 (-) | 2.4 (-) | 1 (-) | 6 (-) |
| <i>Salvia apiana</i> | 38 (17) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Salvia mellifera</i> | 38 (67) | 1.1 (1.3) | 0.2 (0.2) | 2 (2) |
| <i>Rhus integrifolia</i> | 31 (67) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 62 (17) | 8.5 (14) | 1 (14) | 23 (14) |
| * <i>Centaurea melitensis</i> | 38 (33) | 6.2 (4.1) | 0.2 (0.2) | 18 (8) |
| * <i>Hirschfeldia incana</i> | 38 (50) | 5.4 (2.7) | 2 (2) | 15 (4) |

Adenostoma fasciculatum Alliance

Adenostoma fasciculatum-Acmispon glaber Association

Samples Used To Describe Association (n= 25)

Rapid Assessment(s), WR and WSD stands: 1277TK01, 1607MB02, SDG0124, 0603DH04, SDG0004, 0608JJ01, 1333EB01, 0506AK22, 0604JS01, SDRP0423, 0310TK22, 0608DH02, 0505JJ02, 0505DH01, 1393KG01, 0504RB01, 1484TK01, 0423DH01, 0506TK01, 1163MB01, 0423DH03, 0421TK04

Rapid Assessment(s), OC stands (n=3): 1343EK03, 1350EK01, 1456EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 33.4 (34.3) | 15-49 (32-38) | - |
| Herb cover (%) | 14.3 (10.0) | 2-38 (10-10) | 0-2 (0-0.5) |
| Shrub cover (%) | 32.7 (32.7) | 32-33 (32-33) | 0-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description All stands, OC stands

Aspect: SE (2,0), SW (8,1), NW (6,1), NE (4,0), Flat (1,0), E (1,1), Variable (3,0)

Macrotopography: middle to upper 1/3 of slope (2,0), upper 1/3 of slope to ridgetop (4,0), ridge top (1,0), middle 1/3 of slope to ridgetop (2,0), lower to upper 1/3 of slope (4,0), entire slope (1,0), lower to middle 1/3 of slope (4,1), upper 1/3 of slope (2,1), lower 1/3 of slope to ridgetop (1,0), middle 1/3 of slope (3,1)

Microtopography: undulating (14,2), concave (4,1), convex (6,0), flat (1,0)

Parent material: gabbro (1,0), granitic (14,0), metamorphic (2,0), mixed alluvium (1,1), sandstone (3,2)

Soil texture: loam (1,0), sand (2,2), clay loam (2,0), clay (1,0), sandy loam (18,1)

| | Average | Range |
|------------------------------|-------------|----------------------|
| Elevation (ft) | 1515 (1562) | 223-2526 (1093-2034) |
| Slope (degrees) | 16 (12.3) | 0-35 (9-18) |
| Large rock cover (%) | 8.8 (18.3) | 0-40 (0-40) |
| Small rock cover (%) | 31.1 (17.3) | 0-69 (7-35) |
| Bare ground cover (%) | 36.1 (39.7) | 0-65 (23-58) |
| Litter cover (%) | 17.8 (22.7) | 0-71 (10-33) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|----------|
| Shrub | | | | |
| <i>Adenostoma fasciculatum</i> | 100 (100) | 11.7 (19) | 3 (14) | 24 (23) |
| <i>Malosma laurina</i> | 84 (67) | 2.3 (0.2) | 0.2 (0.2) | 12 (0.2) |
| <i>Acmispon glaber</i> | 72 (-) | 5.1 (-) | 0.2 (-) | 19 (-) |
| <i>Helianthemum scoparium</i> | 56 (100) | 6.4 (6.7) | 0.2 (2) | 16 (10) |
| <i>Hesperoyucca whipplei</i> | 56 (100) | 0.3 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Eriogonum fasciculatum</i> | 52 (-) | 2.1 (-) | 0.2 (-) | 9 (-) |
| <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 44 (-) | 0.9 (-) | 0.2 (-) | 2 (-) |
| <i>Salvia apiana</i> | 40 (-) | 0.7 (-) | 0.2 (-) | 3 (-) |
| <i>Xylococcus bicolor</i> | 36 (-) | 1.3 (-) | 0.2 (-) | 3 (-) |
| <i>Artemisia californica</i> | 32 (-) | 0.9 (-) | 0.2 (-) | 6 (-) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 52 (-) | 3.3 (-) | 0.2 (-) | 8 (-) |
| * <i>Festuca myuros</i> | 52 (-) | 3.6 (-) | 0.2 (-) | 19 (-) |
| * <i>Centaurea melitensis</i> | 44 (33) | 2.5 (1) | 0.2 (1) | 15 (1) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 43

| Species Name | Con | Ave | Min | Max |
|-----------------------------|---------|---------|---------|--------|
| * <i>Hypochaeris glabra</i> | 40 (33) | 2.2 (3) | 0.2 (3) | 13 (3) |
| * <i>Avena barbata</i> | 28 (33) | 1.3 (7) | 0.2 (7) | 7 (7) |

***Adenostoma fasciculatum* Alliance**

***Adenostoma fasciculatum*-*Ceanothus crassifolius* Association**

Samples Used To Describe Association (n= 10)

Rapid Assessment(s), WR and WSD stands: 0610RB02, 0610DH02, 0610AK03, 0609TK02, 0609RB04, 0506AK21

Rapid Assessment(s), OC stands (n=4): 1176EK01, 1324EK01, 1324EK02, 1521EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|---------------|
| Total vegetation cover (%) | 47.0 (35.0) | 28-65 (28-40) | - |
| Herb cover (%) | 7.4 (1.8) | 0.2-52 (0.2-5) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 33.8 (33.8) | 23-40 (23-40) | 0.5-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 0.2 (-) | 0.2-0.2 (-) | 10-15 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,0), SW (5,2), Flat (1,1), S (1,1), SE (2,0)

Macrotopography: middle to upper 1/3 of slope (2,0), upper 1/3 of slope (1,0), ridge top (1,0), lower to upper 1/3 of slope (1,0), lower to middle 1/3 of slope (1,0), middle 1/3 of slope (4,4)

Microtopography: undulating (6,2), convex (3,1), flat (1,1)

Parent material: granitic (2,0), metamorphic (4,0), sandstone (4,4)

Soil texture: clay loam (3,0), sand (1,1), sandy loam (4,1), silty loam (1,1)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1993 (1546) | 656-2654 (656-2461) |
| Slope (degrees) | 18 (15.8) | 0-38 (0-38) |
| Large rock cover (%) | 4.8 (3.3) | 0-19 (0-10) |
| Small rock cover (%) | 15.0 (9.3) | 0-32 (0-16) |
| Bare ground cover (%) | 26.9 (48.3) | 6-51 (46-51) |
| Litter cover (%) | 51.4 (37.3) | 25-80 (25-52) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|---------|
| Shrub | | | | |
| <i>Adenostoma fasciculatum</i> | 100 (100) | 28 (20.8) | 9 (9) | 52 (26) |
| <i>Ceanothus crassifolius</i> | 100 (100) | 9.5 (6) | 0.2 (2) | 25 (10) |
| <i>Hesperoyucca whipplei</i> | 100 (100) | 0.7 (0.7) | 0.2 (0.2) | 2 (2) |
| <i>Quercus berberidifolia</i> | 70 (25) | 3.8 (8) | 0.2 (8) | 8 (8) |
| <i>Rhamnus ilicifolia</i> | 60 (-) | 1 (-) | 0.2 (-) | 3 (-) |
| <i>Arctostaphylos glandulosa</i> | 50 (25) | 1.4 (0.2) | 0.2 (0.2) | 6 (0.2) |
| <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 50 (-) | 0.9 (-) | 0.2 (-) | 3 (-) |
| <i>Heteromeles arbutifolia</i> | 50 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Malosma laurina</i> | 50 (75) | 0.9 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Salvia mellifera</i> | 50 (75) | 2.2 (1) | 1 (1) | 5 (1) |
| <i>Mimulus aurantiacus</i> | 40 (-) | 1.2 (-) | 0.2 (-) | 4 (-) |

| | Species Name | Con | Ave | Min | Max |
|------|-------------------------------|---------|-----------|-----------|-----------|
| Herb | <i>Eriogonum fasciculatum</i> | 30 (50) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Stipa lepida</i> | 30 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Adenostoma fasciculatum-Arctostaphylos glandulosa Alliance
Adenostoma fasciculatum-Arctostaphylos glandulosa Association

Samples Used To Describe Association (n= 10)

Rapid Assessment(s), WR and WSD stands: CAHUI.003/606??, CLNF.113, CLNF.121/439, CLNF.097

Rapid Assessment(s), OC stands (n=6): 1180EK03, 1184JD02, 1209EK02, 1351EK02, 1351EK03, 1380EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|---------------|
| Total vegetation cover (%) | 53.2 (50.3) | 35-75 (35-75) | - |
| Herb cover (%) | 0.1 (0.2) | 0-0.2 (0.2-0.2) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 48.4 (39.8) | 35-75 (35-50) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | 3.0 (3.0) | 3-3 (3-3) | 0.5-2 (0.5-1) |
| Overstory tree cover (%) | 2.1 (0.2) | 0.2-6 (0.2-0.2) | 2-5 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: N (2,2), NE (3,1), SE (5,3)

Macrotopography: upper 1/3 of slope (7,5), middle to upper 1/3 of slope (1,1), ridge top (2,0)

Microtopography: undulating (4,3), flat (1,1), concave (1,1), Convex (4,1)

Parent material: granitic (1,0), metamorphic (3,2), sandstone (3,3), sedimentary (2,0)

Soil texture: clay loam (3,2), loam (1,0), sand (2,2), sandy loam (4,2)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3788 (3476) | 2211-4853 (2211-4495) |
| Slope (degrees) | 26 (25.5) | 15-38 (15-38) |
| Large rock cover (%) | 3.2 (0.2) | 0-15 (0-1) |
| Small rock cover (%) | 26.1 (21.3) | 5-70 (5-38) |
| Bare ground cover (%) | 24.3 (26.2) | 1-55 (2-48) |
| Litter cover (%) | 43.3 (49.7) | 2-90 (28-90) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-------------|-----------|-----------|
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 26.6 (27.3) | 8 (8) | 49 (49) |
| | <i>Arctostaphylos glandulosa</i> | 100 (100) | 19.5 (11) | 6 (6) | 45 (16) |
| | <i>Dendromecon rigida</i> | 50 (83) | 0.7 (0.7) | 0.2 (0.2) | 2 (2) |
| | <i>Hesperoyucca whipplei</i> | 40 (50) | 0.4 (0.5) | 0.2 (0.2) | 1 (1) |
| | <i>Quercus berberidifolia</i> | 40 (67) | 5 (5) | 3 (3) | 8 (8) |
| | <i>Yucca whipplei</i> | 40 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |
| | <i>Helianthemum scoparium</i> | 30 (33) | 6.7 (9.5) | 1 (6) | 13 (13) |
| | <i>Heteromeles arbutifolia</i> | 30 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Mimulus aurantiacus</i> | 30 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Adenostoma fasciculatum*-*Arctostaphylos glandulosa* Alliance**

***Adenostoma fasciculatum*-*Arctostaphylos glandulosa*-*Ceanothus crassifolius* Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: VAIL.003/2047, CLNF.064, CLNF.007

Rapid Assessment(s), OC stands (n=1): 1294EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 60.5 (50.0) | 50-72 (50-50) | - |
| Herb cover (%) | 1.0 (-) | 0-2 (-) | 0-0.5 (-) |
| Shrub cover (%) | 60.5 (50.0) | 50-72 (50-50) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1), NW (2,0), Variable (1,0)

Macrotopography: middle 1/3 of slope (1,0), upper 1/3 of slope (3,1)

Microtopography: Convex (3,1), Undulating (1,0)

Parent material: granitic (2,0), sandstone (1,1), sedimentary (1,0)

Soil texture: clay loam (1,0), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3270 (3366) | 2878-3650 (3366-3366) |
| Slope (degrees) | 24 (20.0) | 4-38 (20-20) |
| Large rock cover (%) | 1.3 (-) | 0-3 (-) |
| Small rock cover (%) | 28.8 (15.0) | 15-50 (15-15) |
| Bare ground cover (%) | 27.0 (42.0) | 1-45 (42-42) |
| Litter cover (%) | 38.3 (40.0) | 20-68 (40-40) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus wislizeni</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| Shrub | | | | | |
| | <i>Adenostoma fasciculatum</i> | 100 (100) | 18.8 (30) | 9 (30) | 30 (30) |
| | <i>Arctostaphylos glandulosa</i> | 100 (100) | 21.5 (12) | 5 (12) | 39 (12) |
| | <i>Ceanothus crassifolius</i> | 100 (100) | 21.3 (5) | 5 (5) | 38 (5) |
| | <i>Quercus berberidifolia</i> | 50 (100) | 4 (1) | 1 (1) | 7 (1) |
| | <i>Ceanothus leucodermis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Dendromecon rigida</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Helianthemum scoparium</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| | <i>Yucca whipplei</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | | | | | |
| Herb | | | | | |
| | <i>Camissonia</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Galium angustifolium</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Plagiobothrys</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |

***Adenostoma fasciculatum*-*Arctostaphylos glauca* Alliance**
***Adenostoma fasciculatum*-*Arctostaphylos glauca* Association**

Samples Used To Describe Association (n= 7)

Rapid Assessment(s), WR and WSD stands: BLACK.038/1292, BLACK.054/BPU 107-87, BLACK.058/BPU 1037953-292, BLACK.049/475, BLACK.043/526, SAJAC.010/BPU 11809-121

Rapid Assessment(s), OC stands (n=1): 1180EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|------------------|-------------------|
| Total vegetation cover (%) | 65.4 (72.0) | 55-75 (72-72) | - |
| Herb cover (%) | 9.7 (0.2) | 0.2-25 (0.2-0.2) | 0-1 (0-0.5) |
| Shrub cover (%) | 58.9 (72.0) | 45-72 (72-72) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 2-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (3,0), NW (2,0), SW (2,1)

Macrotopography: lower 1/3 of slope (1,0), middle 1/3 of slope (2,0), upper 1/3 of slope (3,0), upper 1/3 of slope to ridgetop (1,1)

Microtopography: Undulating (7,1)

Parent material: gneiss/biotite gneiss (1,0), granitic (2,0), metamorphic (1,1), sedimentary (2,0)

Soil texture: sand (1,0), sandy loam (4,0)

| | Average | Range |
|------------------------------|----------------|-----------------------|
| Elevation (ft) | 2721 (3596) | 2224-3596 (3596-3596) |
| Slope (degrees) | 25 (40.0) | 14-40 (40-40) |
| Large rock cover (%) | 1.6 (-) | 0-8 (-) |
| Small rock cover (%) | 12.4 (10.0) | 0-38 (10-10) |
| Bare ground cover (%) | 23.3 (17.0) | 0-50 (17-17) |
| Litter cover (%) | 44.7 (70.0) | 0-75 (70-70) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|------------|------------|------------|------------|
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 27.3 (37) | 10 (37) | 45 (37) |
| | <i>Arctostaphylos glauca</i> | 100 (100) | 14.7 (15) | 3 (15) | 33 (15) |
| | <i>Rhus ovata</i> | 100 (100) | 6.3 (1) | 1 (1) | 11 (1) |
| | <i>Salvia mellifera</i> | 71 (100) | 2.6 (1) | 0.2 (1) | 5 (1) |
| | <i>Hesperoyucca whipplei</i> | 57 (100) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Yucca whipplei</i> | 57 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |
| | <i>Eriogonum fasciculatum</i> | 43 (-) | 1.7 (-) | 0.2 (-) | 3 (-) |
| | <i>Heteromeles arbutifolia</i> | 43 (100) | 1.4 (2) | 0.2 (2) | 2 (2) |
| | <i>Quercus berberidifolia</i> | 43 (100) | 1.7 (2) | 1 (2) | 2 (2) |
| | <i>Keckiella antirrhinoides</i> | 29 (-) | 3.5 (-) | 3 (-) | 4 (-) |
| | <i>Rhamnus crocea</i> | 29 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| Herb | <i>*Bromus madritensis</i> | 57 (-) | 3 (-) | 2 (-) | 4 (-) |
| | <i>Cryptantha muricata</i> | 29 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| | <i>Eucrypta chrysanthemifolia</i> | 29 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| | <i>Pterostegia drymarioides</i> | 29 (-) | 7.5 (-) | 7 (-) | 8 (-) |

***Adenostoma fasciculatum-Salvia mellifera* Alliance**
***Adenostoma fasciculatum-Salvia mellifera* Association**

Samples Used To Describe Association (n= 9)

Rapid Assessment(s), WR and WSD stands: SAGE.050, SAGE.056, SATI.010

Rapid Assessment(s), OC stands (n=6): 1019EK01, 1024EK01, 1092EK05, 1229EK02, 1343EK02, 1351EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 45.4 (39.5) | 35-64 (35-50) | - |
| Herb cover (%) | 0.7 (0.6) | 0-2 (0.2-2) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 45.0 (38.8) | 35-65 (35-48) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NE (1,0), SE (4,3), SW (1,0), Variable (1,1), W (1,1)

Macrotopography: lower 1/3 of slope (2,1), middle 1/3 of slope (2,2), middle to upper 1/3 of slope (1,1), upper 1/3 of slope (4,2)

Microtopography: Concave (3,1), flat (3,3), undulating (3,2)

Parent material: granitic (1,0), mixed igneous (1,0), sandstone (6,6), sedimentary (1,0)

Soil texture: loam (3,1), sandy loam (6,5)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1568 (1333) | 623-2457 (623-2457) |
| Slope (degrees) | 18 (20.3) | 5-40 (5-40) |
| Large rock cover (%) | 0.2 (0.3) | 0-2 (0-2) |
| Small rock cover (%) | 6.0 (8.2) | 0-25 (0-25) |
| Bare ground cover (%) | 36.2 (48.5) | 0-66 (29-66) |
| Litter cover (%) | 32.6 (40.5) | 0-68 (5-68) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---------------------------------|-----------|-------------|-----------|---------|
| Shrub | | | | |
| <i>Adenostoma fasciculatum</i> | 100 (100) | 31.1 (30.3) | 24 (24) | 48 (48) |
| <i>Salvia mellifera</i> | 100 (100) | 11.8 (4) | 2 (2) | 30 (6) |
| <i>Hesperoyucca whipplei</i> | 56 (50) | 0.4 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Malosma laurina</i> | 56 (83) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Yucca whipplei</i> | 56 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |
| <i>Helianthemum scoparium</i> | 44 (67) | 1.1 (1.1) | 0.2 (0.2) | 3 (3) |
| <i>Rhus integrifolia</i> | 44 (67) | 1.6 (1.6) | 0.2 (0.2) | 3 (3) |
| <i>Eriodictyon crassifolium</i> | 33 (33) | 2.1 (3) | 0.2 (2) | 4 (4) |

***Adenostoma fasciculatum-Salvia mellifera* Alliance**
***Adenostoma fasciculatum-Salvia mellifera-Ceanothus crassifolius* Association**

Samples Used To Describe Association (n= 9)

Rapid Assessment(s), WR and WSD stands: SMER.033/scrub1.02, SITT.013, RECA.007, RECA.018, CLNF.117/221, SAGE.070, MURR.007, VAIL.008/1809

Rapid Assessment(s), OC stands (n=1): 1161MA01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 63.0 (72.0) | 38-85 (72-72) | - |
| Herb cover (%) | 17.1 (3.0) | 0.2-75 (3-3) | 0-1 (0.5-1) |
| Shrub cover (%) | 52.7 (72.0) | 35-72 (72-72) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Variable (1,0), NE (5,0), SE (2,1), SW (1,0)

Macrotopography: middle to upper 1/3 of slope (1,1), upper 1/3 of slope (5,0), middle 1/3 of slope (1,0), lower 1/3 of slope (1,0), bottom (1,0)

Microtopography: Flat (1,0), Undulating (8,1)

Parent material: conglomerate (1,0), granitic (7,1), mix of two or more rock types (1,0)

Soil texture: loam (2,0), sandy loam (5,0), clay loam (2,1)

| | Average | Range |
|------------------------------|----------------|----------------------|
| Elevation (ft) | 2272 (2628) | 828-2901 (2628-2628) |
| Slope (degrees) | 19 (12.0) | 10-32 (12-12) |
| Large rock cover (%) | 9.8 (-) | 0-30 (-) |
| Small rock cover (%) | 21.1 (25.0) | 0-55 (25-25) |
| Bare ground cover (%) | 27.8 (25.0) | 0-60 (25-25) |
| Litter cover (%) | 27.1 (48.0) | 0-48 (48-48) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--------------------------------|------------|------------|------------|------------|
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 27.2 (55) | 14 (55) | 55 (55) |
| | <i>Ceanothus crassifolius</i> | 100 (100) | 4.5 (2) | 0.2 (2) | 18 (2) |
| | <i>Salvia mellifera</i> | 100 (100) | 17.9 (10) | 10 (10) | 30 (10) |
| | <i>Hesperoyucca whipplei</i> | 56 (100) | 0.5 (1) | 0.2 (1) | 1 (1) |
| | <i>Yucca whipplei</i> | 56 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| | <i>Eriogonum fasciculatum</i> | 44 (100) | 5.5 (0.2) | 3 (0.2) | 10 (0.2) |
| | <i>Rhus ovata</i> | 44 (100) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Helianthemum scoparium</i> | 33 (-) | 2.4 (-) | 0.2 (-) | 4 (-) |
| Herb | <i>*Bromus madritensis</i> | 67 (100) | 5 (1) | 1 (1) | 15 (1) |
| | <i>Cryptantha intermedia</i> | 44 (-) | 2.3 (-) | 0.2 (-) | 5 (-) |
| | <i>Dichelostemma capitatum</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Arctostaphylos glandulosa* Alliance**
***Arctostaphylos glandulosa* Association**

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: 1827MM02

Rapid Assessment(s), OC stands (n=2): 1180EK02, 1209JD01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 68.3 (70.0) | 55-85 (55-85) | - |
| Herb cover (%) | 1.0 (-) | 1-1 (-) | 0-0.5 (-) |
| Shrub cover (%) | 50.0 (50.0) | 50-50 (50-50) | 1-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0-0.5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NW (2,1)

Macrotopography: upper 1/3 of slope (3,2)

Microtopography: concave (1,1), convex (1,1), flat (1,0)

Parent material: metamorphic (1,1)

Soil texture: clay loam (2,1), sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3592 (4057) | 2661-4085 (4029-4085) |
| Slope (degrees) | 20 (22.5) | 15-30 (15-30) |
| Large rock cover (%) | 5.0 (-) | 0-15 (-) |
| Small rock cover (%) | 25.0 (15.0) | 10-45 (10-20) |
| Bare ground cover (%) | 10.0 (15.0) | 0-17 (13-17) |
| Litter cover (%) | 58.3 (67.5) | 40-70 (65-70) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|----------|-------------|-----------|-----------|
| Tree | | | | |
| <i>Quercus wislizeni</i> | 67 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Callitropsis forbesii</i> | 33 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Pinus coulteri</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | |
| <i>Adenostoma fasciculatum</i> | 67 (100) | 3.6 (3.6) | 0.2 (0.2) | 7 (7) |
| <i>Arctostaphylos glandulosa</i> | 67 (100) | 61.5 (61.5) | 53 (53) | 70 (70) |
| <i>Garrya flavescens</i> | 67 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Quercus berberidifolia</i> | 67 (100) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Rhus ovata</i> | 67 (50) | 1.1 (0.2) | 0.2 (0.2) | 2 (0.2) |
| <i>Acmispon glaber</i> var. <i>glaber</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Arctostaphylos glandulosa</i> subsp. <i>glandulosa</i> | 33 (-) | 20 (-) | 20 (-) | 20 (-) |
| <i>Ceanothus crassifolius</i> | 33 (50) | 2 (2) | 2 (2) | 2 (2) |
| <i>Ceanothus oliganthus</i> | 33 (50) | 2 (2) | 2 (2) | 2 (2) |
| <i>Ceanothus tomentosus</i> | 33 (-) | 30 (-) | 30 (-) | 30 (-) |
| <i>Comarostaphylis diversifolia</i> subsp. <i>diversifolia</i> | 33 (-) | 25 (-) | 25 (-) | 25 (-) |
| <i>Dendromecon rigida</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|--|---------|-----------|-----------|-----------|
| <i>Ericameria cuneata</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Eriogonum fasciculatum</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 33 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Hazardia squarrosa</i> var. <i>grindelioides</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Helianthemum scoparium</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Hesperoyucca whipplei</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lonicera subspicata</i> var. <i>denudata</i> | 33 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Mimulus aurantiacus</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Quercus</i> | 33 (-) | 10 (-) | 10 (-) | 10 (-) |
| <i>Rhamnus ilicifolia</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Salvia mellifera</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>Galium angustifolium</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Artemisia californica* Alliance** ***Artemisia californica* Association**

Samples Used To Describe Association (n= 33)

Rapid Assessment(s), WR and WSD stands: 0708AK02, 1701JD01, 0625DH01, 1705JD02, 1634JD02, 0707RB03, SDRP0236, 1627MB02, SDRP0087, SDRP0280, SDRP0139, SDRP0138, 0709MG03, 0709DH01

Rapid Assessment(s), OC stands (n=19): 1003EK04, 1004EK01, 1015EK02, 1047EK03, 1093EK01, 1114EK01, 1137EK01, 1187EK02, 1204EK01, 1223EK01, 1366EK01, 1408EK01, 1419EK01, 1419EK02, 1472EK03, 1478EK01, 1494EK02, a1069EK01, ek3.8.1

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 48.2 (50.7) | 25-70 (35-70) | - |
| Herb cover (%) | 12.0 (11.8) | 0.2-45 (1-45) | 0-2 (0-1) |
| Shrub cover (%) | 43.6 (43.6) | 20-65 (20-65) | 0-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: E (3,3), SE (5,2), W (1,1), Variable (2,2), SW (6,3), NW (6,2), NE (6,2), N (1,1), S (3,3)

Macrotopography: middle to upper 1/3 of slope (7,4), upper 1/3 of slope to ridgetop (1,1), upper 1/3 of slope (9,8), entire slope (1,0), ridge top (1,0), middle 1/3 of slope (4,4), lower to upper 1/3 of slope (1,0), lower 1/3 of slope (2,0), lower to middle 1/3 of slope (3,1), middle 1/3 of slope to ridgetop (3,0)

Microtopography: flat (9,5), convex (7,5), concave (5,3), undulating (12,6)

Parent material: granitic (3,0), metamorphic (2,0), sandstone (20,17), sedimentary (1,0), semi-schist (1,1)

Soil texture: sandy loam (12,6), sand (2,2), clay (2,1), clay loam (11,7), silt (1,0), loam (2,1)

| | Average | Range |
|------------------------------|-------------|------------------|
| Elevation (ft) | 530 (630) | 0-1404 (75-1404) |
| Slope (degrees) | 15 (16.7) | 1-40 (1-40) |
| Large rock cover (%) | 0.2 (0.1) | 0-5 (0-1) |
| Small rock cover (%) | 8.6 (5.7) | 0-55 (0-55) |
| Bare ground cover (%) | 39.9 (43.3) | 0-74 (17-62) |
| Litter cover (%) | 35.9 (48.8) | 0-75 (25-70) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------------|-----------|-------------|-----------|---------|
| Shrub | <i>Artemisia californica</i> | 100 (100) | 31.2 (34.2) | 10 (10) | 55 (55) |
| | <i>Eriogonum fasciculatum</i> | 64 (58) | 2.2 (2.7) | 0.2 (0.2) | 11 (11) |
| | <i>Rhus integrifolia</i> | 61 (68) | 3.2 (2.8) | 0.2 (0.2) | 17 (10) |
| | <i>Baccharis pilularis</i> | 55 (68) | 2.7 (2.4) | 0.2 (0.2) | 11 (10) |
| | <i>Malosma laurina</i> | 42 (37) | 3.8 (4) | 0.2 (0.2) | 12 (12) |
| | <i>Mimulus aurantiacus</i> | 42 (53) | 0.6 (0.7) | 0.2 (0.2) | 2 (2) |
| | <i>Acmispon glaber</i> | 36 (37) | 1.7 (2.7) | 0.2 (0.2) | 15 (15) |
| | <i>Opuntia littoralis</i> | 36 (47) | 0.7 (0.4) | 0.2 (0.2) | 4 (1) |
| | | | | | |
| Herb | <i>Pseudognaphalium californicum</i> | 48 (58) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| | * <i>Bromus madritensis</i> | 36 (26) | 3.9 (0.5) | 0.2 (0.2) | 35 (1) |
| | * <i>Brassica nigra</i> | 33 (53) | 0.9 (0.5) | 0.2 (0.2) | 5 (2) |
| | * <i>Centaurea melitensis</i> | 30 (37) | 2.4 (1.9) | 0.2 (0.2) | 10 (10) |

Artemisia californica Alliance

Artemisia californica-*Eriogonum fasciculatum*-*Opuntia littoralis* Provisional Association Association

Samples Used To Describe Association (n= 18)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=18): 1017EK01, 1032EK02, 1056EK07, 1057EK04, 1064EK01, 1066EK01, 1066EK02, 1066EK06, 1128EK02, 1128EK03, 1128EK04, 1197EK02, 1344EK01, 1410EK01, 1450EK03, 1463EK01, 1513EK01, 3.8.2

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|---------------|
| Total vegetation cover (%) | 42.3 (42.3) | 31-60 (31-60) | - |
| Herb cover (%) | 4.4 (4.4) | 0.2-18 (0.2-18) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 39.7 (39.7) | 25-60 (25-60) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1), S (7,7), SE (8,8), SW (1,1), W (1,1)

Macrotopography: lower 1/3 of slope (1,1), lower to middle 1/3 of slope (1,1), middle 1/3 of slope (5,5), middle to upper 1/3 of slope (6,6), upper 1/3 of slope (3,3), upper 1/3 of slope to ridgetop (2,2)

Microtopography: convex (2,2), flat (4,4), concave (3,3), undulating (9,9)

Parent material: mixed alluvium (3,3), sandstone (11,11)

Soil texture: clay loam (2,2), silty loam (1,1), sandy loam (10,10), loam (1,1), clay (1,1), sand (2,2)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 693 (693) | 148-1027 (148-1027) |
| Slope (degrees) | 22 (21.9) | 3-50 (3-50) |
| Large rock cover (%) | 0.8 (0.8) | 0-5 (0-5) |
| Small rock cover (%) | 9.4 (9.4) | 0-42 (0-42) |
| Bare ground cover (%) | 50.6 (50.6) | 31-63 (31-63) |
| Litter cover (%) | 37.1 (37.1) | 15-55 (15-55) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---------------------------------------|-----------|-----------|-----------|-----------|
| Shrub | <i>Artemisia californica</i> | 100 (100) | 19 (19) | 1 (1) | 37 (37) |
| | <i>Eriogonum fasciculatum</i> | 100 (100) | 7.1 (7.1) | 0.2 (0.2) | 20 (20) |
| | <i>Opuntia littoralis</i> | 100 (100) | 7.2 (7.2) | 1 (1) | 20 (20) |
| | <i>Malosma laurina</i> | 83 (83) | 1 (1) | 0.2 (0.2) | 5 (5) |
| | <i>Acmispon glaber</i> | 78 (78) | 2.3 (2.3) | 0.2 (0.2) | 12 (12) |
| | <i>Rhus integrifolia</i> | 72 (72) | 2 (2) | 0.2 (0.2) | 7 (7) |
| | <i>Cuscuta occidentalis</i> | 50 (50) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| | <i>Mimulus aurantiacus</i> | 50 (50) | 1.4 (1.4) | 0.2 (0.2) | 7 (7) |
| | <i>Salvia mellifera</i> | 33 (33) | 0.5 (0.5) | 0.2 (0.2) | 2 (2) |
| | <i>Baccharis pilularis</i> | 28 (28) | 0.8 (0.8) | 0.2 (0.2) | 3 (3) |
| | <i>Brickellia californica</i> | 28 (28) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cylindropuntia prolifera</i> | 28 (28) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhamnus ilicifolia</i> | 28 (28) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Sambucus nigra subsp. caerulea</i> | 28 (28) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| Herb | <i>Pseudognaphalium californicum</i> | 56 (56) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Bromus madritensis</i> | 44 (44) | 0.8 (0.8) | 0.2 (0.2) | 3 (3) |
| | <i>Stipa pulchra</i> | 39 (39) | 0.3 (0.3) | 0.2 (0.2) | 1 (1) |
| | * <i>Centaurea melitensis</i> | 33 (33) | 1.6 (1.6) | 0.2 (0.2) | 8 (8) |
| | <i>Dudleya pulverulenta</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Avena barbata</i> | 28 (28) | 0.7 (0.7) | 0.2 (0.2) | 1 (1) |
| | <i>Scrophularia californica</i> | 28 (28) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Artemisia californica Alliance

Artemisia californica-*Mimulus aurantiacus*-*Elymus condensatus* Provisional Association

Samples Used To Describe Association (n= 11)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=11): 1010EK01, 1010EK03, 1041EK03, 1045JD01, 1045JD02, 1076MA02, 1079EK02, 1089EK02, 1137MA01, 1187EK07, 1413EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 59.5 (59.5) | 50-75 (50-75) | - |
| Herb cover (%) | 6.1 (6.1) | 1-20 (1-20) | 0-2 (0-2) |
| Shrub cover (%) | 54.4 (54.4) | 42-60 (42-60) | 0.5-5 (0.5-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 53

Local Environmental Description (All stands, OC stands)

Aspect: NW (3,3), E (1,1), N (1,1), NE (6,6)

Macrotopography: middle 1/3 of slope (2,2), middle to upper 1/3 of slope (1,1), lower to middle 1/3 of slope (3,3), upper 1/3 of slope (2,2), lower to upper 1/3 of slope (3,3)

Microtopography: convex (2,2), flat (3,3), undulating (5,5)

Parent material: sandstone (8,8), sedimentary (2,2)

Soil texture: clay loam (1,1), loam (2,2), sandy loam (2,2)

| | Average | Range |
|-----------------------|-------------|-------------------|
| Elevation (ft) | 483 (483) | 59-1240 (59-1240) |
| Slope (degrees) | 30 (29.5) | 15-52 (15-52) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 33.2 (33.2) | 5-70 (5-70) |
| Litter cover (%) | 64.9 (64.9) | 30-94 (30-94) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---------------------------------------|-----------|-------------|-----------|-----------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 21.5 (21.5) | 4 (4) | 40 (40) |
| <i>Mimulus aurantiacus</i> | 100 (100) | 14.1 (14.1) | 0.2 (0.2) | 33 (33) |
| <i>Malosma laurina</i> | 82 (82) | 15.2 (15.2) | 0.2 (0.2) | 45 (45) |
| <i>Baccharis pilularis</i> | 55 (55) | 2.9 (2.9) | 0.2 (0.2) | 8 (8) |
| <i>Sambucus nigra subsp. caerulea</i> | 55 (64) | 0.5 (0.5) | 0.2 (0.2) | 2 (2) |
| <i>Heteromeles arbutifolia</i> | 45 (45) | 2.9 (2.9) | 0.2 (0.2) | 9 (9) |
| <i>Rhus integrifolia</i> | 45 (45) | 4 (4) | 0.2 (0.2) | 8 (8) |
| <i>Toxicodendron diversilobum</i> | 36 (36) | 6.8 (6.8) | 3 (3) | 12 (12) |
| <i>Salvia mellifera</i> | 27 (27) | 2.3 (2.3) | 2 (2) | 3 (3) |
| Herb | | | | |
| <i>Elymus condensatus</i> | 82 (82) | 6.1 (6.1) | 0.2 (0.2) | 20 (20) |
| <i>Pseudognaphalium californicum</i> | 27 (27) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Artemisia californica* Alliance**

***Artemisia californica*-*Nolina cismontana* Provisional Association**

Samples Used To Describe Association (n= 2)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=2): 1187EK01, 1187EK06

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 45.0 (45.0) | 43-47 (43-47) | - |
| Herb cover (%) | 0.6 (0.6) | 0.2-1 (0.2-1) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 45.0 (45.0) | 43-47 (43-47) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (1,1), W (1,1)

Macrotopography: middle 1/3 of slope (1,1), middle to upper 1/3 of slope (1,1)

Microtopography: concave (2,2)

Parent material: sandstone (2,2)

Soil texture: sandy loam (1,1)

| | Average | Range |
|------------------------------|----------------|-----------------------|
| Elevation (ft) | 1163 (1163) | 1161-1165 (1161-1165) |
| Slope (degrees) | 38 (37.5) | 35-40 (35-40) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 51.5 (51.5) | 45-58 (45-58) |
| Litter cover (%) | 46.0 (46.0) | 40-52 (40-52) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|------------|------------|------------|------------|
| Shrub | | | | | |
| | <i>Artemisia californica</i> | 100 (100) | 20 (20) | 17 (17) | 23 (23) |
| | <i>Malosma laurina</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Nolina cismontana</i> | 100 (100) | 16 (16) | 15 (15) | 17 (17) |
| | <i>Rhus integrifolia</i> | 100 (100) | 2 (2) | 1 (1) | 3 (3) |
| | <i>Salvia mellifera</i> | 100 (100) | 4.5 (4.5) | 4 (4) | 5 (5) |
| | <i>Acmispon glaber</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Encelia californica</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Eriophyllum confertiflorum</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Hesperoyucca whipplei</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Mirabilis laevis</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Quercus berberidifolia</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Salvia apiana</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | | |
| | <i>*Centaurea melitensis</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Chamaesyce polycarpa</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Linanthus</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Pseudognaphalium biolettii</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Artemisia californica* Alliance**

***Artemisia californica*-*Salvia apiana* Association**

Samples Used To Describe Association (n= 17)

Rapid Assessment(s), WR and WSD stands: SMER.024, SMER.029/435/scrub1.06, SMER.011, SMER.018/Scrub1.20, SMER.027, SAAN.018, SMER.028, SMER.044/scrub1.19, WILD.013, SAAN.014, WILD.014, SAAN.013, SMER.003

Rapid Assessment(s), OC stands (n=4): 1187EK05, 1213EK01, 1340EK01, 1350EK02

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 55

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 58.9 (43.0) | 38-90 (38-46) | - |
| Herb cover (%) | 16.2 (7.8) | 0-60 (2-15) | 0-1 (0-0.5) |
| Shrub cover (%) | 46.5 (37.5) | 33-75 (36-38) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-15 (-) |
| Overstory tree cover (%) | 1.0 (-) | 1-1 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (2,2), NE (2,1), NW (2,0), S (1,1), SE (7,0), SW (1,0), Variable (2,0)

Macrotopography: middle to upper 1/3 of slope (2,2), ridge top (3,0), middle 1/3 of slope (4,0), bottom (1,0), upper 1/3 of slope (7,2)

Microtopography: Convex (6,2), flat (1,1), Concave (3,0), undulating (7,1)

Parent material: granitic (8,0), igneous (1,0), metamorphic (1,1), mix of two or more rock types (1,0), sandstone (3,3), sedimentary (3,0)

Soil texture: clay loam (5,1), silt (1,0), clay (1,1), sandy loam (6,1), loam (3,0)

| | Average | Range |
|------------------------------|----------------|----------------------|
| Elevation (ft) | 1381 (1596) | 872-2315 (1316-1880) |
| Slope (degrees) | 23 (28.3) | 5-45 (16-45) |
| Large rock cover (%) | 3.3 (-) | 0-15 (-) |
| Small rock cover (%) | 9.6 (4.0) | 0-35 (0-12) |
| Bare ground cover (%) | 31.6 (41.7) | 0-58 (28-55) |
| Litter cover (%) | 33.0 (51.3) | 0-69 (30-69) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-------------------------------|------------|------------|------------|------------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 25.7 (21) | 11 (11) | 55 (26) |
| <i>Salvia apiana</i> | 100 (100) | 15.2 (7) | 5 (5) | 27 (8) |
| <i>Malosma laurina</i> | 76 (25) | 3.7 (14) | 0.2 (14) | 14 (14) |
| <i>Eriogonum fasciculatum</i> | 71 (75) | 1.7 (1.4) | 0.2 (0.2) | 4 (3) |
| <i>Hesperoyucca whipplei</i> | 35 (25) | 1.1 (0.2) | 0.2 (0.2) | 4 (0.2) |
| <i>Salvia mellifera</i> | 35 (25) | 2.4 (2) | 0.2 (2) | 5 (2) |
| <i>Yucca whipplei</i> | 35 (-) | 1.1 (-) | 0.2 (-) | 4 (-) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 47 (50) | 8 (1) | 0.2 (1) | 30 (1) |
| <i>*Centaurea melitensis</i> | 41 (25) | 5.4 (2) | 1 (2) | 22 (2) |
| <i>*Hirschfeldia incana</i> | 41 (25) | 8 (0.2) | 0.2 (0.2) | 20 (0.2) |
| <i>Galium angustifolium</i> | 29 (100) | 2 (2.3) | 0.2 (0.2) | 5 (5) |

***Artemisia californica*-*Eriogonum fasciculatum* Alliance**

***Artemisia californica*-*Eriogonum fasciculatum*-*Malosma laurina* Association**

Samples Used To Describe Association (n= 50)

Rapid Assessment(s), WR and WSD stands: SDG0038, SDRP0261, 1617MB01, 1548MB01, 1700PG02, SDG0115, SDG0046, SRPL.019/675, SDG0066, SDRP0320, 1035MB01, SDRP0467, CLNF.001, SDRP0190, SAAN.012, SDRP0319, 1353MB01, 1437KH01, SDG0010, SDG0012, 1549KH01, 0519TK01, SDG0065, SDRP0090, 0505RB03, SDG0016, 0521RB03, 0520RB03, SDRP0125, SDRP0132, 0521TK01, CLNF.068, SDG0019, SDG0070, SDRP0130, 0603DH01, SDG0037, SDG0007, SDG0022, SDG0026, SDG0029, SDG0036, 0610AK01
Rapid Assessment(s), OC stands (n=7): 1056EK06, 1062EK02, 1076EK01, 1108EK01, 1407EK01, 1414EK01, 1516EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 50.2 (49.7) | 15-75 (38-70) | - |
| Herb cover (%) | 18.5 (15.7) | 0-55 (0.2-55) | 0-1 (0-1) |
| Shrub cover (%) | 39.2 (33.1) | 10-59 (10-48) | 0-5 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 0.2 (-) | 0.2-0.2 (-) | 5-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SE (15,0), W (2,1), SW (11,1), S (3,3), NW (6,1), NE (5,0), Flat (3,0), E (2,1), (0,0), Variable (2,0)
Macrotopography: middle 1/3 of slope (10,1), upper 1/3 of slope (6,3), middle to upper 1/3 of slope (6,1), lower to upper 1/3 of slope (5,0), lower to middle 1/3 of slope (7,1), lower 1/3 of slope (12,1), bottom (3,0), ridge top (1,0)
Microtopography: concave (5,0), convex (10,1), flat (6,2), undulating (28,4)
Parent material: gabbro (1,0), granitic (15,0), igneous (2,0), metamorphic (3,0), sandstone (7,7), sedimentary (2,0)
Soil texture: clay loam (12,1), loam (2,0), sandy loam (21,5), silty loam (1,1)

| | Average | Range |
|------------------------------|-------------|--------------------|
| Elevation (ft) | 802 (478) | 115-3062 (115-751) |
| Slope (degrees) | 14 (8.0) | 0-40 (3-32) |
| Large rock cover (%) | 1.1 (0.7) | 0-23 (0-5) |
| Small rock cover (%) | 4.9 (1.9) | 0-30 (0-8) |
| Bare ground cover (%) | 25.5 (44.1) | 0-85 (27-58) |
| Litter cover (%) | 25.2 (50.9) | 0-78 (35-70) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-------------------------------|-----------|-------------|-----------|---------|
| Shrub | <i>Artemisia californica</i> | 100 (100) | 15.9 (12.7) | 0.2 (3) | 38 (26) |
| | <i>Eriogonum fasciculatum</i> | 94 (100) | 10.4 (14.9) | 0.2 (1) | 33 (33) |
| | <i>Malosma laurina</i> | 78 (71) | 6.4 (0.7) | 0.2 (0.2) | 35 (2) |
| | <i>Salvia apiana</i> | 40 (14) | 1.2 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Bahiopsis laciniata</i> | 32 (-) | 1.9 (-) | 0.2 (-) | 7 (-) |
| | <i>Acmispon glaber</i> | 28 (71) | 0.7 (1.2) | 0.2 (0.2) | 5 (5) |
| | | | | | |
| Herb | <i>*Bromus madritensis</i> | 56 (43) | 7.3 (4.1) | 0.2 (0.2) | 50 (10) |
| | <i>*Centaurea melitensis</i> | 54 (57) | 3.5 (1.4) | 0.2 (0.2) | 25 (5) |
| | <i>*Hirschfeldia incana</i> | 30 (14) | 2.8 (1) | 0.2 (1) | 25 (1) |
| | <i>*Erodium</i> | 28 (-) | 9.5 (-) | 0.2 (-) | 40 (-) |

***Artemisia californica*-*Salvia mellifera* Alliance**
***Artemisia californica*-*Salvia mellifera* Association**

Samples Used To Describe Association (n= 52)

Rapid Assessment(s), WR and WSD stands: SDRP0075, 0625JJ01, 1769JD01, 0709RB03, 0709DH02, 0708JS02, 0708DH04, 0707DH04, 1710KH05, 0706AK01, 1769KH01, SDRP0271, 1710JD05, SDRP0062, 1470JD01, 1202JD01, 1737JD02, 1710KH03, SDRP0278, 0625RB03, 1336FS01, 1710JD11, SDRP0272, 0624AK01, 0623RB05, 1737KH02, SDG0113, 1513KH01, SDRP0142, 1710KH01, SDRP0277, SDRP0353, SDRP0354, SDRP0275, 1737JD06, SDRP0276, SDRP0402, 1698FS01, SDRP0403, 0623RB02

Rapid Assessment(s), OC stands (n=12): 1015EK05, 1047EK01, 1047EK04, 1056EK05, 1056EK08, 1088EK05, 1173EK01, 1413EK03, 1465MA01, 1520MA02, 1524EK01, 3.8jd01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-----------------|-------------|
| Total vegetation cover (%) | 51.9 (52.8) | 15-92 (39-85) | - |
| Herb cover (%) | 10.0 (8.6) | 0.2-40 (0.2-28) | 0-2 (0-0.5) |
| Shrub cover (%) | 46.6 (46.6) | 26-77 (26-77) | 0-5 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 2.1 (0.2) | 0.2-4 (0.2-0.2) | 2-10 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), W (1,0), Variable (1,1), SW (9,2), SE (11,4), S (1,1), NE (12,0), E (1,1), NW (15,2)

Macrotopography: entire slope (1,0), middle 1/3 of slope to ridgetop (2,0), upper 1/3 of slope (3,1), ridge top (2,1), upper 1/3 of slope to ridgetop (1,0), middle to upper 1/3 of slope (10,5), middle 1/3 of slope (8,2), lower to upper 1/3 of slope (6,0), lower to middle 1/3 of slope (8,2), lower 1/3 of slope (8,0), bottom (1,0), lower 1/3 of slope to ridgetop (1,1)

Microtopography: concave (5,1), convex (12,3), flat (10,4), undulating (25,4)

Parent material: conglomerate (1,0), diorite (1,0), granitic (3,0), metamorphic (2,0), mixed alluvium (4,0), mixed sedimentary (1,0), sandstone (18,7), sandy alluvium (2,0), sedimentary (3,0)

Soil texture: clay (1,0), clay loam (14,3), loam (3,2), sand (3,1), sandy loam (23,4), silt (1,0)

| | Average | Range |
|------------------------------|-------------|-------------------|
| Elevation (ft) | 376 (721) | 0-1453 (148-1453) |
| Slope (degrees) | 19 (19.5) | 2-38 (2-38) |
| Large rock cover (%) | 0.5 (-) | 0-17 (-) |
| Small rock cover (%) | 10.6 (11.1) | 0-80 (0-50) |
| Bare ground cover (%) | 32.2 (36.5) | 0-83 (5-58) |
| Litter cover (%) | 33.8 (50.3) | 0-90 (10-90) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-------------------------------|-----------|-------------|-----------|---------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 14.9 (21.3) | 2 (9) | 40 (40) |
| <i>Salvia mellifera</i> | 100 (100) | 13.2 (14.4) | 2 (4) | 45 (45) |
| <i>Rhus integrifolia</i> | 73 (67) | 8.2 (5.6) | 0.2 (0.2) | 31 (30) |
| <i>Malosma laurina</i> | 58 (67) | 3.9 (3.9) | 0.2 (0.2) | 12 (12) |
| <i>Eriogonum fasciculatum</i> | 56 (58) | 2.7 (2) | 0.2 (0.2) | 9 (7) |
| <i>Mimulus aurantiacus</i> | 52 (58) | 3.2 (1.3) | 0.2 (0.2) | 20 (7) |
| <i>Opuntia littoralis</i> | 44 (58) | 1.2 (1.4) | 0.2 (0.2) | 6 (6) |
| <i>Acmispon glaber</i> | 35 (33) | 2.6 (9.6) | 0.2 (0.2) | 37 (37) |
| Herb | | | | |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 58

| Species Name | Con | Ave | Min | Max |
|-------------------------------|---------|-----------|-----------|--------|
| * <i>Bromus madritensis</i> | 46 (25) | 4.2 (1.8) | 0.2 (0.2) | 15 (5) |
| * <i>Centaurea melitensis</i> | 37 (17) | 2.1 (4) | 0.2 (1) | 8 (7) |

***Atriplex amnicola* Alliance**

***Atriplex amnicola* Provisional Semi-Natural Stands**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1517EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 60.0 (60.0) | 60-60 (60-60) | - |
| Herb cover (%) | 7.0 (7.0) | 7-7 (7-7) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 55.0 (55.0) | 55-55 (55-55) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1)

Macrotopography: lower to middle 1/3 of slope (1,1)

Microtopography: undulating (1,1)

Parent material: sandstone (1,1)

Soil texture:

| | Average | Range |
|-----------------------|-------------|---------------|
| Elevation (ft) | 10 (10) | 10-10 (10-10) |
| Slope (degrees) | 15 (15.0) | 15-15 (15-15) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 38.0 (38.0) | 38-38 (38-38) |
| Litter cover (%) | 60.0 (60.0) | 60-60 (60-60) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--------------------------------|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| * <i>Acacia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Artemisia californica</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| * <i>Atriplex amnicola</i> | 100 (100) | 50 (50) | 50 (50) | 50 (50) |
| <i>Atriplex lentiformis</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Baccharis pilularis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Encelia californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Isocoma menziesii</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Anagallis arvensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus diandrus</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| * <i>Bromus hordeaceus</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| * <i>Carduus pycnocephalus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Centaurea melitensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|-----------------------------|-----------|-----------|-----------|-----------|
| <i>*Erodium cicutarium</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Euphorbia peplus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Medicago polymorpha</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Atriplex lentiformis* Alliance**
***Atriplex lentiformis* Association**

Samples Used To Describe Association (n= 2)

Rapid Assessment(s), WR and WSD stands: 0310AK23

Rapid Assessment(s), OC stands (n=1): 1404EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|---------------|
| Total vegetation cover (%) | 45.0 (50.0) | 40-50 (50-50) | - |
| Herb cover (%) | 6.1 (12.0) | 0.2-12 (12-12) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 38.0 (38.0) | 38-38 (38-38) | 1-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: S (1,1), SE (1,0)

Macrotopography: lower 1/3 of slope (1,0), lower to middle 1/3 of slope (1,1)

Microtopography: flat (2,1)

Parent material: mixed alluvium (1,0), sandstone (1,1)

Soil texture: clay loam (1,0), sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|---------------|
| Elevation (ft) | 33 (66) | 0-66 (66-66) |
| Slope (degrees) | 5 (3.0) | 3-6 (3-3) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 3.5 (5.0) | 2-5 (5-5) |
| Bare ground cover (%) | 30.0 (43.0) | 17-43 (43-43) |
| Litter cover (%) | 65.0 (50.0) | 50-80 (50-50) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-------------------------------|-----------|-----------|-----------|-----------|
| Shrub | <i>Atriplex lentiformis</i> | 100 (100) | 35 (30) | 30 (30) | 40 (30) |
| | <i>Artemisia californica</i> | 50 (100) | 5 (5) | 5 (5) | 5 (5) |
| | <i>Baccharis pilularis</i> | 50 (100) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Eriogonum fasciculatum</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Lycium californicum</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhus integrifolia</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | | | | | |
| Herb | <i>*Anagallis arvensis</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Brassica nigra</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Bromus madritensis</i> | 50 (100) | 10 (10) | 10 (10) | 10 (10) |
| | <i>*Carduus pycnocephalus</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Centaurea melitensis</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | | | | | |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 60

| | Species Name | Con | Ave | Min | Max |
|-----------|--------------------------------------|----------|-----------|-----------|-----------|
| | <i>Cressa truxillensis</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Festuca myuros</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Foeniculum vulgare</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Frankenia salina</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Galium parisiense</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Pseudognaphalium californicum</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Sonchus asper</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Cryptogam | | | | | |
| | * <i>Moss</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| Algae | | | | | |
| | <i>Algae</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Baccharis pilularis* Alliance**

***Baccharis pilularis*/Herbaceous Association**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: SDRP0129, 1177JD01, 1186JD01, 0625AK03

Rapid Assessment(s), OC stands (n=2): 1033JD01, 1089EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 41.8 (51.5) | 30-55 (48-55) | - |
| Herb cover (%) | 20.6 (6.0) | 6-30 (6-6) | 0-5 (0-0.5) |
| Shrub cover (%) | 48.5 (48.5) | 42-55 (42-55) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 0.2 (-) | 0.2-0.2 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (2,0), NE (2,1), NW (1,0), SE (1,1)

Macrotopography: lower 1/3 of slope (1,0), lower to middle 1/3 of slope (1,1), bottom (3,0), middle 1/3 of slope (1,1)

Microtopography: concave (2,1), convex (1,1), flat (3,0)

Parent material: clayey alluvium (1,0), granitic (1,0), sandstone (1,1)

Soil texture: clay (1,0), clay loam (1,1), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|----------------|
| Elevation (ft) | 316 (194) | 0-597 (20-367) |
| Slope (degrees) | 7 (15.5) | 0-26 (5-26) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 2.8 (-) | 0-9 (-) |
| Bare ground cover (%) | 37.7 (32.5) | 0-65 (32-33) |
| Litter cover (%) | 41.0 (65.0) | 0-65 (65-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | * <i>Schinus molle</i> | 33 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Baccharis pilularis</i> | 100 (100) | 26 (39) | 8 (26) | 52 (52) |
| | <i>Artemisia californica</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 61

| Species Name | Con | Ave | Min | Max |
|--------------------------------------|----------|-----------|-----------|-----------|
| <i>Isocoma menziesii</i> | 50 (50) | 5.7 (15) | 1 (15) | 15 (15) |
| <i>Eriogonum fasciculatum</i> | 33 (-) | 2.1 (-) | 0.2 (-) | 4 (-) |
| <i>Mimulus aurantiacus</i> | 33 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Bromus hordeaceus</i> | 67 (50) | 4.1 (5) | 0.2 (5) | 10 (5) |
| * <i>Foeniculum vulgare</i> | 67 (50) | 6.8 (0.2) | 0.2 (0.2) | 20 (0.2) |
| * <i>Hirschfeldia incana</i> | 50 (-) | 0.8 (-) | 0.2 (-) | 2 (-) |
| <i>Pseudognaphalium californicum</i> | 50 (100) | 1.1 (0.2) | 0.2 (0.2) | 3 (0.2) |
| <i>Ambrosia psilostachya</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| * <i>Brassica nigra</i> | 33 (50) | 4.1 (0.2) | 0.2 (0.2) | 8 (0.2) |
| * <i>Bromus madritensis</i> | 33 (-) | 7.5 (-) | 3 (-) | 12 (-) |
| * <i>Centaurea melitensis</i> | 33 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| * <i>Helminthotheca echioides</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Baccharis salicifolia* Alliance**
***Baccharis salicifolia* Association**

Samples Used To Describe Association (n= 25)

Rapid Assessment(s), WR and WSD stands: 0520DH06, 0709AK01, SDRP0078, 1361JD02, 1626KH01, SDRP0200, 0504MJ23, SDRP0081, 1784FS01, SDRP0150, SDRP0316, 1604LS02, SDG0106, 1019DH01, 1545EB01, 0506RB04, 1766KH02, 1163FS01, 0506AK05, SDRP0067

Rapid Assessment(s), OC stands (n=5): 1014EK01, 1243MA01, 1426MA01, 1499EK01, 1510EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-------------------|-------------|
| Total vegetation cover (%) | 49.7 (42.6) | 15-90 (23-77) | - |
| Herb cover (%) | 19.0 (7.6) | 0.2-72 (0.2-14) | 0-2 (0-1) |
| Shrub cover (%) | 35.4 (35.4) | 18-74 (18-74) | 0-10 (1-5) |
| Tall shrub / low tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 1-2 (1-2) |
| Overstory tree cover (%) | 2.0 (1.4) | 0.2-8 (0.2-2) | 2-15 (2-10) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (3,0), Variable (2,0), SW (6,1), S (3,3), NE (2,0), Flat (8,0), SE (1,1)

Macrotopography: bottom (19,4), bottom to lower 1/3 of slope (1,0), bottom to mid 1/3 of slope (1,0), lower 1/3 of slope (2,0), middle 1/3 of slope (1,0)

Microtopography: concave (8,3), flat (13,2), undulating (4,0)

Parent material: clayey alluvium (1,0), gravelly alluvium (1,0), mixed alluvium (6,1), sandstone (3,3), sandy alluvium (6,0), sedimentary (1,0), silty alluvium (3,0)

Soil texture: sandy loam (5,0), clay loam (1,0), silty loam (2,0), loam (1,0), clay (2,0), sand (9,4)

| | Average | Range |
|------------------------------|-------------|-------------------|
| Elevation (ft) | 546 (793) | 0-1263 (381-1079) |
| Slope (degrees) | 2 (1.8) | 0-15 (1-3) |
| Large rock cover (%) | 5.0 (-) | 0-45 (-) |
| Small rock cover (%) | 15.6 (17.2) | 0-78 (0-45) |
| Bare ground cover (%) | 21.1 (48.4) | 0-75 (4-75) |
| Litter cover (%) | 33.9 (32.2) | 0-95 (5-88) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 62

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------------|-----------|-------------|-----------|----------|
| Tree | | | | | |
| | <i>Salix lasiolepis</i> | 32 (20) | 1.5 (2) | 0.2 (2) | 3 (2) |
| Shrub | | | | | |
| | <i>Baccharis salicifolia</i> | 100 (100) | 26.9 (28.4) | 8 (15) | 72 (55) |
| | <i>Artemisia californica</i> | 32 (80) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>Sambucus nigra subsp. caerulea</i> | 28 (40) | 2.3 (3.5) | 0.2 (2) | 6 (5) |
| | * <i>Tamarix</i> | 28 (20) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| Herb | | | | | |
| | <i>Ambrosia psilostachya</i> | 40 (60) | 1 (0.5) | 0.2 (0.2) | 4 (1) |
| | * <i>Hirschfeldia incana</i> | 40 (20) | 1.4 (4) | 0.2 (4) | 4 (4) |
| | * <i>Centaurea melitensis</i> | 32 (40) | 0.8 (0.6) | 0.2 (0.2) | 2 (1) |
| | * <i>Avena barbata</i> | 28 (40) | 5.5 (0.2) | 0.2 (0.2) | 31 (0.2) |
| | * <i>Bromus diandrus</i> | 28 (60) | 3.3 (2) | 0.2 (1) | 8 (3) |

***Ceanothus crassifolius* Alliance**
***Ceanothus crassifolius* Association**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: SDRP0479, SDRP0465, SDRP0438, 0605DH02, 0506AK01
Rapid Assessment(s), OC stands (n=1): 1235EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|---------------|
| Total vegetation cover (%) | 60.7 (38.0) | 34-95 (38-38) | - |
| Herb cover (%) | 17.8 (3.0) | 3-65 (3-3) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 37.0 (37.0) | 37-37 (37-37) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SE (1,0), W (1,1), NW (2,0), NE (2,0)
Macrotopography: lower to upper 1/3 of slope (4,0), middle 1/3 of slope (2,1)
Microtopography: flat (1,0), undulating (5,1)
Parent material: granitic (5,0), sandstone (1,1)
Soil texture: loam (2,0), sandy loam (3,0), clay loam (1,1)

| | Average | Range |
|------------------------------|-------------|----------------------|
| Elevation (ft) | 1134 (1795) | 689-1795 (1795-1795) |
| Slope (degrees) | 24 (40.0) | 7-40 (40-40) |
| Large rock cover (%) | 1.7 (-) | 0-8 (-) |
| Small rock cover (%) | 10.2 (43.0) | 0-43 (43-43) |
| Bare ground cover (%) | 20.0 (21.0) | 5-39 (21-21) |
| Litter cover (%) | 58.8 (33.0) | 33-80 (33-33) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 63

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---|-----------|-----------|-----------|-----------|
| Shrub | <i>Ceanothus crassifolius</i> | 100 (100) | 34.5 (6) | 6 (6) | 60 (6) |
| | <i>Malosma laurina</i> | 83 (100) | 2.8 (1) | 1 (1) | 5 (1) |
| | <i>Xylococcus bicolor</i> | 83 (-) | 3.3 (-) | 0.2 (-) | 8 (-) |
| | <i>Salvia mellifera</i> | 67 (100) | 4.6 (1) | 0.2 (1) | 12 (1) |
| | <i>Acmispon glaber</i> | 50 (100) | 3.8 (11) | 0.2 (11) | 11 (11) |
| | <i>Adenostoma fasciculatum</i> | 50 (-) | 5.3 (-) | 1 (-) | 10 (-) |
| | <i>Artemisia californica</i> | 50 (100) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Hazardia squarrosa</i> | 50 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Rhus ovata</i> | 50 (100) | 1.4 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>Ceanothus tomentosus</i> | 33 (-) | 4.5 (-) | 2 (-) | 7 (-) |
| | <i>Cneoridium dumosum</i> | 33 (-) | 2.6 (-) | 0.2 (-) | 5 (-) |
| | <i>Eriogonum fasciculatum</i> | 33 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| | <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Mimulus aurantiacus</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Quercus berberidifolia</i> | 33 (-) | 5.5 (-) | 3 (-) | 8 (-) |
| Herb | <i>Marah macrocarpus</i> var. <i>macrocarpus</i> | 83 (-) | 3.4 (-) | 0.2 (-) | 6 (-) |
| | * <i>Bromus madritensis</i> | 50 (-) | 8 (-) | 1 (-) | 15 (-) |
| | * <i>Bromus diandrus</i> | 33 (-) | 6.5 (-) | 3 (-) | 10 (-) |
| | <i>Cryptantha</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Galium angustifolium</i> | 33 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Helianthus gracilentus</i> | 33 (100) | 0.6 (1) | 0.2 (1) | 1 (1) |
| | * <i>Hirschfeldia incana</i> | 33 (100) | 1.5 (1) | 1 (1) | 2 (1) |
| | <i>Pseudognaphalium californicum</i> | 33 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Ceanothus megacarpus* Alliance**

***Ceanothus megacarpus* Association**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1521EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-------------------|---------------|
| Total vegetation cover (%) | 65.0 (65.0) | 65-65 (65-65) | - |
| Herb cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 65.0 (65.0) | 65-65 (65-65) | 2-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 64

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,1)

Macrotopography: middle 1/3 of slope (1,1)

Microtopography: convex (1,1)

Parent material: sandstone (1,1)

Soil texture: sandy loam (1,1)

| | Average | Range |
|------------------------------|----------------|-------------------|
| Elevation (ft) | 627 (627) | 627-627 (627-627) |
| Slope (degrees) | 40 (40.0) | 40-40 (40-40) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 5.0 (5.0) | 5-5 (5-5) |
| Litter cover (%) | 92.0 (92.0) | 92-92 (92-92) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--------------------------------|------------|------------|------------|------------|
| Shrub | | | | |
| <i>Adenostoma fasciculatum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Ceanothus megacarpus</i> | 100 (100) | 63 (63) | 63 (63) | 63 (63) |
| <i>Heteromeles arbutifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Keckiella cordifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Malosma laurina</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Marah</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Quercus berberidifolia</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Rhus integrifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Ceanothus tomentosus* Alliance**

***Ceanothus tomentosus* Association**

Samples Used To Describe Association (n= 28)

Rapid Assessment(s), WR and WSD stands: 1155JD01, 0605RB02, 0519DH02, 1361JD01, 0311DH22, 0505MG01, 1329EB02, 0602DH03, 1249TK01, 0519AK02, 0624AK04, 1362MB01, 0422AK01, 1607JD02, 0506AK02, 1604EB03, SDRP0425, 1486MM02, 1827MM01, 0708TK01, 1146JD01, 0422DH01, 1361JD03, 0504RB23, 1381JD01, 0609DH01

Rapid Assessment(s), OC stands (n=2): 1341EK02, 1343EK05

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|----------------|-------------------|
| Total vegetation cover (%) | 57.3 (47.5) | 25-85 (47-48) | - |
| Herb cover (%) | 12.6 (0.2) | 0-30 (0.2-0.2) | 0-1 (0-0.5) |
| Shrub cover (%) | 47.5 (47.5) | 47-48 (47-48) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-2 (-) |
| Overstory tree cover (%) | 5.0 (-) | 1-8 (-) | 2-5 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 65

Local Environmental Description (All stands, OC stands)

Aspect: NE (9,1), SW (3,0), NW (14,0), SE (2,1)

Macrotopography: middle to upper 1/3 of slope (6,0), middle 1/3 of slope to ridgetop (1,0), middle 1/3 of slope (6,1), lower to upper 1/3 of slope (6,0), lower to middle 1/3 of slope (1,0), lower 1/3 of slope (3,0), bottom to lower 1/3 of slope (2,0), bottom to upper 1/3 of slope (1,0)

Microtopography: flat (6,0), undulating (9,1), convex (10,0), concave (3,1)

Parent material: diorite (1,0), gabbro (3,0), granitic (8,0), metamorphic (5,0), sandstone (2,2), volcanic flow (1,0)

Soil texture: loam (1,0), silty loam (1,1), sandy loam (15,1), clay loam (9,0), muck (1,0)

| | Average | Range |
|------------------------------|----------------|----------------------|
| Elevation (ft) | 1416 (1920) | 174-2684 (1867-1972) |
| Slope (degrees) | 24 (27.0) | 2-42 (25-29) |
| Large rock cover (%) | 8.1 (0.5) | 0-35 (0-1) |
| Small rock cover (%) | 22.8 (-) | 0-68 (-) |
| Bare ground cover (%) | 25.4 (42.5) | 0-84 (36-49) |
| Litter cover (%) | 36.9 (54.0) | 0-70 (48-60) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--|------------|------------|------------|------------|
| Shrub | <i>Ceanothus tomentosus</i> | 100 (100) | 32.4 (36) | 7 (32) | 60 (40) |
| | <i>Malosma laurina</i> | 75 (100) | 3.7 (1.5) | 0.2 (1) | 15 (2) |
| | <i>Adenostoma fasciculatum</i> | 54 (100) | 3.5 (2) | 0.2 (1) | 10 (3) |
| | <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 43 (-) | 0.8 (-) | 0.2 (-) | 3 (-) |
| | <i>Xylococcus bicolor</i> | 43 (-) | 2.7 (-) | 0.2 (-) | 8 (-) |
| | <i>Hazardia squarrosa</i> | 36 (-) | 0.5 (-) | 0.2 (-) | 2 (-) |
| | <i>Acemispion glaber</i> | 32 (50) | 3.1 (2) | 0.2 (2) | 10 (2) |
| | <i>Malacothamnus fasciculatus</i> | 29 (100) | 1.6 (1) | 0.2 (1) | 4 (1) |
| | <i>Eriogonum fasciculatum</i> | 25 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| | <i>Heteromeles arbutifolia</i> | 25 (50) | 0.9 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>Mimulus aurantiacus</i> | 25 (-) | 1.1 (-) | 0.2 (-) | 5 (-) |
| | <i>Rhamnus crocea</i> | 25 (-) | 1 (-) | 0.2 (-) | 3 (-) |
| | <i>Rhus ovata</i> | 25 (-) | 3.3 (-) | 0.2 (-) | 15 (-) |
| Herb | <i>*Bromus madritensis</i> | 43 (-) | 6.4 (-) | 0.2 (-) | 20 (-) |
| | <i>*Gastridium phleoides</i> | 32 (-) | 0.4 (-) | 0.2 (-) | 2 (-) |
| | <i>Calystegia macrostegia</i> | 29 (-) | 1 (-) | 0.2 (-) | 3 (-) |
| | <i>*Centaurea melitensis</i> | 25 (-) | 3.2 (-) | 0.2 (-) | 10 (-) |
| | <i>*Festuca myuros</i> | 25 (-) | 7.9 (-) | 2 (-) | 18 (-) |

***Cercocarpus betuloides* Alliance**
***Cercocarpus betuloides* Association**

Samples Used To Describe Association (n= 1)
Rapid Assessment(s), WR and WSD stands: None
Rapid Assessment(s), OC stands (n=1): 1231JD01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 62.0 (62.0) | 62-62 (62-62) | - |
| Herb cover (%) | - (-) | - (-) | - (-) |
| Shrub cover (%) | - (-) | - (-) | - (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (1,1)
Macrotopography: middle to upper 1/3 of slope (1,1)
Microtopography: undulating (1,1)
Parent material:
Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 5289 (5289) | 5289-5289 (5289-5289) |
| Slope (degrees) | 26 (26.0) | 26-26 (26-26) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 10.0 (10.0) | 10-10 (10-10) |
| Bare ground cover (%) | 33.0 (33.0) | 33-33 (33-33) |
| Litter cover (%) | 55.0 (55.0) | 55-55 (55-55) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>Ceanothus leucodermis</i> | 100 (100) | 12 (12) | 12 (12) | 12 (12) |
| <i>Cercocarpus betuloides</i> var. <i>betuloides</i> | 100 (100) | 45 (45) | 45 (45) | 45 (45) |
| <i>Eriophyllum confertiflorum</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Garrya flavescens</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Keckiella antirrhinoides</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Stephanomeria cichoriacea</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| Herb | | | | |
| <i>Cirsium occidentale</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Clarkia purpurea</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Encelia californica* Alliance**
***Encelia californica* (Alliance Only)**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None
Rapid Assessment(s), OC stands (n=1): 1404EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 54.0 (54.0) | 54-54 (54-54) | - |
| Herb cover (%) | 4.0 (4.0) | 4-4 (4-4) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 50.0 (50.0) | 50-50 (50-50) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: S (1,1)
Macrotopography: upper 1/3 of slope (1,1)
Microtopography: concave (1,1)
Parent material: sandstone (1,1)
Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|---------------|
| Elevation (ft) | 39 (39) | 39-39 (39-39) |
| Slope (degrees) | 30 (30.0) | 30-30 (30-30) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 53.0 (53.0) | 53-53 (53-53) |
| Litter cover (%) | 45.0 (45.0) | 45-45 (45-45) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>*Atriplex amnicola</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Atriplex semibaccata</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Baccharis pilularis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Cylindropuntia prolifera</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Encelia californica</i> | 100 (100) | 45 (45) | 45 (45) | 45 (45) |
| <i>Eriogonum fasciculatum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lycium californicum</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| <i>Mirabilis laevis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Peritoma arborea</i> var. <i>arborea</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Chorizanthe</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Distichlis spicata</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>*Mesembryanthemum crystallinum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Salsola tragus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Encelia californica Alliance

Encelia californica-Artemisia californica Association

Samples Used To Describe Association (n= 18)

Rapid Assessment(s), WR and WSD stands: 0310RB24, 0708MG01, HASP.016, CORN.009/392, 1737KH01, LAMA.019, SDG0087, 0309RB24, 1520JD01, 0309RB25, LAMA.006, 0707MG01, SDG0083

Rapid Assessment(s), OC stands (n=5): 1075EK01, 1168EK01, 1407EK02, 1471EK02, 1477EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|-------------|
| Total vegetation cover (%) | 48.2 (54.2) | 25-75 (50-61) | - |
| Herb cover (%) | 11.0 (0.4) | 0.2-45 (0.2-1) | 0-2 (0-0.5) |
| Shrub cover (%) | 46.4 (54.0) | 25-60 (50-60) | 0-2 (0-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-2 (-) |
| Overstory tree cover (%) | 2.0 (-) | 2-2 (-) | 10-15 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (9,3), NE (1,0), NW (2,1), S (1,1), SE (5,0)

Macrotopography: entire slope (1,0), lower 1/3 of slope (3,0), lower to middle 1/3 of slope (1,1), lower to upper 1/3 of slope (1,0), middle 1/3 of slope (6,2), ridge top (1,0), upper 1/3 of slope (4,1)

Microtopography: convex (8,0), flat (5,3), concave (3,1), Undulating (2,1)

Parent material: granitic (5,0), mixed sedimentary (1,0), sandstone (7,5), sedimentary (1,0), volcanic flow (1,0)

Soil texture: sand (5,2), sandy loam (9,2), clay loam (3,0)

| | Average | Range |
|-----------------------|-------------|-----------------|
| Elevation (ft) | 439 (70) | 0-2095 (20-121) |
| Slope (degrees) | 20 (16.4) | 2-50 (2-50) |
| Large rock cover (%) | 2.9 (-) | 0-15 (-) |
| Small rock cover (%) | 10.3 (-) | 0-60 (-) |
| Bare ground cover (%) | 44.6 (49.3) | 4-74 (33-58) |
| Litter cover (%) | 39.2 (48.8) | 3-94 (40-65) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---------------------------------|-----------|-------------|-----------|----------|
| Shrub | | | | |
| <i>Encelia californica</i> | 100 (100) | 24.1 (29.4) | 2 (8) | 48 (47) |
| <i>Artemisia californica</i> | 94 (100) | 9.3 (14.8) | 0.2 (1) | 37 (37) |
| <i>Eriogonum fasciculatum</i> | 56 (40) | 2.3 (1) | 0.2 (1) | 8 (1) |
| <i>Rhus integrifolia</i> | 50 (80) | 1.4 (0.8) | 0.2 (0.2) | 2 (1) |
| <i>Cylindropuntia prolifera</i> | 28 (20) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 50 (20) | 4.3 (1) | 0.2 (1) | 18 (1) |
| * <i>Brassica nigra</i> | 39 (60) | 3.8 (0.2) | 0.2 (0.2) | 17 (0.2) |

***Ericameria palmeri* Alliance**

***Ericameria palmeri/Stipa pulchra* Provisional Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=4): 1110EK04, 1298MA02, 1449EK02, 1456MA01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|---------------|
| Total vegetation cover (%) | 58.0 (58.0) | 40-77 (40-77) | - |
| Herb cover (%) | 25.8 (25.8) | 14-35 (14-35) | 0-1 (0-1) |
| Shrub cover (%) | 33.5 (33.5) | 26-37 (26-37) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (1,1), NE (2,2), Flat (1,1)

Macrotopography: lower 1/3 of slope (4,4)

Microtopography: flat (2,2), undulating (2,2)

Parent material: sandstone (2,2)

Soil texture: clay loam (2,2)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 835 (835) | 600-1020 (600-1020) |
| Slope (degrees) | 8 (8.3) | 1-15 (1-15) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 2.0 (2.0) | 0-5 (0-5) |
| Bare ground cover (%) | 18.3 (18.3) | 9-31 (9-31) |
| Litter cover (%) | 77.0 (77.0) | 66-85 (66-85) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|-----------|-------------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Ericameria palmeri</i> var. <i>pachylepis</i> | 50 (100) | 35 (32.8) | 35 (26) | 35 (35) |
| | <i>Ericameria palmeri</i> var. <i>pachylepis</i> | 50 (100) | 30.5 (32.8) | 26 (26) | 35 (35) |
| | <i>Isocoma menziesii</i> | 50 (75) | 0.6 (0.5) | 0.2 (0.2) | 1 (1) |
| | <i>Baccharis pilularis</i> | 25 (25) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Isocoma menziesii</i> | 25 (75) | 0.2 (0.5) | 0.2 (0.2) | 0.2 (1) |
| | <i>Mimulus aurantiacus</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhamnus ilicifolia</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Sambucus nigra</i> subsp. <i>caerulea</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | | |
| | * <i>Bromus diandrus</i> | 100 (100) | 17 (17) | 4 (4) | 32 (32) |
| | <i>Stipa pulchra</i> | 100 (100) | 8.3 (8.3) | 3 (3) | 14 (14) |
| | * <i>Brassica nigra</i> | 75 (75) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| | * <i>Avena barbata</i> | 50 (50) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | * <i>Cynara cardunculus</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Elymus condensatus</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Amsinckia intermedia</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|--------------------------------------|---------|-----------|-----------|-----------|
| * <i>Avena fatua</i> | 25 (25) | 10 (10) | 10 (10) | 10 (10) |
| * <i>Bromus hordeaceus</i> | 25 (25) | 5 (5) | 5 (5) | 5 (5) |
| * <i>Centaurea melitensis</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Dichelostemma capitatum</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Erodium botrys</i> | 25 (25) | 1 (1) | 1 (1) | 1 (1) |
| * <i>Erodium cicutarium</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Festuca perennis</i> | 25 (25) | 3 (3) | 3 (3) | 3 (3) |
| * <i>Hirschfeldia incana</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Lactuca serriola</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Marrubium vulgare</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Medicago polymorpha</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Phacelia</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium californicum</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Rumex crispus</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Sisyrinchium bellum</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Torilis arvensis</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Vicia villosa</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Eriodictyon crassifolium* Alliance**

***Eriodictyon crassifolium* (Alliance Only)**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: LAFU.024, CLNF.039, BLACK.006/339, BLACK.008/976

Rapid Assessment(s), OC stands (n=2): 1161EK01, 1363EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 37.2 (33.0) | 27-50 (27-39) | - |
| Herb cover (%) | 14.4 (6.0) | 1-30 (6-6) | 0-0.5 (-) |
| Shrub cover (%) | 29.8 (30.0) | 20-39 (21-39) | 0-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 1.0 (-) | 1-1 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: E (1,1), S (1,1), SW (1,0), Variable (3,0)

Macrotopography: middle 1/3 of slope (4,0), middle to upper 1/3 of slope (1,1), upper 1/3 of slope (1,1)

Microtopography: Convex (2,0), Flat (1,0), undulating (3,2)

Parent material: granitic (3,1), sandstone (1,1), sedimentary (2,0)

Soil texture: clay loam (1,0), loam (1,0), sand (1,1), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3435 (2400) | 2159-5629 (2159-2641) |
| Slope (degrees) | 14 (28.0) | 2-30 (26-30) |
| Large rock cover (%) | 2.0 (1.5) | 0-5 (0-3) |
| Small rock cover (%) | 42.5 (20.5) | 1-70 (1-40) |
| Bare ground cover (%) | 37.2 (59.0) | 7-80 (38-80) |
| Litter cover (%) | 16.2 (17.5) | 10-25 (15-20) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 71

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|----------------------------------|-----------|------------|-----------|----------|
| Tree | | | | | |
| | <i>Quercus wislizeni</i> | 33 (-) | 1.5 (-) | 1 (-) | 2 (-) |
| Shrub | | | | | |
| | <i>Eriodictyon crassifolium</i> | 100 (100) | 14 (11) | 9 (9) | 20 (13) |
| | <i>Adenostoma fasciculatum</i> | 83 (100) | 5.6 (11.5) | 0.2 (7) | 16 (16) |
| | <i>Eriogonum fasciculatum</i> | 67 (-) | 4.6 (-) | 0.2 (-) | 8 (-) |
| | <i>Ceanothus crassifolius</i> | 50 (100) | 0.8 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>Rhus ovata</i> | 50 (50) | 1.7 (1) | 0.2 (1) | 4 (1) |
| | <i>Salvia mellifera</i> | 50 (100) | 2.7 (0.6) | 0.2 (0.2) | 7 (1) |
| | <i>Helianthemum scoparium</i> | 33 (100) | 4.6 (4.6) | 0.2 (0.2) | 9 (9) |
| Herb | | | | | |
| | <i>*Bromus madritensis</i> | 50 (50) | 3.5 (0.2) | 0.2 (0.2) | 10 (0.2) |
| | <i>*Bromus tectorum</i> | 33 (-) | 6 (-) | 2 (-) | 10 (-) |
| | <i>Chaenactis artemisiifolia</i> | 33 (50) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>Cryptantha muricata</i> | 33 (-) | 2 (-) | 1 (-) | 3 (-) |
| | <i>Gilia</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Eriogonum fasciculatum* Alliance**

***Eriogonum fasciculatum*/*Salvia columbariae*-*Mirabilis laevis* Association**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: SDG0067, 0507TK02, SDRP0421, 0506DH21, SDRP0441

Rapid Assessment(s), OC stands (n=1): 1513EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|--------------|-------------|
| Total vegetation cover (%) | 29.0 (10.0) | 8-55 (10-10) | - |
| Herb cover (%) | 24.5 (5.0) | 5-50 (5-5) | 0-2 (0-0.5) |
| Shrub cover (%) | 5.0 (5.0) | 5-5 (5-5) | 0-2 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 2-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: S (1,1), SE (1,0), SW (3,0), Variable (1,0)

Macrotopography: lower to middle 1/3 of slope (1,0), middle 1/3 of slope (3,1), middle to upper 1/3 of slope (2,0)

Microtopography: concave (1,0), undulating (2,1), convex (1,0), flat (2,0)

Parent material: conglomerate (1,0), granitic (3,0), sandstone (2,1)

Soil texture: sand (1,1), sandy loam (4,0), silty loam (1,0)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 948 (974) | 715-1227 (974-974) |
| Slope (degrees) | 26 (45.0) | 12-45 (45-45) |
| Large rock cover (%) | 0.0 (-) | 0-0.2 (-) |
| Small rock cover (%) | 10.9 (-) | 0-50 (-) |
| Bare ground cover (%) | 53.8 (94.0) | 8-94 (94-94) |
| Litter cover (%) | 25.7 (5.0) | 5-80 (5-5) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---|-----------|-----------|-----------|-----------|
| Shrub | <i>Eriogonum fasciculatum</i> | 100 (100) | 3.7 (2) | 1 (2) | 6 (2) |
| | <i>Artemisia californica</i> | 83 (100) | 0.9 (1) | 0.2 (1) | 2 (1) |
| | <i>Mirabilis laevis</i> | 83 (100) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Acmispon glaber</i> | 50 (100) | 1.1 (2) | 0.2 (2) | 2 (2) |
| | <i>Hesperoyucca whipplei</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Malosma laurina</i> | 50 (-) | 3.7 (-) | 2 (-) | 5 (-) |
| | <i>Cneoridium dumosum</i> | 33 (-) | 2 (-) | 1 (-) | 3 (-) |
| | <i>Salvia mellifera</i> | 33 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| Herb | <i>*Bromus madritensis</i> | 83 (100) | 8.8 (0.2) | 0.2 (0.2) | 23 (0.2) |
| | <i>*Centaurea melitensis</i> | 83 (-) | 4.2 (-) | 0.2 (-) | 8 (-) |
| | <i>*Hirschfeldia incana</i> | 67 (-) | 3.6 (-) | 0.2 (-) | 8 (-) |
| | <i>Salvia columbariae</i> | 67 (100) | 3.3 (0.2) | 0.2 (0.2) | 9 (0.2) |
| | <i>Selaginella bigelovii</i> | 67 (-) | 3.6 (-) | 0.2 (-) | 7 (-) |
| | <i>*Avena fatua</i> | 50 (-) | 10.3 (-) | 4 (-) | 20 (-) |
| | <i>*Erodium cicutarium</i> | 50 (-) | 1.4 (-) | 0.2 (-) | 2 (-) |
| | <i>Phacelia parryi</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Stephanomeria virgata</i> | 50 (100) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Calystegia macrostegia</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Chaenactis glabriuscula</i> | 33 (100) | 2.5 (1) | 1 (1) | 4 (1) |
| | <i>Chamaesyce polycarpa</i> | 33 (100) | 0.6 (1) | 0.2 (1) | 1 (1) |
| | <i>Cryptantha</i> | 33 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Cryptantha intermedia</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>*Hypochaeris glabra</i> | 33 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Lepidium nitidum</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Mimulus brevipes</i> | 33 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Muhlenbergia microsperma</i> | 33 (100) | 0.6 (1) | 0.2 (1) | 1 (1) |
| | <i>Phacelia cicutaria</i> var. <i>hispida</i> | 33 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| | <i>Pterostegia drymarioides</i> | 33 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |

Eriogonum fasciculatum Alliance

Eriogonum fasciculatum-*Salvia apiana* Association

Samples Used To Describe Association (n= 18)

Rapid Assessment(s), WR and WSD stands: SAGE.038, 1125PG01, CLNF.050, SDRP0195, SAGE.068, CLNF.042, SDRP0188, 1542MM01, SDRP0083, 1614MB01, CLNF.072, CORS.002/394/BPU 12856-240, WINC.008/226, WINC.009, MURR.004, CLNF.091, CLNF.017
Rapid Assessment(s), OC stands (n=1): 1346EK01

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 73

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 43.3 (32.0) | 27-75 (32-32) | - |
| Herb cover (%) | 14.7 (2.0) | 0-47 (2-2) | 0-1 (0-0.5) |
| Shrub cover (%) | 34.8 (30.0) | 14-58 (30-30) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (2,0), Variable (5,0), SE (5,0), NE (2,0), SW (4,1)
Macrotopography: lower to middle 1/3 of slope (3,0), lower to upper 1/3 of slope (2,0), middle 1/3 of slope (5,1), middle to upper 1/3 of slope (1,0), ridge top (1,0), upper 1/3 of slope (6,0)
Microtopography: Convex (7,0), Flat (2,1), undulating (9,0)
Parent material: granitic (7,0), igneous (1,0), metamorphic (2,0), sandstone (1,1), sedimentary (5,0)
Soil texture: clay loam (9,1), sandy loam (7,0), loam (1,0)

| | Average | Range |
|------------------------------|----------------|--------------------|
| Elevation (ft) | 1879 (961) | 423-3499 (961-961) |
| Slope (degrees) | 23 (8.0) | 8-44 (8-8) |
| Large rock cover (%) | 3.7 (-) | 0-23 (-) |
| Small rock cover (%) | 25.1 (-) | 0-75 (-) |
| Bare ground cover (%) | 23.8 (33.0) | 0-45 (33-33) |
| Litter cover (%) | 19.7 (65.0) | 0-65 (65-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------|------------|------------|------------|------------|
| Shrub | <i>Salvia apiana</i> | 100 (100) | 10.1 (8) | 1 (8) | 26 (8) |
| | <i>Eriogonum fasciculatum</i> | 89 (100) | 18.4 (15) | 4 (15) | 37 (15) |
| | <i>Artemisia californica</i> | 78 (100) | 4 (6) | 0.2 (6) | 17 (6) |
| | <i>Hesperoyucca whipplei</i> | 39 (-) | 1.1 (-) | 0.2 (-) | 3 (-) |
| | <i>Malosma laurina</i> | 39 (-) | 4.6 (-) | 0.2 (-) | 14 (-) |
| | <i>Yucca whipplei</i> | 39 (-) | 1.1 (-) | 0.2 (-) | 3 (-) |
| | <i>Acmispon glaber</i> | 33 (100) | 1.3 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Keckiella antirrhinoides</i> | 33 (-) | 2.6 (-) | 0.2 (-) | 5 (-) |
| | <i>Lotus scoparius</i> | 33 (-) | 1.3 (-) | 0.2 (-) | 5 (-) |
| | <i>Adenostoma fasciculatum</i> | 28 (-) | 1.1 (-) | 0.2 (-) | 4 (-) |
| | <i>Rhamnus crocea</i> | 28 (-) | 2.2 (-) | 1 (-) | 5 (-) |
| | | | | | |
| Herb | <i>*Hirschfeldia incana</i> | 33 (100) | 1.3 (0.2) | 0.2 (0.2) | 4 (0.2) |
| | <i>*Centaurea melitensis</i> | 28 (-) | 2.7 (-) | 0.2 (-) | 7 (-) |

***Isocoma menziesii* Alliance**
***Isocoma menziesii* Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: 1019DH02, 0312RB22, SDRP0242

Rapid Assessment(s), OC stands (n=1): 1057EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 44.8 (60.0) | 18-60 (60-60) | - |
| Herb cover (%) | 27.0 (20.0) | 10-45 (20-20) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 45.0 (45.0) | 45-45 (45-45) | 0-5 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (2,0), SW (2,1)

Macrotopography: bottom to lower 1/3 of slope (1,0), lower 1/3 of slope (3,1)

Microtopography: convex (1,0), flat (3,1)

Parent material: granitic (1,0), sandstone (1,1), sandy alluvium (1,0), silty alluvium (1,0)

Soil texture: clay loam (1,0), sandy loam (2,1), silt (1,0)

| | Average | Range |
|-----------------------|-------------|-----------------|
| Elevation (ft) | 119 (105) | 0-371 (105-105) |
| Slope (degrees) | 3 (2.0) | 0-10 (2-2) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 0.9 (3.0) | 0-3 (3-3) |
| Bare ground cover (%) | 31.8 (30.0) | 10-55 (30-30) |
| Litter cover (%) | 54.5 (65.0) | 0-88 (65-65) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>Isocoma menziesii</i> | 100 (100) | 20.5 (37) | 7 (37) | 37 (37) |
| <i>Eriogonum fasciculatum</i> | 75 (100) | 1.1 (3) | 0.2 (3) | 3 (3) |
| <i>Ambrosia salsola</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Artemisia californica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Astragalus trichopodus</i> var. <i>lonchus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Atriplex semibaccata</i> | 25 (-) | 8 (-) | 8 (-) | 8 (-) |
| <i>Baccharis pilularis</i> | 25 (100) | 10 (10) | 10 (10) | 10 (10) |
| <i>Baccharis sarothroides</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Cylindropuntia prolifera</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Gutierrezia californica</i> | 25 (-) | 4 (-) | 4 (-) | 4 (-) |
| <i>Lycium californicum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Salvia leucophylla</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 100 (100) | 10.1 (5) | 0.2 (5) | 30 (5) |
| <i>*Brassica nigra</i> | 75 (100) | 0.7 (1) | 0.2 (1) | 1 (1) |
| <i>*Centaurea melitensis</i> | 75 (100) | 5.4 (10) | 0.2 (10) | 10 (10) |
| <i>Ambrosia psilostachya</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 75

| Species Name | Con | Ave | Min | Max |
|---------------------------------------|----------|-----------|-----------|-----------|
| <i>*Erodium cicutarium</i> | 50 (-) | 7.5 (-) | 1 (-) | 14 (-) |
| <i>*Festuca myuros</i> | 50 (-) | 14 (-) | 3 (-) | 25 (-) |
| <i>Ambrosia acanthicarpa</i> | 25 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>*Bromus diandrus</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>*Bromus hordeaceus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Croton setiger</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Cynara cardunculus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Elymus triticoides</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Erodium</i> | 25 (-) | 4 (-) | 4 (-) | 4 (-) |
| <i>*Foeniculum vulgare</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Hedypnois cretica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Hypochaeris glabra</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Lamarckia aurea</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Lolium multiflorum</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>*Medicago polymorpha</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>*Mesembryanthemum crystallinum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Oxalis pes-caprae</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Raphanus sativus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Sonchus oleraceus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

Keckiella antirrhinoides Alliance

Keckiella antirrhinoides-Artemisia californica Association

Samples Used To Describe Association (n= 24)

Rapid Assessment(s), WR and WSD stands: SUNN.005, SDRP0440, SDRP0480, SDG0069, SDRP0135, CLNF.059, RECA.022, SDRP0084, SDRP0463, 0611TK01, 1478MB01, CLNF.060, SDRP0318, 1493PG01, RECA.012, SDRP0070, SDRP0429, 1352EB01, 1548JD01, 0520TK03, 0520AK02, SDRP0464, SDRP0263

Rapid Assessment(s), OC stands (n=1): 1279EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|------------|
| Total vegetation cover (%) | 55.2 (55.0) | 37-90 (55-55) | - |
| Herb cover (%) | 19.4 (-) | 3-50 (-) | 0-2 (-) |
| Shrub cover (%) | 52.7 (55.0) | 37-75 (55-55) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 0.5 (-) | 0.2-1 (-) | 2-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (15,0), NW (6,0), SE (2,1), Variable (1,0)

Macrotopography: bottom to mid 1/3 of slope (1,0), upper 1/3 of slope (3,0), middle to upper 1/3 of slope (1,0), middle 1/3 of slope (9,1), lower to upper 1/3 of slope (2,0), lower 1/3 of slope (2,0), lower to middle 1/3 of slope (6,0)

Microtopography: undulating (7,1), flat (2,0), Concave (3,0), convex (12,0)

Parent material: granitic (20,0), metamorphic (2,1), mixed igneous (1,0)

Soil texture: clay loam (6,0), loam (3,0), sandy loam (12,0), clay (1,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 76

| | Average | Range |
|-----------------------|------------|----------------------|
| Elevation (ft) | 967 (1975) | 302-2129 (1975-1975) |
| Slope (degrees) | 27 (38.0) | 4-45 (38-38) |
| Large rock cover (%) | 9.1 (-) | 0-48 (-) |
| Small rock cover (%) | 4.1 (-) | 0-18 (-) |
| Bare ground cover (%) | 30.2 (-) | 0-85 (-) |
| Litter cover (%) | 32.7 (-) | 0-83 (-) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---|-----------|-----------|-----------|---------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 10.8 (8) | 0.2 (8) | 45 (8) |
| <i>Keckiella antirrhinoides</i> | 96 (100) | 18.3 (36) | 4 (36) | 80 (36) |
| <i>Malosma laurina</i> | 75 (100) | 6.6 (1) | 1 (1) | 23 (1) |
| <i>Eriogonum fasciculatum</i> | 71 (-) | 3.8 (-) | 0.2 (-) | 14 (-) |
| <i>Salvia apiana</i> | 58 (100) | 2.6 (8) | 0.2 (8) | 15 (8) |
| <i>Mimulus aurantiacus</i> | 42 (100) | 1.4 (0.2) | 0.2 (0.2) | 4 (0.2) |
| <i>Adenostoma fasciculatum</i> | 33 (-) | 3.5 (-) | 0.2 (-) | 10 (-) |
| <i>Rhus ovata</i> | 29 (-) | 4.9 (-) | 0.2 (-) | 26 (-) |
| <i>Cneoridium dumosum</i> | 25 (-) | 4.5 (-) | 1 (-) | 9 (-) |
| <i>Rhamnus crocea</i> | 25 (-) | 2.7 (-) | 0.2 (-) | 10 (-) |
| <i>Sambucus mexicana</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Sambucus nigra subsp. caerulea</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 71 (-) | 6.3 (-) | 0.2 (-) | 18 (-) |
| <i>*Hirschfeldia incana</i> | 50 (-) | 2.4 (-) | 0.2 (-) | 7 (-) |
| <i>*Centaurea melitensis</i> | 29 (-) | 1.7 (-) | 0.2 (-) | 6 (-) |
| <i>*Hypochaeris glabra</i> | 25 (-) | 2.8 (-) | 0.2 (-) | 15 (-) |
| <i>Marah macrocarpus</i> var. | | | | |
| <i>macrocarpus</i> | 25 (-) | 1.3 (-) | 0.2 (-) | 7 (-) |
| <i>Phacelia cicutaria</i> var. <i>hispida</i> | 25 (-) | 2.1 (-) | 0.2 (-) | 7 (-) |

Lepidospartum squamatum Alliance

Lepidospartum squamatum (Alliance Only)

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1250EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 17.0 (17.0) | 17-17 (17-17) | - |
| Herb cover (%) | 10.0 (10.0) | 10-10 (10-10) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 7.0 (7.0) | 7-7 (7-7) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 77

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1)

Macrotopography: bottom (1,1)

Microtopography: undulating (1,1)

Parent material: mixed alluvium (1,1)

Soil texture: sand (1,1)

| | Average | Range |
|------------------------------|-------------|-----------------------|
| Elevation (ft) | 1152 (1152) | 1152-1152 (1152-1152) |
| Slope (degrees) | 2 (2.0) | 2-2 (2-2) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 75.0 (75.0) | 75-75 (75-75) |
| Bare ground cover (%) | 19.0 (19.0) | 19-19 (19-19) |
| Litter cover (%) | 5.0 (5.0) | 5-5 (5-5) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|-----------|-----------|-----------|-----------|
| Tree | <i>*Nicotiana glauca</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Platanus racemosa</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | | | | | |
| Shrub | <i>Acmispon glaber</i> | 100 (100) | 6 (6) | 6 (6) | 6 (6) |
| | <i>Acmispon heermannii</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Artemisia californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Astragalus trichopodus</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Baccharis salicifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Brickellia californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriophyllum confertiflorum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Lepidospartum squamatum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | <i>Acmispon strigosus</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>*Anagallis arvensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Artemisia douglasiana</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>*Bromus madritensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Calystegia macrostegia</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| | <i>*Centaurea melitensis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cryptantha muricata</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Hirschfeldia incana</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Hoita</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Logfia gallica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Lupinus hirsutissimus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Marrubium vulgare</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Melilotus albus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Melilotus indicus</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>*Oxalis californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Phacelia cicutaria</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Phacelia minor</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Plantago major</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Salvia columbariae</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| <i>Scrophularia californica</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Sonchus asper</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Stipa miliacea</i> var. <i>miliacea</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Trifolium obtusiflorum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Verbascum virgatum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Lepidospartum squamatum* Alliance**

***Lepidospartum squamatum*-*Baccharis salicifolia* Association**

Samples Used To Describe Association (n= 8)

Rapid Assessment(s), WR and WSD stands: BEAU.016, BEAU.008/574, LAEL.003/511, ANZA.069

Rapid Assessment(s), OC stands (n=4): 1056EK02, 1088EK01, 1092EK01, 1133EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 18.1 (19.3) | 11-28 (15-28) | - |
| Herb cover (%) | 5.5 (6.3) | 1-10 (3-8) | 0-5 (0-0.5) |
| Shrub cover (%) | 13.5 (13.0) | 8-25 (8-25) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 2-5 (-) |
| Overstory tree cover (%) | 2.0 (-) | 2-2 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,0), SW (3,1), S (3,3), SE (1,0)

Macrotopography: bottom (8,4)

Microtopography: Flat (5,1), undulating (3,3)

Parent material: granitic (1,0), mixed alluvium (1,1), sandstone (1,1), sandy alluvium (3,0)

Soil texture: sand (8,4)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 1338 (554) | 440-3219 (440-705) |
| Slope (degrees) | 1 (0.8) | 0-2 (0-1) |
| Large rock cover (%) | 1.6 (-) | 0-10 (-) |
| Small rock cover (%) | 66.3 (58.7) | 45-98 (45-75) |
| Bare ground cover (%) | 28.6 (40.0) | 0-53 (24-53) |
| Litter cover (%) | 1.4 (0.3) | 0-5 (0-1) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | * <i>Nicotiana glauca</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Platanus racemosa</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Schinus</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Baccharis salicifolia</i> | 100 (100) | 2.9 (0.9) | 0.2 (0.2) | 11 (2) |
| | <i>Lepidospartum squamatum</i> | 100 (100) | 7.9 (7.8) | 2 (2) | 19 (19) |
| | <i>Eriogonum fasciculatum</i> | 75 (100) | 0.8 (0.9) | 0.2 (0.2) | 3 (3) |
| | <i>Acmispon glaber</i> | 50 (75) | 2.8 (3.3) | 1 (1) | 6 (6) |
| | <i>Artemisia californica</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Lotus scoparius</i> | 50 (-) | 2.8 (-) | 1 (-) | 6 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 79

| Species Name | Con | Ave | Min | Max |
|--------------------------------------|----------|-----------|-----------|-----------|
| <i>Brickellia californica</i> | 38 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Acmispon heermannii</i> | 25 (50) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| <i>Bebbia juncea</i> | 25 (25) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Opuntia littoralis</i> | 25 (50) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Salvia apiana</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 88 (75) | 0.7 (0.2) | 0.2 (0.2) | 3 (0.2) |
| <i>*Brassica nigra</i> | 50 (100) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| <i>*Bromus diandrus</i> | 50 (100) | 1.3 (1.3) | 0.2 (0.2) | 2 (2) |
| <i>*Centaurea melitensis</i> | 50 (100) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| <i>*Erodium cicutarium</i> | 50 (75) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Phacelia minor</i> | 50 (75) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>*Sonchus asper</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Acmispon strigosus</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Ambrosia psilostachya</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Artemisia douglasiana</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Calystegia macrostegia</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Cryptantha muricata</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Hirschfeldia incana</i> | 38 (-) | 1.7 (-) | 0.2 (-) | 4 (-) |
| <i>*Hypochaeris glabra</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Logfia gallica</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lupinus bicolor</i> | 38 (75) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Anagallis arvensis</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Arundo donax</i> | 25 (25) | 2.6 (0.2) | 0.2 (0.2) | 5 (0.2) |
| <i>*Avena barbata</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Bromus tectorum</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| <i>Camissoniopsis bistorta</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Chaenactis glabriuscula</i> | 25 (25) | 1 (1) | 1 (1) | 1 (1) |
| <i>Corethrogyne filaginifolia</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Cryptantha intermedia</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Cynodon dactylon</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Heterotheca grandiflora</i> | 25 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lessingia filaginifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Lupinus hirsutissimus</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lupinus sparsiflorus</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Melilotus indicus</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Phacelia cicutaria</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium californicum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium stramineum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Scrophularia californica</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stachys</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Stipa miliacea var. miliacea</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Trifolium obtusiflorum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Lepidospartum squamatum* Alliance**

***Lepidospartum squamatum-Eriogonum fasciculatum* Association**

Samples Used To Describe Association (n= 7)

Rapid Assessment(s), WR and WSD stands: AGUA.008, AGUA.009/848, BEAU.004/BPU#222075/12, ANZA.014
Rapid Assessment(s), OC stands (n=3): 1088EK02, 1133EK04, 1142EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|---------------|
| Total vegetation cover (%) | 41.4 (30.3) | 20-60 (20-42) | - |
| Herb cover (%) | 16.4 (5.7) | 3-40 (3-8) | 0-1 (0-0.5) |
| Shrub cover (%) | 26.9 (24.7) | 17-43 (17-34) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 2-5 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (3,1), NW (1,0), SE (1,0), SW (2,2)

Macrotopography: bottom (5,2), bottom to lower 1/3 of slope (1,1), lower 1/3 of slope (1,0)

Microtopography: flat (6,2), undulating (1,1)

Parent material: granitic (3,0), mixed alluvium (2,2), sandstone (1,1), sandy alluvium (1,0)

Soil texture: sandy loam (2,0), sand (5,3)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 2154 (672) | 551-3828 (551-820) |
| Slope (degrees) | 1 (0.7) | 0-3 (0-1) |
| Large rock cover (%) | 1.1 (-) | 0-6 (-) |
| Small rock cover (%) | 54.3 (39.0) | 22-84 (22-65) |
| Bare ground cover (%) | 33.6 (52.7) | 1-77 (25-77) |
| Litter cover (%) | 8.6 (6.7) | 0-18 (0-12) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--------------------------------|-----------|-----------|-----------|---------|
| Shrub | | | | |
| <i>Eriogonum fasciculatum</i> | 100 (100) | 9.4 (14) | 2 (8) | 25 (25) |
| <i>Lepidospartum squamatum</i> | 100 (100) | 8.4 (4) | 1 (1) | 22 (7) |
| <i>Acmispon glaber</i> | 57 (67) | 1.6 (1.1) | 0.2 (0.2) | 3 (2) |
| <i>Lotus scoparius</i> | 57 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| <i>Hesperoyucca whipplei</i> | 43 (100) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Opuntia littoralis</i> | 43 (100) | 1.7 (1.7) | 0.2 (0.2) | 3 (3) |
| <i>Opuntia parryi</i> | 43 (-) | 6.3 (-) | 3 (-) | 11 (-) |
| <i>Sambucus mexicana</i> | 43 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Artemisia californica</i> | 29 (67) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Baccharis salicifolia</i> | 29 (67) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Brickellia californica</i> | 29 (67) | 1.6 (1.6) | 0.2 (0.2) | 3 (3) |
| <i>Malosma laurina</i> | 29 (67) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| <i>Prunus ilicifolia</i> | 29 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| <i>Ribes quercetorum</i> | 29 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| <i>Salvia apiana</i> | 29 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Senecio flaccidus</i> | 29 (-) | 2 (-) | 2 (-) | 2 (-) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 71 (100) | 1.5 (0.5) | 0.2 (0.2) | 4 (1) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 81

| Species Name | Con | Ave | Min | Max |
|--------------------------------|---------|-----------|-----------|-----------|
| <i>*Bromus diandrus</i> | 57 (33) | 2.6 (0.2) | 0.2 (0.2) | 4 (0.2) |
| <i>Cryptantha intermedia</i> | 57 (67) | 2.1 (2.1) | 0.2 (0.2) | 4 (4) |
| <i>*Avena barbata</i> | 43 (67) | 6.8 (0.2) | 0.2 (0.2) | 20 (0.2) |
| <i>Camissonia bistorta</i> | 43 (-) | 2.1 (-) | 0.2 (-) | 5 (-) |
| <i>Camissoniopsis bistorta</i> | 43 (33) | 2.1 (0.2) | 0.2 (0.2) | 5 (0.2) |
| <i>*Hirschfeldia incana</i> | 43 (33) | 1.1 (1) | 0.2 (1) | 2 (1) |
| <i>*Brassica nigra</i> | 29 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Bromus tectorum</i> | 29 (-) | 13 (-) | 12 (-) | 14 (-) |
| <i>*Centaurea melitensis</i> | 29 (67) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| <i>*Erodium cicutarium</i> | 29 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Festuca myuros</i> | 29 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Filago gallica</i> | 29 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| <i>*Hypochaeris glabra</i> | 29 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Logfia gallica</i> | 29 (33) | 1.6 (0.2) | 0.2 (0.2) | 3 (0.2) |
| <i>Lotus strigosus</i> | 29 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Lupinus concinnus</i> | 29 (-) | 1.5 (-) | 1 (-) | 2 (-) |

***Lycium californicum* Alliance**
***Lycium californicum* Association**

Samples Used To Describe Association (n= 8)

Rapid Assessment(s), WR and WSD stands: 1710JD01, 0312TK22, 0312RB21, 0312TK21, 0503AK21, 0707TK04
Rapid Assessment(s), OC stands (n=2): 1071EK01, 1406EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|---------------|
| Total vegetation cover (%) | 40.8 (34.5) | 24-58 (24-45) | - |
| Herb cover (%) | 20.6 (2.1) | 0.2-44 (0.2-4) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 32.5 (32.5) | 23-42 (23-42) | 0-2 (0-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,0), SE (2,0), SW (4,2), Variable (1,0)

Macrotopography: bottom to lower 1/3 of slope (1,0), lower 1/3 of slope (3,1), lower to middle 1/3 of slope (1,1), lower to upper 1/3 of slope (1,0), middle to upper 1/3 of slope (1,0), bottom (1,0)

Microtopography: concave (1,1), convex (2,0), flat (1,0), undulating (4,1)

Parent material: mixed alluvium (1,0), sandstone (3,2), sandy alluvium (2,0), siltstone (1,0)

Soil texture: sandy loam (3,0), sand (2,2), clay (1,0), clay loam (2,0)

| | Average | Range |
|-----------------------|-------------|--------------|
| Elevation (ft) | 35 (5) | 0-148 (0-10) |
| Slope (degrees) | 23 (65.0) | 0-70 (60-70) |
| Large rock cover (%) | 0.6 (2.5) | 0-5 (0-5) |
| Small rock cover (%) | 4.8 (15.0) | 0-20 (10-20) |
| Bare ground cover (%) | 45.3 (51.5) | 0-86 (49-54) |
| Litter cover (%) | 42.6 (29.5) | 5-85 (20-39) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|---|-----------|-------------|-----------|-----------|
| Shrub | <i>Lycium californicum</i> | 100 (100) | 11.9 (12.5) | 2 (5) | 23 (20) |
| | <i>Cylindropuntia prolifera</i> | 88 (100) | 2.8 (0.2) | 0.2 (0.2) | 8 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 63 (50) | 3.6 (3) | 0.2 (3) | 7 (3) |
| | <i>Artemisia californica</i> | 38 (-) | 2.7 (-) | 1 (-) | 4 (-) |
| | * <i>Atriplex semibaccata</i> | 38 (50) | 0.5 (1) | 0.2 (1) | 1 (1) |
| | <i>Encelia californica</i> | 38 (100) | 0.7 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>Isocoma menziesii</i> | 38 (50) | 1.5 (0.2) | 0.2 (0.2) | 4 (0.2) |
| | <i>Opuntia littoralis</i> | 25 (50) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Peritoma arborea</i> var. <i>arborea</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhus integrifolia</i> | 25 (100) | 15.1 (15.1) | 0.2 (0.2) | 30 (30) |
| Herb | * <i>Bromus madritensis</i> | 63 (-) | 19.8 (-) | 4 (-) | 30 (-) |
| | <i>Distichlis spicata</i> | 63 (50) | 1.1 (3) | 0.2 (3) | 3 (3) |
| | <i>Arthrocnemum subterminale</i> | 50 (-) | 3.3 (-) | 0.2 (-) | 7 (-) |
| | <i>Frankenia salina</i> | 50 (-) | 2.8 (-) | 0.2 (-) | 7 (-) |
| | * <i>Oxalis pes-caprae</i> | 38 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Amblyopappus pusillus</i> | 25 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| | <i>Antirrhinum nuttallianum</i> | 25 (-) | 5 (-) | 2 (-) | 8 (-) |
| | * <i>Avena barbata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Bromus diandrus</i> | 25 (-) | 6 (-) | 4 (-) | 8 (-) |
| | * <i>Centaurea melitensis</i> | 25 (-) | 3 (-) | 2 (-) | 4 (-) |
| | <i>Cressa truxillensis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Deinandra fasciculata</i> | 25 (-) | 3.5 (-) | 3 (-) | 4 (-) |
| | * <i>Festuca myuros</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Pterostegia drymarioides</i> | 25 (-) | 4 (-) | 3 (-) | 5 (-) |
| | * <i>Salsola tragus</i> | 25 (50) | 1.1 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | * <i>Sonchus oleraceus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Sporobolus airoides</i> | 25 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Suaeda taxifolia</i> | 25 (100) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |

***Malacothamnus fasciculatus* Alliance** ***Malacothamnus fasciculatus* Association**

Samples Used To Describe Association (n= 11)

Rapid Assessment(s), WR and WSD stands: 1526MM02, 1817TK01, 1525EB01, 0422RB02, 1531EB03

Rapid Assessment(s), OC stands (n=6): 1204EK02, 1204EK03, 1253EK01, 1289MA02, 1298EK01, 1426EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|------------|
| Total vegetation cover (%) | 52.4 (52.0) | 31-70 (31-70) | - |
| Herb cover (%) | 10.6 (6.0) | 0.2-30 (0.2-21) | 0-1 (0-1) |
| Shrub cover (%) | 47.8 (47.8) | 30-67 (30-67) | 1-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 2-5 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (2,1), NW (1,0), S (2,2), SE (4,1), W (1,1), N (1,1)

Macrotopography: lower to upper 1/3 of slope (1,0), middle 1/3 of slope (3,3), lower to middle 1/3 of slope (1,1), lower 1/3 of slope (2,0), bottom to lower 1/3 of slope (1,0), bottom (1,0), middle to upper 1/3 of slope (2,2)

Microtopography: undulating (3,1), concave (2,1), convex (1,0), flat (5,4)

Parent material: diorite (1,0), sandstone (5,5)

Soil texture: clay loam (4,3), loam (2,2), sandy loam (5,1)

| | Average | Range |
|------------------------------|----------------|---------------------|
| Elevation (ft) | 755 (904) | 128-1270 (538-1270) |
| Slope (degrees) | 23 (30.8) | 0-48 (9-48) |
| Large rock cover (%) | 9.1 (-) | 0-45 (-) |
| Small rock cover (%) | 16.6 (1.1) | 0-67 (0-3) |
| Bare ground cover (%) | 33.8 (36.0) | 2-79 (12-52) |
| Litter cover (%) | 38.6 (60.4) | 2-82 (45-82) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------------|------------|-------------|------------|------------|
| Shrub | <i>Malacothamnus fasciculatus</i> | 100 (100) | 27.5 (28.8) | 10 (10) | 60 (60) |
| | <i>Malosma laurina</i> | 100 (100) | 3.8 (1.9) | 0.2 (0.2) | 9 (5) |
| | <i>Artemisia californica</i> | 91 (100) | 2.7 (3.2) | 0.2 (1) | 8 (8) |
| | <i>Salvia mellifera</i> | 73 (100) | 2.5 (0.7) | 0.2 (0.2) | 15 (1) |
| | <i>Acmispon glaber</i> | 55 (83) | 10.3 (11.4) | 5 (5) | 22 (22) |
| | <i>Rhus integrifolia</i> | 36 (50) | 1.3 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>Salvia apiana</i> | 36 (17) | 0.4 (1) | 0.2 (1) | 1 (1) |
| | <i>Eriogonum fasciculatum</i> | 27 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 27 (33) | 4.7 (6.5) | 1 (1) | 12 (12) |
| | <i>Mimulus aurantiacus</i> | 27 (17) | 2.1 (1) | 0.2 (1) | 5 (1) |
| | <i>Sambucus nigra subsp. caerulea</i> | 27 (50) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| Herb | <i>*Bromus madritensis</i> | 45 (50) | 3.6 (3.1) | 0.2 (0.2) | 7 (5) |
| | <i>*Hirschfeldia incana</i> | 45 (67) | 0.8 (0.2) | 0.2 (0.2) | 3 (0.2) |
| | <i>*Centaurea melitensis</i> | 36 (33) | 1.5 (2) | 1 (2) | 2 (2) |
| | <i>Calystegia macrostegia</i> | 27 (-) | 1.1 (-) | 0.2 (-) | 3 (-) |

***Malosma laurina* Alliance**

***Malosma laurina* (Alliance Only)**

Samples Used To Describe Association (n= 5)

Rapid Assessment(s), WR and WSD stands: CORN.005/436, CLNF.009, CLNF.120

Rapid Assessment(s), OC stands (n=2): 1110EK05, 1425MA01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|-------------|
| Total vegetation cover (%) | 61.6 (76.0) | 35-77 (75-77) | - |
| Herb cover (%) | 22.4 (25.6) | 0.2-51 (0.2-51) | 0-1 (0-1) |
| Shrub cover (%) | 39.6 (47.5) | 20-75 (20-75) | 0.5-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 2-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,0), N (1,1), NE (1,0), NW (1,0), SE (1,1)
Macrotopography: upper 1/3 of slope (3,1), bottom (1,0), lower to middle 1/3 of slope (1,1)
Microtopography: concave (1,1), Convex (1,0), Flat (1,0), undulating (2,1)
Parent material: granitic (1,0), sandstone (1,1), sedimentary (1,0), silty alluvium (1,0)
Soil texture: loam (1,0), clay loam (1,1), sandy loam (3,1)

| | Average | Range |
|-----------------------|-------------|--------------------|
| Elevation (ft) | 1331 (727) | 702-2949 (702-751) |
| Slope (degrees) | 19 (23.0) | 1-26 (20-26) |
| Large rock cover (%) | 12.0 (-) | 0-45 (-) |
| Small rock cover (%) | 15.6 (1.0) | 0-38 (0-2) |
| Bare ground cover (%) | 20.4 (18.5) | 5-32 (5-32) |
| Litter cover (%) | 47.6 (77.0) | 5-89 (65-89) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------|-----------|-----------|-----------|-----------|
| Shrub | <i>Malosma laurina</i> | 100 (100) | 21.6 (31) | 11 (12) | 50 (50) |
| | <i>Artemisia californica</i> | 40 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Heteromeles arbutifolia</i> | 40 (50) | 10.5 (15) | 6 (15) | 15 (15) |
| | <i>Mimulus aurantiacus</i> | 40 (50) | 18.5 (30) | 7 (30) | 30 (30) |
| | <i>Rhus integrifolia</i> | 40 (100) | 3 (3) | 2 (2) | 4 (4) |
| | <i>Romneya coulteri</i> | 40 (-) | 3.6 (-) | 0.2 (-) | 7 (-) |
| | <i>Salvia mellifera</i> | 40 (-) | 1.6 (-) | 0.2 (-) | 3 (-) |
| Herb | <i>*Bromus madritensis</i> | 40 (50) | 6.5 (12) | 1 (12) | 12 (12) |
| | <i>*Centaurea melitensis</i> | 40 (50) | 4.5 (7) | 2 (7) | 7 (7) |
| | <i>Elymus condensatus</i> | 40 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Erodium cicutarium</i> | 40 (-) | 3.6 (-) | 0.2 (-) | 7 (-) |
| | <i>*Hirschfeldia incana</i> | 40 (-) | 3.6 (-) | 0.2 (-) | 7 (-) |
| | <i>Marah macrocarpus</i> | 40 (-) | 4 (-) | 2 (-) | 6 (-) |

***Malosma laurina* Alliance**

***Malosma laurina-Acmispon glaber* Association**

Samples Used To Describe Association (n= 37)

Rapid Assessment(s), WR and WSD stands: SDRP0427, 1529MM01, 1604EB01, 0422RB01, 0422AK04, SDRP0477, 1393KG02, SDG0020, 0506TK02, 1539PJ01, 0422RB03, 0611AK01, 1327KH01, 1069FS02, 0519AK01, 0507DH02, 0519RB01, SDG0103, 0422AK02, 1216MM01, SDG0052, SDG0039, SDG0008, SDG0116, SDG0031, SDG0040, SDG0108, 1536EB01, 0421RB01, SDG0109, 1539PG02, SDG0048, 0603JJ02, 1216MB01, 1538JD01, 0423AK01

Rapid Assessment(s), OC stands (n=1): 1218EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 42.0 (34.0) | 10-85 (34-34) | - |
| Herb cover (%) | 22.9 (6.0) | 2-65 (6-6) | 0-1 (0.5-1) |
| Shrub cover (%) | 28.0 (28.0) | 28-28 (28-28) | 0-5 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 5-10 (-) |
| Overstory tree cover (%) | 2.1 (-) | 0.2-5 (-) | 2-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SE (14,0), SW (11,1), NW (4,0), NE (7,0), Flat (1,0)
Macrotopography: middle 1/3 of slope to ridgetop (2,0), upper 1/3 of slope (2,0), ridge top (1,0), middle to upper 1/3 of slope (5,1), middle 1/3 of slope (12,0), lower to upper 1/3 of slope (4,0), lower to middle 1/3 of slope (2,0), bottom to lower 1/3 of slope (1,0), upper 1/3 of slope to ridgetop (1,0), lower 1/3 of slope (6,0)
Microtopography: flat (2,0), undulating (22,0), convex (6,0), concave (6,1)
Parent material: gabbro (2,0), granitic (16,0), metamorphic (1,0), phyllite (1,0), quartz diorite (1,0), sandstone (1,1), sedimentary (1,0)
Soil texture: clay loam (7,0), loam (3,0), sand (1,0), sandy loam (18,1), silty loam (1,0), clay (1,0)

| | Average | Range |
|-----------------------|-------------|----------------------|
| Elevation (ft) | 1019 (1467) | 371-1654 (1467-1467) |
| Slope (degrees) | 20 (28.0) | 0-45 (28-28) |
| Large rock cover (%) | 9.0 (-) | 0-63 (-) |
| Small rock cover (%) | 17.9 (2.0) | 0-75 (2-2) |
| Bare ground cover (%) | 30.4 (56.0) | 0-94 (56-56) |
| Litter cover (%) | 21.7 (40.0) | 0-82 (40-40) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|---------|
| Shrub | | | | |
| <i>Malosma laurina</i> | 100 (100) | 8.9 (10) | 2 (10) | 23 (10) |
| <i>Artemisia californica</i> | 76 (100) | 1.2 (1) | 0.2 (1) | 5 (1) |
| <i>Acmispon glaber</i> | 65 (100) | 8 (8) | 0.2 (8) | 26 (8) |
| <i>Salvia apiana</i> | 62 (100) | 1.1 (0.2) | 0.2 (0.2) | 4 (0.2) |
| <i>Eriogonum fasciculatum</i> | 51 (-) | 1.7 (-) | 0.2 (-) | 8 (-) |
| <i>Hesperoyucca whipplei</i> | 38 (100) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Rhamnus crocea</i> | 35 (-) | 2 (-) | 0.2 (-) | 15 (-) |
| <i>Mimulus aurantiacus</i> | 30 (100) | 0.6 (0.2) | 0.2 (0.2) | 4 (0.2) |
| <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 27 (-) | 1.9 (-) | 0.2 (-) | 6 (-) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 57 (-) | 6.4 (-) | 1 (-) | 15 (-) |
| * <i>Centaurea melitensis</i> | 49 (100) | 2.4 (1) | 0.2 (1) | 10 (1) |
| * <i>Hirschfeldia incana</i> | 43 (-) | 6.3 (-) | 0.2 (-) | 50 (-) |
| * <i>Festuca myuros</i> | 41 (-) | 3.7 (-) | 0.2 (-) | 12 (-) |
| * <i>Avena barbata</i> | 35 (-) | 5 (-) | 0.2 (-) | 23 (-) |
| <i>Calystegia macrostegia</i> | 32 (-) | 4 (-) | 0.2 (-) | 18 (-) |
| * <i>Hypochaeris glabra</i> | 30 (-) | 5.1 (-) | 0.2 (-) | 20 (-) |
| <i>Marah macrocarpus</i> var. <i>macrocarpus</i> | 27 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |

***Malosma laurina* Alliance**

***Malosma laurina-Artemisia californica* Association**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1218EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 35.0 (35.0) | 35-35 (35-35) | - |
| Herb cover (%) | 3.0 (3.0) | 3-3 (3-3) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 32.0 (32.0) | 32-32 (32-32) | 1-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1)

Macrotopography:

Microtopography: flat (1,1)

Parent material: sandstone (1,1)

Soil texture: sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 1368 (1368) | 1368-1368 (1368-1368) |
| Slope (degrees) | 32 (32.0) | 32-32 (32-32) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 2.0 (2.0) | 2-2 (2-2) |
| Bare ground cover (%) | 66.0 (66.0) | 66-66 (66-66) |
| Litter cover (%) | 30.0 (30.0) | 30-30 (30-30) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>Acmispon glaber</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Artemisia californica</i> | 100 (100) | 15 (15) | 15 (15) | 15 (15) |
| <i>Cylindropuntia prolifera</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Eriogonum fasciculatum</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Malacothamnus fasciculatus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Malosma laurina</i> | 100 (100) | 15 (15) | 15 (15) | 15 (15) |
| <i>Mimulus aurantiacus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Opuntia littoralis</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Salvia mellifera</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Bromus diandrus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus madritensis</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |

***Opuntia littoralis* Alliance**

***Opuntia littoralis-Eriogonum fasciculatum-Malosma laurina* Association**

Samples Used To Describe Association (n= 7)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=7): 1086EK01, 1092EK02, 1347EK01, 1463EK02, 1472EK01, 1479EK01, 1513EK06

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|-----------------|-------------------|
| Total vegetation cover (%) | 34.7 (34.7) | 18-65 (18-65) | - |
| Herb cover (%) | 4.7 (4.7) | 0.2-10 (0.2-10) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 31.0 (31.0) | 15-65 (15-65) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 2.1 (2.1) | 0.2-4 (0.2-4) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (2,2), S (4,4), SE (1,1)

Macrotopography: lower to middle 1/3 of slope (1,1), middle 1/3 of slope (3,3), lower 1/3 of slope (1,1), bottom to lower 1/3 of slope (1,1), upper 1/3 of slope (1,1)

Microtopography: undulating (5,5), flat (2,2)

Parent material: sandstone (7,7)

Soil texture: sand (3,3), sandy loam (2,2)

| | Average | Range |
|------------------------------|----------------|-----------------|
| Elevation (ft) | 582 (582) | 0-1056 (0-1056) |
| Slope (degrees) | 27 (27.3) | 1-75 (1-75) |
| Large rock cover (%) | 10.7 (10.7) | 0-64 (0-64) |
| Small rock cover (%) | 3.2 (3.2) | 0-10 (0-10) |
| Bare ground cover (%) | 50.3 (50.3) | 25-67 (25-67) |
| Litter cover (%) | 33.8 (33.8) | 8-65 (8-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------------|------------|-------------|------------|------------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 29 (29) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| Shrub | | | | | |
| | <i>Eriogonum fasciculatum</i> | 100 (100) | 5 (5) | 1 (1) | 14 (14) |
| | <i>Opuntia littoralis</i> | 86 (86) | 9 (9) | 4 (4) | 20 (20) |
| | <i>Rhus integrifolia</i> | 71 (71) | 2 (2) | 1 (1) | 5 (5) |
| | <i>Artemisia californica</i> | 57 (57) | 0.9 (0.9) | 0.2 (0.2) | 2 (2) |
| | <i>Malosma laurina</i> | 57 (57) | 7.1 (7.1) | 0.2 (0.2) | 25 (25) |
| | <i>Sambucus nigra subsp. caerulea</i> | 57 (57) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Acmispon glaber</i> | 43 (43) | 3.4 (3.4) | 0.2 (0.2) | 8 (8) |
| | <i>Hesperoyucca whipplei</i> | 43 (43) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Baccharis pilularis</i> | 29 (29) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>Brickellia californica</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cylindropuntia prolifera</i> | 29 (29) | 3.5 (3.5) | 2 (2) | 5 (5) |
| | <i>Encelia californica</i> | 29 (29) | 13.5 (13.5) | 1 (1) | 26 (26) |
| | <i>Mimulus aurantiacus</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Salvia mellifera</i> | 29 (29) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |

| | Species Name | Con | Ave | Min | Max |
|------|--------------------------------------|---------|-----------|-----------|-----------|
| Herb | <i>*Avena barbata</i> | 43 (43) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| | <i>*Bromus diandrus</i> | 43 (43) | 3 (3) | 1 (1) | 6 (6) |
| | <i>*Brassica nigra</i> | 29 (29) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>*Bromus madritensis</i> | 29 (29) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>*Carduus pycnocephalus</i> | 29 (29) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>*Centaurea melitensis</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Crassula connata</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Dudleya pulverulenta</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Logfia gallica</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Pseudognaphalium californicum</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Scrophularia californica</i> | 29 (29) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Quercus berberidifolia* Alliance** ***Quercus berberidifolia* Association**

Samples Used To Describe Association (n= 33)

Rapid Assessment(s), WR and WSD stands: 1696LS01, 1710JD12, 1702MB01, 0609AK03, SDG0053, 0506RB02, 0506RB23, 0609RB01, 1737JD04, 1304JD02, 1710JD04, 1561JD01, 1384JD01, 0610TK03, 1607MM01, SDRP0197, 1361KH01, 1613MB01, 1817TK02, 0610DH03, 1710JD09, 1702PG01, 1710JD07, 1607MM03, 1038MB01, 1645FS02

Rapid Assessment(s), OC stands (n=7): 1031EK01, 1090EK01, 1092EK04, 1100MA01, 1128MA01, 1187EK03, 1510EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 60.4 (58.7) | 28-88 (39-88) | - |
| Herb cover (%) | 7.2 (2.2) | 0-60 (0.2-6) | 0-2 (0-1) |
| Shrub cover (%) | 57.3 (57.3) | 39-85 (39-85) | 0-10 (0-10) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-2 (-) |
| Overstory tree cover (%) | 2.0 (4.0) | 1-4 (4-4) | 2-15 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: W (1,1), N (4,4), NE (11,1), NW (12,1), SE (2,0), SW (2,0), Variable (1,0)

Macrotopography: lower to upper 1/3 of slope (4,1), middle 1/3 of slope (4,1), upper 1/3 of slope (5,1), middle to upper 1/3 of slope (7,3), middle 1/3 of slope to ridgetop (1,0), lower 1/3 of slope (1,0), entire slope (1,0), bottom to upper 1/3 of slope (2,0), bottom (1,0), lower to middle 1/3 of slope (6,1)

Microtopography: convex (4,0), flat (10,2), undulating (13,2), concave (6,3)

Parent material: diorite (1,0), gabbro (2,0), granitic (5,0), igneous (1,0), metamorphic (5,0), sandstone (5,5)

Soil texture: clay (1,0), clay loam (8,4), loam (4,1), sandy loam (19,2), silty loam (1,0)

| | Average | Range |
|------------------------------|-------------|---------------------|
| Elevation (ft) | 1082 (786) | 213-3153 (499-1145) |
| Slope (degrees) | 21 (28.9) | 4-50 (12-50) |
| Large rock cover (%) | 4.9 (-) | 0-40 (-) |
| Small rock cover (%) | 15.6 (2.9) | 0-64 (0-10) |
| Bare ground cover (%) | 18.6 (18.4) | 1-65 (5-45) |
| Litter cover (%) | 55.0 (77.0) | 0-90 (43-90) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|----------|-----------|-----------|----------|
| Shrub | <i>Quercus berberidifolia</i> | 61 (100) | 45.1 (43) | 15 (24) | 70 (65) |
| | <i>Heteromeles arbutifolia</i> | 55 (57) | 2.7 (5.3) | 0.2 (1) | 12 (12) |
| | <i>Rhus integrifolia</i> | 55 (100) | 4.2 (7.2) | 0.2 (0.2) | 20 (20) |
| | <i>Adenostoma fasciculatum</i> | 42 (14) | 3.2 (0.2) | 0.2 (0.2) | 15 (0.2) |
| | <i>Mimulus aurantiacus</i> | 39 (57) | 0.9 (0.9) | 0.2 (0.2) | 3 (2) |
| | <i>Quercus acutidens</i> | 39 (-) | 33.6 (-) | 6 (-) | 75 (-) |
| | <i>Lonicera subspicata</i> | 30 (-) | 1.2 (-) | 0.2 (-) | 5 (-) |
| | <i>Toxicodendron diversilobum</i> | 30 (29) | 2.4 (2.6) | 0.2 (0.2) | 6 (5) |
| | <i>Acmispon glaber</i> | 27 (29) | 2.1 (0.2) | 0.2 (0.2) | 7 (0.2) |
| | <i>Artemisia californica</i> | 27 (71) | 2.5 (2.5) | 0.2 (0.2) | 8 (5) |
| | <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 27 (-) | 1.4 (-) | 0.2 (-) | 6 (-) |
| Herb | <i>Marah macrocarpus</i> var. <i>macrocarpus</i> | 30 (-) | 0.6 (-) | 0.2 (-) | 3 (-) |
| | <i>*Festuca myuros</i> | 27 (-) | 2.7 (-) | 0.2 (-) | 7 (-) |
| | <i>Stipa lepida</i> | 27 (29) | 0.5 (1.1) | 0.2 (0.2) | 2 (2) |
| | | | | | |

Quercus berberidifolia Alliance

Quercus berberidifolia-*Arctostaphylos glandulosa* Association

Samples Used To Describe Association (n= 2)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=2): 1180MA01, 1215EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|---------------|
| Total vegetation cover (%) | 77.0 (77.0) | 74-80 (74-80) | - |
| Herb cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0.5-1 (0.5-1) |
| Shrub cover (%) | 80.0 (80.0) | 80-80 (80-80) | 2-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1), S (1,1)

Macrotopography: upper 1/3 of slope (1,1), upper 1/3 of slope to ridgetop (1,1)

Microtopography: undulating (1,1), concave (1,1)

Parent material: granitic (1,1)

Soil texture: clay loam (1,1), sand (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3596 (3596) | 3596-3596 (3596-3596) |
| Slope (degrees) | 40 (40.0) | 40-40 (40-40) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 2.5 (2.5) | 0-5 (0-5) |
| Bare ground cover (%) | 12.0 (12.0) | 7-17 (7-17) |
| Litter cover (%) | 82.5 (82.5) | 80-85 (80-85) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--|-----------|-------------|-----------|-----------|
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 6.5 (6.5) | 5 (5) | 8 (8) |
| | <i>Arctostaphylos glandulosa</i> | 100 (100) | 12.5 (12.5) | 10 (10) | 15 (15) |
| | <i>Heteromeles arbutifolia</i> | 100 (100) | 4 (4) | 1 (1) | 7 (7) |
| | <i>Quercus berberidifolia</i> | 100 (100) | 45 (45) | 35 (35) | 55 (55) |
| | <i>Rhus ovata</i> | 100 (100) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>Acmispon glaber</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Ceanothus crassifolius</i> | 50 (50) | 5 (5) | 5 (5) | 5 (5) |
| | <i>Ceanothus leucodermis</i> | 50 (50) | 9 (9) | 9 (9) | 9 (9) |
| | <i>Cercocarpus betuloides</i> var. <i>betuloides</i> | 50 (50) | 7 (7) | 7 (7) | 7 (7) |
| | <i>Clematis lasiantha</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Eriophyllum confertiflorum</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Garrya flavescens</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Hesperoyucca whipplei</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhamnus ilicifolia</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Salvia mellifera</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| Herb | <i>Stipa coronata</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Quercus berberidifolia Alliance

Quercus berberidifolia-*Ceanothus tomentosus* Association

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1363EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|------------|
| Total vegetation cover (%) | 71.0 (71.0) | 71-71 (71-71) | - |
| Herb cover (%) | - (-) | - (-) | - (-) |
| Shrub cover (%) | - (-) | - (-) | - (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,1)

Macrotopography: upper 1/3 of slope (1,1)

Microtopography: flat (1,1)

Parent material: sandstone (1,1)

Soil texture: silty loam (1,1)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 91

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 2047 (2047) | 2047-2047 (2047-2047) |
| Slope (degrees) | 5 (5.0) | 5-5 (5-5) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 7.0 (7.0) | 7-7 (7-7) |
| Bare ground cover (%) | 12.0 (12.0) | 12-12 (12-12) |
| Litter cover (%) | 78.0 (78.0) | 78-78 (78-78) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>Acmispon glaber</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Adenostoma fasciculatum</i> | 100 (100) | 5 (5) | 5 (5) | 5 (5) |
| <i>Ceanothus tomentosus</i> | 100 (100) | 15 (15) | 15 (15) | 15 (15) |
| <i>Cercocarpus betuloides</i> var. <i>betuloides</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>Eriodictyon crassifolium</i> | 100 (100) | 3 (3) | 3 (3) | 3 (3) |
| <i>Heteromeles arbutifolia</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| <i>Malosma laurina</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Quercus berberidifolia</i> | 100 (100) | 40 (40) | 40 (40) | 40 (40) |
| <i>Rhus ovata</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Quercus berberidifolia* Alliance**

***Quercus berberidifolia*-*Fraxinus dipetala*-*Heteromeles arbutifolia* Association**

Samples Used To Describe Association (n= 17)

Rapid Assessment(s), WR and WSD stands: CORS.004/404/BPU 19994-239, CAHUI.028/265, CAHUI.002, CLNF.038, ANZA.122/165, CORS.010/(BPU 34102-295), CLNF.013, CAHUI.027/106, CORS.008/BPU, CLNF.066, CLNF.071, CLNF.040, CLNF.044

Rapid Assessment(s), OC stands (n=4): 1179EK01, 1179MA01, 1189EK01, 1294EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|------------------|-------------|
| Total vegetation cover (%) | 70.9 (66.3) | 51-90 (60-77) | - |
| Herb cover (%) | 10.3 (0.2) | 0.2-50 (0.2-0.2) | 0-1 (0-0.5) |
| Shrub cover (%) | 63.1 (53.5) | 14-89 (14-77) | 0.5-5 (1-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | 55.0 (55.0) | 55-55 (55-55) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: N (2,2), NE (8,1), NW (6,1), Variable (1,0)

Macrotopography: middle 1/3 of slope (4,1), middle 1/3 of slope to ridgetop (1,1), lower to upper 1/3 of slope (1,1), upper 1/3 of slope (11,1)

Microtopography: Convex (2,0), Undulating (15,4)

Parent material: granitic (5,0), metamorphic (3,2), sandstone (2,2), sedimentary (7,0)

Soil texture: clay loam (2,1), loam (4,0), sand (1,0), sandy loam (6,0), unknown (1,0)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 92

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3187 (2720) | 1224-4495 (1224-3287) |
| Slope (degrees) | 31 (36.0) | 18-40 (26-40) |
| Large rock cover (%) | 2.6 (-) | 0-22 (-) |
| Small rock cover (%) | 10.2 (1.3) | 0-60 (0-5) |
| Bare ground cover (%) | 16.9 (20.5) | 3-33 (5-32) |
| Litter cover (%) | 65.5 (75.0) | 15-90 (65-87) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|----------------------------------|-----------|-------------|-----------|-----------|
| Shrub | | | | |
| <i>Fraxinus dipetala</i> | 100 (100) | 7.3 (9.8) | 0.2 (0.2) | 20 (20) |
| <i>Quercus berberidifolia</i> | 100 (100) | 39.4 (47.3) | 15 (35) | 75 (65) |
| <i>Heteromeles arbutifolia</i> | 88 (100) | 6.3 (4) | 0.2 (3) | 15 (6) |
| <i>Adenostoma fasciculatum</i> | 65 (50) | 1.8 (1.1) | 0.2 (0.2) | 4 (2) |
| <i>Cercocarpus betuloides</i> | 53 (-) | 6.4 (-) | 0.2 (-) | 23 (-) |
| <i>Arctostaphylos glandulosa</i> | 41 (25) | 1.1 (3) | 0.2 (3) | 3 (3) |
| <i>Garrya flavescens</i> | 41 (50) | 6.1 (0.2) | 0.2 (0.2) | 30 (0.2) |
| <i>Rhamnus ilicifolia</i> | 35 (-) | 2.9 (-) | 0.2 (-) | 7 (-) |
| <i>Rhus ovata</i> | 35 (50) | 2.2 (2.5) | 0.2 (2) | 6 (3) |
| <i>Keckiella cordifolia</i> | 29 (75) | 1.4 (1.4) | 0.2 (0.2) | 3 (3) |
| <i>Lonicera subspicata</i> | 29 (25) | 1.8 (2) | 0.2 (2) | 3 (2) |
| <i>Mimulus aurantiacus</i> | 29 (25) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>*Bromus madritensis</i> | 29 (-) | 1.5 (-) | 0.2 (-) | 3 (-) |

***Quercus berberidifolia*-*Adenostoma fasciculatum* Alliance**
***Quercus berberidifolia*-*Adenostoma fasciculatum* Association**

Samples Used To Describe Association (n= 12)

Rapid Assessment(s), WR and WSD stands: SRPL.020, CLNF.109, POTR.002/374, SAJAC.001/62/BPU#595578/143, BLACK.051, WILD.004/506/BPU 403-108, SRPL.011/790, CLNF.049, JRTR.003, SAJAC.011

Rapid Assessment(s), OC stands (n=2): 1205EK02, 1520EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 67.2 (52.0) | 44-85 (44-60) | - |
| Herb cover (%) | 17.5 (4.5) | 0-52 (3-6) | 0-1 (0-0.5) |
| Shrub cover (%) | 56.9 (49.5) | 27-80 (44-55) | 0.5-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 1-5 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 93

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), NE (4,0), NW (2,0), SE (1,0), SW (1,0), Variable (2,0), W (1,1)

Macrotopography: lower 1/3 of slope (4,1), lower to middle 1/3 of slope (1,1), middle 1/3 of slope (1,0), upper 1/3 of slope (6,0)

Microtopography: undulating (9,1), Concave (1,0), Convex (1,0), flat (1,1)

Parent material: basalt (1,0), granitic (6,0), igneous (1,0), mixed alluvium (1,1), sandstone (1,1), sedimentary (1,0)

Soil texture: unknown (1,0), clay (1,0), clay loam (2,1), loam (1,0), sand (1,1), sandy loam (5,0)

| | Average | Range |
|------------------------------|----------------|-----------------------|
| Elevation (ft) | 2335 (1760) | 1240-3665 (1240-2280) |
| Slope (degrees) | 22 (17.5) | 3-45 (5-30) |
| Large rock cover (%) | 6.8 (5.0) | 0-30 (0-10) |
| Small rock cover (%) | 16.9 (22.5) | 0-48 (0-45) |
| Bare ground cover (%) | 19.8 (19.5) | 0-60 (7-32) |
| Litter cover (%) | 35.2 (50.0) | 0-70 (35-65) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-----------------------------------|------------|------------|------------|------------|
| Shrub | <i>Adenostoma fasciculatum</i> | 100 (100) | 20.2 (18) | 2 (12) | 39 (24) |
| | <i>Quercus berberidifolia</i> | 100 (100) | 24 (16) | 8 (12) | 55 (20) |
| | <i>Rhamnus ilicifolia</i> | 67 (100) | 3.9 (1) | 0.2 (1) | 15 (1) |
| | <i>Lonicera subspicata</i> | 58 (100) | 1.4 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 50 (100) | 1.5 (1) | 0.2 (1) | 5 (1) |
| | <i>Eriogonum fasciculatum</i> | 42 (50) | 3.2 (10) | 0.2 (10) | 10 (10) |
| | <i>Malosma laurina</i> | 33 (50) | 3.3 (1) | 1 (1) | 9 (1) |
| | <i>Rhus ovata</i> | 33 (-) | 1.9 (-) | 0.2 (-) | 6 (-) |
| | <i>Salvia mellifera</i> | 33 (-) | 2.5 (-) | 1 (-) | 5 (-) |
| | <i>Acmispon glaber</i> | 25 (100) | 4 (5) | 2 (5) | 5 (5) |
| | <i>Artemisia californica</i> | 25 (100) | 9.7 (2.1) | 0.2 (0.2) | 25 (4) |
| | <i>Eriophyllum confertiflorum</i> | 25 (100) | 2.4 (3.5) | 0.2 (1) | 6 (6) |
| | <i>Hesperoyucca whipplei</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Keckiella cordifolia</i> | 25 (50) | 0.5 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Lotus scoparius</i> | 25 (-) | 4 (-) | 2 (-) | 5 (-) |
| | <i>Toxicodendron diversilobum</i> | 25 (50) | 2.4 (0.2) | 0.2 (0.2) | 6 (0.2) |
| | <i>Yucca whipplei</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| Herb | <i>*Bromus madritensis</i> | 33 (-) | 9.3 (-) | 0.2 (-) | 15 (-) |
| | <i>*Bromus diandrus</i> | 25 (-) | 7.7 (-) | 2 (-) | 16 (-) |
| | <i>*Hirschfeldia incana</i> | 25 (-) | 5.3 (-) | 3 (-) | 10 (-) |
| | <i>Phacelia distans</i> | 25 (-) | 4.3 (-) | 2 (-) | 6 (-) |
| | <i>Unknown</i> | 25 (-) | 5.4 (-) | 0.2 (-) | 10 (-) |

***Rhus integrifolia* Alliance**
***Rhus integrifolia* Association**

Samples Used To Describe Association (n= 22)

Rapid Assessment(s), WR and WSD stands: 1602KH01, 0706TK05, 1520KH02, SDRP0284, SDRP0251, 1620JD01, SDRP0256, 0625JS01, 0623DH03, 0625DH04, 1625MB01, SDRP0096, 0706DH01, 1710JD10

Rapid Assessment(s), OC stands (n=8): 1003EK02, 1003EK03, 1056EK01, 1058EK01, 1062EK01, 1064EK02, 1099EK03, 1479EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 61.0 (49.8) | 30-95 (30-72) | - |
| Herb cover (%) | 5.0 (4.6) | 0.2-20 (1-20) | 0-5 (0-1) |
| Shrub cover (%) | 46.1 (46.1) | 10-75 (10-75) | 0-10 (0.5-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 0.9 (0.5) | 0.2-2 (0.2-1) | 2-10 (2-10) |

Local Environmental Description (All stands, OC stands)

Aspect: E (2,2), NE (8,4), NW (7,0), S (1,1), SE (1,1), SW (2,0), Variable (1,0)

Macrotopography: bottom (1,0), middle 1/3 of slope (1,1), upper 1/3 of slope to ridgetop (1,1), upper 1/3 of slope (4,3), lower to middle 1/3 of slope (5,1), entire slope (1,0), lower 1/3 of slope (3,0), lower to upper 1/3 of slope (5,2)

Microtopography: undulating (11,2), concave (5,2), convex (2,1), flat (4,3)

Parent material: conglomerate (1,0), igneous (1,0), metamorphic (1,0), sandstone (10,6), sandy alluvium (1,0)

Soil texture: loam (1,0), silty loam (1,0), sand (1,0), clay loam (4,3), clay (1,0), sandy loam (10,2)

| | Average | Range |
|------------------------------|----------------|-----------------|
| Elevation (ft) | 323 (525) | 13-732 (13-732) |
| Slope (degrees) | 21 (23.9) | 3-65 (5-65) |
| Large rock cover (%) | 0.7 (-) | 0-10 (-) |
| Small rock cover (%) | 6.6 (1.0) | 0-42 (0-7) |
| Bare ground cover (%) | 29.3 (47.0) | 0-74 (22-74) |
| Litter cover (%) | 43.3 (50.0) | 0-95 (25-75) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--------------------------------|------------|-------------|------------|------------|
| Shrub | <i>Rhus integrifolia</i> | 100 (100) | 41.5 (29.6) | 5 (5) | 85 (55) |
| | <i>Artemisia californica</i> | 82 (75) | 4.5 (3.8) | 0.2 (1) | 20 (8) |
| | <i>Mimulus aurantiacus</i> | 68 (75) | 1.3 (1.2) | 0.2 (0.2) | 8 (2) |
| | <i>Heteromeles arbutifolia</i> | 64 (63) | 4.8 (5.8) | 0.2 (2) | 20 (15) |
| | <i>Malosma laurina</i> | 55 (63) | 4.4 (3.2) | 0.2 (1) | 13 (7) |
| | <i>Salvia mellifera</i> | 50 (38) | 2.3 (0.8) | 0.2 (0.2) | 9 (2) |
| | <i>Opuntia littoralis</i> | 32 (38) | 1.7 (1.1) | 0.2 (0.2) | 5 (3) |
| | <i>Encelia californica</i> | 27 (-) | 0.8 (-) | 0.2 (-) | 3.2 (-) |
| | <i>Eriogonum fasciculatum</i> | 27 (38) | 3.2 (4.7) | 0.2 (1) | 8 (8) |
| | | | | | |
| Herb | <i>*Bromus madritensis</i> | 41 (38) | 3.4 (3.5) | 0.2 (0.2) | 10 (10) |
| | <i>Elymus condensatus</i> | 36 (50) | 0.8 (0.6) | 0.2 (0.2) | 2 (1) |
| | <i>Stipa lepida</i> | 27 (25) | 1.6 (2.6) | 0.2 (0.2) | 5 (5) |

Rhus integrifolia/Dudleya edulis-Carpobrotus spp. Provisional Alliance

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=3): 1001EK01, 1001EK02, 1001JD01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 17.3 (17.3) | 10-26 (10-26) | - |
| Herb cover (%) | 17.0 (17.0) | 9-25 (9-25) | 0-0.5 (0-0.5) |
| Shrub cover (%) | 4.5 (4.5) | 2-7 (2-7) | 0.5-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SE (1,1), W (2,2)

Macrotopography: lower 1/3 of slope (2,2), lower to upper 1/3 of slope (1,1)

Microtopography: undulating (3,3)

Parent material: sandstone (2,2)

Soil texture: sand (1,1), sandy loam (1,1)

| | Average | Range |
|-----------------------|-------------|---------------|
| Elevation (ft) | 76 (76) | 69-89 (69-89) |
| Slope (degrees) | 47 (47.0) | 26-75 (26-75) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 16.5 (16.5) | 3-30 (3-30) |
| Bare ground cover (%) | 76.0 (76.0) | 66-86 (66-86) |
| Litter cover (%) | 6.0 (6.0) | 2-10 (2-10) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>*Myoporum laetum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Isocoma menziesii</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Acmispon glaber</i> | 67 (67) | 1.5 (1.5) | 1 (1) | 2 (2) |
| | <i>Artemisia californica</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Eriogonum fasciculatum</i> | 67 (67) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>Eriogonum parvifolium</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Rhus integrifolia</i> | 67 (67) | 3 (3) | 2 (2) | 4 (4) |
| | <i>Atriplex canescens</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Atriplex semibaccata</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Baccharis pilularis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cylindropuntia prolifera</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Opuntia littoralis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Peritoma arborea var. arborea</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | | |
| | <i>*Bromus diandrus</i> | 100 (100) | 3.3 (3.3) | 1 (1) | 8 (8) |
| | <i>Dudleya edulis</i> | 100 (100) | 4.4 (4.4) | 0.2 (0.2) | 12 (12) |
| | <i>Atriplex californica</i> | 67 (67) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| | <i>*Carpobrotus edulis</i> | 67 (67) | 2 (2) | 1 (1) | 3 (3) |

| Species Name | Con | Ave | Min | Max |
|--------------------------------|---------|-----------|-----------|-----------|
| <i>*Malephora crocea</i> | 67 (67) | 1 (1) | 1 (1) | 1 (1) |
| <i>*Carpobrotus chilensis</i> | 33 (33) | 6 (6) | 6 (6) | 6 (6) |
| <i>Dichelostemma capitatum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Eriogonum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Festuca perennis</i> | 33 (33) | 3 (3) | 3 (3) | 3 (3) |
| <i>Heterotheca grandiflora</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Lupinus</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Medicago polymorpha</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Oenothera</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Oxalis pes-caprae</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Polypodium californicum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Salsola tragus</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Sonchus asper</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Salix lasiolepis* Alliance**

***Salix lasiolepis* (Alliance Only)**

Samples Used To Describe Association (n= 18)

Rapid Assessment(s), WR and WSD stands: 0707RB02, SDRP0339, 0709RB01, 0421TK03, SDRP0066, 1766KH01, 0624RB01, 0625DH03, 0709JS02, 1702FS01, 0310TK21, SDRP0095, 0504TK21, 0311TK25, 1730KH02, 0625DH05
Rapid Assessment(s), OC stands (n=2): 1028EK01, 1413EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-----------------|--------------|
| Total vegetation cover (%) | 55.6 (50.5) | 24-90 (45-56) | - |
| Herb cover (%) | 17.3 (5.5) | 0.2-45 (2-9) | 0-5 (0-0.5) |
| Shrub cover (%) | 27.7 (27.7) | 0.4-55 (0.4-55) | 0-10 (0.5-5) |
| Tall shrub / low tree cover (%) | 20.0 (20.0) | 20-20 (20-20) | 0.5-2 (1-2) |
| Overstory tree cover (%) | 7.9 (-) | 0.2-25 (-) | 2-20 (2-5) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (4,0), NW (4,0), S (1,1), SE (1,0), SW (8,1)

Macrotopography: bottom (18,2)

Microtopography: flat (9,0), undulating (1,0), concave (8,2)

Parent material: granitic (1,0), igneous (1,0), mixed alluvium (4,0), sandstone (2,1), sandy alluvium (4,0), silty alluvium (3,0)

Soil texture: clay loam (1,0), muck (2,0), sand (4,0), sandy loam (2,0), silt (1,0), silty loam (2,0), clay (1,0)

| | Average | Range |
|------------------------------|-------------|------------------|
| Elevation (ft) | 209 (154) | 0-1014 (121-187) |
| Slope (degrees) | 2 (1.5) | 0-3 (1-2) |
| Large rock cover (%) | 2.1 (-) | 0-30 (-) |
| Small rock cover (%) | 3.1 (2.5) | 0-14 (0-5) |
| Bare ground cover (%) | 21.4 (51.0) | 0-78 (47-55) |
| Litter cover (%) | 52.2 (45.0) | 0-88 (40-50) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 97

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------------|-----------|-------------|-----------|----------|
| Tree | <i>Salix lasiolepis</i> | 100 (100) | 34.4 (40.5) | 9 (35) | 88 (46) |
| | <i>Salix gooddingii</i> | 39 (50) | 3.6 (0.2) | 0.2 (0.2) | 8 (0.2) |
| | <i>Platanus racemosa</i> | 33 (-) | 1.4 (-) | 0.2 (-) | 5 (-) |
| | | | | | |
| Shrub | <i>Baccharis salicifolia</i> | 61 (50) | 8.4 (0.2) | 0.2 (0.2) | 30 (0.2) |
| | <i>Baccharis pilularis</i> | 33 (100) | 1 (0.6) | 0.2 (0.2) | 4 (1) |
| Herb | <i>Anemopsis californica</i> | 44 (-) | 2.5 (-) | 0.2 (-) | 10 (-) |
| | <i>Artemisia douglasiana</i> | 33 (50) | 1.4 (0.2) | 0.2 (0.2) | 6 (0.2) |
| | <i>Carex spissa</i> | 28 (-) | 8.5 (-) | 0.2 (-) | 37 (-) |
| | <i>Juncus acutus subsp. leopoldii</i> | 28 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |

***Salix lasiolepis* Alliance**

***Salix lasiolepis/Baccharis salicifolia* Association**

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=3): 1045EK01, 1057EK02, 1298MA01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|-----------------|-------------|
| Total vegetation cover (%) | 52.3 (52.3) | 45-62 (45-62) | - |
| Herb cover (%) | 15.0 (15.0) | 10-20 (10-20) | 0-1 (0-1) |
| Shrub cover (%) | 32.5 (32.5) | 10-55 (10-55) | 1-5 (1-5) |
| Tall shrub / low tree cover (%) | 17.5 (17.5) | 15-20 (15-20) | 1-2 (1-2) |
| Overstory tree cover (%) | 17.4 (17.4) | 0.2-45 (0.2-45) | 2-10 (2-10) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,1), SE (1,1), SW (1,1)

Macrotopography: bottom (3,3)

Microtopography: concave (2,2), flat (1,1)

Parent material: mixed alluvium (2,2), sandy alluvium (1,1)

Soil texture: sand (3,3)

| | Average | Range |
|------------------------------|-------------|-----------------|
| Elevation (ft) | 389 (389) | 66-961 (66-961) |
| Slope (degrees) | 1 (1.3) | 0-2 (0-2) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 8.7 (8.7) | 0-25 (0-25) |
| Bare ground cover (%) | 29.7 (29.7) | 7-52 (7-52) |
| Litter cover (%) | 37.7 (37.7) | 35-43 (35-43) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|-----------------------------------|-----------|-------------|-----------|-----------|
| Tree | <i>Salix lasiolepis</i> | 100 (100) | 34.7 (34.7) | 22 (22) | 52 (52) |
| | <i>Platanus racemosa</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Populus nigra</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Quercus agrifolia</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Washingtonia robusta</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | <i>Baccharis salicifolia</i> | 100 (100) | 13.3 (13.3) | 10 (10) | 15 (15) |
| | <i>Baccharis pilularis</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Artemisia californica</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Toxicodendron diversilobum</i> | 33 (33) | 2 (2) | 2 (2) | 2 (2) |
| Herb | <i>Artemisia douglasiana</i> | 67 (67) | 3 (3) | 2 (2) | 4 (4) |
| | * <i>Arundo donax</i> | 67 (67) | 8.5 (8.5) | 5 (5) | 12 (12) |
| | * <i>Melilotus indicus</i> | 67 (67) | 5.1 (5.1) | 0.2 (0.2) | 10 (10) |
| | <i>Typha</i> | 67 (67) | 2.6 (2.6) | 0.2 (0.2) | 5 (5) |
| | * <i>Brassica nigra</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Bromus diandrus</i> | 33 (33) | 7 (7) | 7 (7) | 7 (7) |
| | * <i>Centaurea melitensis</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| | * <i>Cortaderia jubata</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| | * <i>Cortaderia sellosana</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Urtica dioica</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Salvia apiana Alliance

Salvia apiana-*Artemisia californica* Association

Samples Used To Describe Association (n= 17)

Rapid Assessment(s), WR and WSD stands: SDG0030, SDG0009, 1633MB01, 1720FS01, SDG0042, 1686FS01, SDRP0097, SDRP0471, 0603RB01, 1073FS01, SDRP0140, SDG0033, 1102MB01, SDRP0136
Rapid Assessment(s), OC stands (n=3): 1161EK02, 1161MA02, 1172EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|-------------|
| Total vegetation cover (%) | 49.5 (47.7) | 22-75 (44-54) | - |
| Herb cover (%) | 20.5 (2.1) | 0.2-52 (0.2-4) | 0-1 (0-0.5) |
| Shrub cover (%) | 47.7 (47.7) | 44-54 (44-54) | 0-2 (0-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 2.6 (-) | 0.2-5 (-) | 2-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), SW (2,0), Variable (1,0), W (1,0), NE (5,0), NW (3,0), SE (4,2)

Macrotopography: lower to middle 1/3 of slope (3,0), ridge top (1,0), middle to upper 1/3 of slope (5,3), middle 1/3 of slope (1,0), lower 1/3 of slope (3,0), bottom (1,0), lower to upper 1/3 of slope (3,0)

Microtopography: convex (6,1), flat (5,0), undulating (4,1), concave (2,1)

Parent material: gabbro (2,0), granitic (10,3)

Soil texture: loam (2,0), sandy loam (5,0), silty loam (1,0), clay loam (4,3)

| | Average | Range |
|-----------------------|-------------|----------------------|
| Elevation (ft) | 1283 (3259) | 374-3346 (3189-3346) |
| Slope (degrees) | 20 (29.3) | 0-42 (20-42) |
| Large rock cover (%) | 4.5 (-) | 0-40 (-) |
| Small rock cover (%) | 8.5 (8.7) | 0-39 (0-16) |
| Bare ground cover (%) | 22.5 (26.0) | 0-70 (21-30) |
| Litter cover (%) | 22.2 (62.7) | 0-70 (58-70) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|-----------|------------|-----------|----------|
| Shrub | | | | |
| <i>Salvia apiana</i> | 100 (100) | 13.5 (29) | 4 (25) | 37 (37) |
| <i>Artemisia californica</i> | 94 (67) | 11.5 (4.5) | 0.2 (4) | 35 (5) |
| <i>Eriogonum fasciculatum</i> | 71 (33) | 2.9 (0.2) | 0.2 (0.2) | 15 (0.2) |
| <i>Malosma laurina</i> | 71 (33) | 3.8 (1) | 1 (1) | 12 (1) |
| <i>Mimulus aurantiacus</i> | 41 (67) | 4.5 (14) | 0.2 (5) | 23 (23) |
| <i>Acemispion glaber</i> | 29 (33) | 3 (14) | 0.2 (14) | 14 (14) |
| <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i> | 29 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Rhamnus crocea</i> | 29 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |
| Herb | | | | |
| * <i>Hirschfeldia incana</i> | 47 (33) | 2.2 (1) | 0.2 (1) | 10 (1) |
| * <i>Bromus madritensis</i> | 35 (-) | 2.2 (-) | 0.2 (-) | 5 (-) |
| * <i>Centaurea melitensis</i> | 29 (-) | 1.3 (-) | 0.2 (-) | 3.2 (-) |
| * <i>Erodium</i> | 29 (-) | 27.8 (-) | 2 (-) | 45 (-) |

Salvia leucophylla Alliance

Salvia leucophylla-*Artemisia californica* Association

Samples Used To Describe Association (n= 3)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=3): 1300MA01, 1341EK01, 1343EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 45.7 (45.7) | 36-60 (36-60) | - |
| Herb cover (%) | 15.7 (15.7) | 1-26 (1-26) | 0-1 (0-1) |
| Shrub cover (%) | 30.0 (30.0) | 9-41 (9-41) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: N (1,1), SE (2,2)

Macrotopography: middle 1/3 of slope (2,2), upper 1/3 of slope (1,1)

Microtopography: concave (1,1), flat (2,2)

Parent material: sandstone (2,2)

Soil texture: clay loam (3,3)

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 100

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 1419 (1419) | 1024-1703 (1024-1703) |
| Slope (degrees) | 35 (34.7) | 14-50 (14-50) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 0.7 (0.7) | 0-2 (0-2) |
| Bare ground cover (%) | 36.7 (36.7) | 10-58 (10-58) |
| Litter cover (%) | 60.0 (60.0) | 40-85 (40-85) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--------------------------------------|-----------|-----------|-----------|-----------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 9 (9) | 3 (3) | 19 (19) |
| <i>Malacothamnus fasciculatus</i> | 100 (100) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Salvia leucophylla</i> | 100 (100) | 21 (21) | 6 (6) | 35 (35) |
| <i>Malosma laurina</i> | 67 (67) | 1 (1) | 1 (1) | 1 (1) |
| <i>Eriogonum fasciculatum</i> | 33 (33) | 3 (3) | 3 (3) | 3 (3) |
| <i>Hesperoyucca whipplei</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Mirabilis laevis</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Nolina cismontana</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Salvia mellifera</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| <i>*Hirschfeldia incana</i> | 67 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stipa lepida</i> | 67 (67) | 8.6 (8.6) | 0.2 (0.2) | 17 (17) |
| <i>*Avena barbata</i> | 33 (33) | 25 (25) | 25 (25) | 25 (25) |
| <i>*Brachypodium distachyon</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Brassica nigra</i> | 33 (33) | 2 (2) | 2 (2) | 2 (2) |
| <i>Calochortus splendens</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Centaurea melitensis</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| <i>Elymus condensatus</i> | 33 (33) | 2 (2) | 2 (2) | 2 (2) |
| <i>Galium angustifolium</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Phacelia cicutaria</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Pseudognaphalium californicum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Solanum xanti</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stipa pulchra</i> | 33 (33) | 1 (1) | 1 (1) | 1 (1) |
| <i>Trifolium ciliolatum</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Uropappus lindleyi</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Salvia mellifera* Alliance**

***Salvia mellifera-Eriogonum fasciculatum* Association**

Samples Used To Describe Association (n= 30)

Rapid Assessment(s), WR and WSD stands: 1730KH01, SRPL.008/625, WINC.013, LAEL.014, SAGE.055, 0625RB01, ANZA.066, 1312KH01, 0707TK05, ANZA.064, SITT.006/51, JRTR.001, 0625RB02, WINC.015/difficult AA, SDG0099, 1126FS01, 1115PG01, SDRP0334, 0706TK01, SDG0093

Rapid Assessment(s), OC stands (n=10): 1099EK01, 1099EK04, 1099EK05, 1099MA01, 1101EK01, 1101EK02, 1128EK01, 1406EK01, 3.8jd05, ek3.8.4

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 101

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|----------------|-------------|
| Total vegetation cover (%) | 45.5 (48.2) | 21-75 (34-65) | - |
| Herb cover (%) | 7.4 (0.5) | 0.2-60 (0.2-1) | 0-1 (0-0.5) |
| Shrub cover (%) | 43.3 (48.1) | 20-65 (34-65) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-5 (-) |
| Overstory tree cover (%) | 0.6 (-) | 0.2-1 (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: E (2,2), NE (2,0), NW (3,0), SE (8,3), SW (10,5), Variable (5,0)

Macrotopography: middle to upper 1/3 of slope (7,5), upper 1/3 of slope to ridgetop (1,0), ridge top (2,0), middle 1/3 of slope (6,2), lower to upper 1/3 of slope (2,1), lower 1/3 of slope (2,0), bottom (1,0), upper 1/3 of slope (8,2)

Microtopography: undulating (8,2), concave (8,4), convex (8,2), flat (6,2)

Parent material: calcareous sandstone (1,0), granitic (9,0), mixed alluvium (1,1), sandstone (11,7), sedimentary (2,1), volcanic flow (1,0)

Soil texture: unknown (1,1), clay loam (4,1), loam (4,2), sand (3,0), sandy loam (14,6), silty loam (1,0)

| | Average | Range |
|-----------------------|-------------|------------------|
| Elevation (ft) | 989 (618) | 62-2721 (62-889) |
| Slope (degrees) | 20 (25.5) | 2-55 (2-55) |
| Large rock cover (%) | 2.2 (2.2) | 0-18 (0-18) |
| Small rock cover (%) | 14.1 (6.8) | 0-58 (0-35) |
| Bare ground cover (%) | 40.3 (45.6) | 0-73 (15-73) |
| Litter cover (%) | 33.7 (43.5) | 0-95 (25-65) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-------------------------------|-----------|-------------|-----------|----------|
| Shrub | | | | |
| <i>Salvia mellifera</i> | 100 (100) | 19.3 (22.4) | 5 (13) | 45 (35) |
| <i>Eriogonum fasciculatum</i> | 87 (100) | 8.3 (8.8) | 1 (2) | 30 (30) |
| <i>Artemisia californica</i> | 80 (100) | 5.6 (8.4) | 0.2 (0.2) | 20 (20) |
| <i>Rhus integrifolia</i> | 50 (100) | 5.4 (4.5) | 0.2 (0.2) | 20 (20) |
| <i>Malosma laurina</i> | 47 (60) | 3.5 (3.1) | 0.2 (0.2) | 10 (7) |
| <i>Opuntia littoralis</i> | 33 (40) | 0.3 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Acmispon glaber</i> | 30 (50) | 0.6 (0.7) | 0.2 (0.2) | 2 (2) |
| <i>Lotus scoparius</i> | 30 (-) | 0.6 (-) | 0.2 (-) | 2 (-) |
| <i>Mimulus aurantiacus</i> | 27 (60) | 2 (1.4) | 0.2 (0.2) | 5 (5) |
| Herb | | | | |
| * <i>Bromus madritensis</i> | 50 (30) | 3.9 (0.2) | 0.2 (0.2) | 22 (0.2) |

Salvia mellifera Alliance

Salvia mellifera-Lotus scoparius Association

Samples Used To Describe Association (n= 11)

Rapid Assessment(s), WR and WSD stands: LABO.008, SATI.003, SAGE.027, SAGE.035, SCWP.001, SITT.018

Rapid Assessment(s), OC stands (n=5): 1197EK04, 1380EK02, 1463EK05, 1499EK02, 1499KH02

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 102

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-----------------|---------------|
| Total vegetation cover (%) | 37.6 (43.6) | 13-60 (37-48) | - |
| Herb cover (%) | 9.7 (4.2) | 0.2-45 (0.2-15) | 0-1 (0-0.5) |
| Shrub cover (%) | 29.8 (39.6) | 12-46 (30-46) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-2 (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: E (1,1), W (1,1), SW (3,2), SE (2,1), NE (3,0), NW (1,0)

Macrotopography: middle 1/3 of slope (3,3), ridge top (2,0), upper 1/3 of slope (5,1), upper 1/3 of slope to ridgetop (1,1)

Microtopography: Undulating (4,1), convex (5,2), flat (2,2)

Parent material: granitic (2,0), mixed igneous (1,0), sandstone (3,3), sedimentary (2,0)

Soil texture: clay loam (2,1), sandy loam (9,4)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 1752 (1348) | 951-2338 (951-1923) |
| Slope (degrees) | 21 (19.4) | 3-30 (3-30) |
| Large rock cover (%) | 2.5 (-) | 0-25 (-) |
| Small rock cover (%) | 12.1 (0.8) | 0-78 (0-3) |
| Bare ground cover (%) | 45.5 (61.6) | 3-87 (42-87) |
| Litter cover (%) | 36.1 (35.4) | 0-90 (9-55) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|-----------|-------------|-----------|-----------|
| Shrub | | | | |
| <i>Acmispon glaber</i> | 100 (100) | 6.5 (10.4) | 0.2 (3) | 24 (24) |
| <i>Lotus scoparius</i> | 100 (-) | 6.5 (-) | 0.2 (-) | 24 (-) |
| <i>Salvia mellifera</i> | 100 (100) | 14.1 (15.6) | 4 (8) | 25 (21) |
| <i>Artemisia californica</i> | 82 (100) | 3 (2.4) | 1 (1) | 5 (4) |
| <i>Hesperoyucca whipplei</i> | 45 (40) | 0.9 (1.5) | 0.2 (1) | 2 (2) |
| <i>Malacothamnus fasciculatus</i> | 45 (80) | 6.4 (7.8) | 0.2 (0.2) | 15 (15) |
| <i>Malosma laurina</i> | 45 (80) | 2.1 (2.6) | 0.2 (0.2) | 4 (4) |
| <i>Yucca whipplei</i> | 45 (-) | 0.9 (-) | 0.2 (-) | 2 (-) |
| <i>Adenostoma fasciculatum</i> | 36 (40) | 3 (2) | 1 (1) | 5 (3) |
| <i>Eriogonum fasciculatum</i> | 36 (40) | 1.9 (1.5) | 0.2 (1) | 5 (2) |
| <i>Rhus ovata</i> | 36 (20) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Nolina cismontana</i> | 27 (60) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Opuntia littoralis</i> | 27 (60) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Centaurea melitensis</i> | 64 (100) | 0.8 (0.9) | 0.2 (0.2) | 3 (3) |
| * <i>Hirschfeldia incana</i> | 45 (40) | 1.3 (0.2) | 0.2 (0.2) | 4 (0.2) |
| * <i>Brassica nigra</i> | 27 (60) | 2.4 (2.4) | 0.2 (0.2) | 6 (6) |
| * <i>Bromus madritensis</i> | 27 (20) | 4 (2) | 1 (2) | 9 (2) |

***Salvia mellifera* Alliance**

***Salvia mellifera*-*Malosma laurina* Association**

Samples Used To Describe Association (n= 29)

Rapid Assessment(s), WR and WSD stands: 0311TK24, 1732JD01, SDRP0329, SDRP0359, SDRP0270, 1339FS01, 1278LS01, 1485MM01, SDRP0235, 1493FS01, SDG0117, 0507RB01, 0311AK22, 0507MG01, 0507DH01, SDRP0061, 0623RB03, 0708JS01, 0624RB03, 1700FS01, 1439FS01, 0708AK01, 0624DH02, SDG0081, SDRP0059

Rapid Assessment(s), OC stands (n=4): 1032EK01, 1088EK04, 1343EK04, 1413EK04

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|----------------|-------------|
| Total vegetation cover (%) | 48.0 (39.3) | 14-93 (37-42) | - |
| Herb cover (%) | 7.2 (2.8) | 0.2-35 (0.2-8) | 0-2 (0-0.5) |
| Shrub cover (%) | 37.0 (37.0) | 33-42 (33-42) | 0-5 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 2.0 (-) | 2-2 (-) | 5-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Variable (4,0), N (1,1), NE (2,0), NW (5,0), S (1,1), SE (8,2), SW (8,0)

Macrotopography: middle 1/3 of slope (4,2), upper 1/3 of slope (4,2), upper 1/3 of slope to ridgetop (3,0), middle 1/3 of slope to ridgetop (1,0), lower to middle 1/3 of slope (2,0), lower 1/3 of slope to ridgetop (1,0), lower 1/3 of slope (2,0), bottom to mid 1/3 of slope (1,0), middle to upper 1/3 of slope (7,0), lower to upper 1/3 of slope (4,0)

Microtopography: concave (2,0), convex (11,1), flat (6,3), undulating (10,0)

Parent material: conglomerate (3,0), general volcanic extrusives (1,0), granitic (6,0), metamorphic (2,0), mixed sedimentary (2,0), sandstone (5,4), sedimentary (3,0), volcanic flow (1,0)

Soil texture: clay (2,0), clay loam (4,1), loam (3,0), sand (2,0), sandy loam (16,1)

| | Average | Range |
|------------------------------|-------------|---------------------|
| Elevation (ft) | 684 (932) | 112-1588 (466-1588) |
| Slope (degrees) | 20 (22.0) | 3-38 (12-30) |
| Large rock cover (%) | 2.1 (-) | 0-15 (-) |
| Small rock cover (%) | 18.9 (12.8) | 0-58 (0-37) |
| Bare ground cover (%) | 30.8 (48.8) | 0-68 (31-63) |
| Litter cover (%) | 28.8 (36.3) | 0-87 (30-45) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--------------------------------------|-----------|-------------|-----------|----------|
| Shrub | <i>Salvia mellifera</i> | 100 (100) | 26.1 (26.8) | 5 (17) | 60 (37) |
| | <i>Malosma laurina</i> | 97 (100) | 6.2 (2.3) | 0.2 (0.2) | 30 (4) |
| | <i>Artemisia californica</i> | 52 (25) | 3.1 (4) | 0.2 (4) | 8 (4) |
| | <i>Eriogonum fasciculatum</i> | 48 (25) | 1.5 (1) | 0.2 (1) | 4 (1) |
| | <i>Mimulus aurantiacus</i> | 45 (25) | 10.2 (0.2) | 0.2 (0.2) | 40 (0.2) |
| | <i>Rhus integrifolia</i> | 41 (50) | 4.5 (5) | 0.2 (3) | 20 (7) |
| | <i>Acmispon glaber</i> | 28 (25) | 4.1 (1) | 0.2 (1) | 11 (1) |
| | <i>Heteromeles arbutifolia</i> | 28 (50) | 1.4 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>Malacothamnus fasciculatus</i> | 28 (50) | 0.5 (1.1) | 0.2 (0.2) | 2 (2) |
| | | | | | |
| Herb | <i>*Bromus madritensis</i> | 52 (25) | 5.8 (3) | 0.2 (3) | 35 (3) |
| | <i>*Centaurea melitensis</i> | 38 (50) | 1.4 (0.2) | 0.2 (0.2) | 11 (0.2) |
| | <i>Stipa lepida</i> | 34 (75) | 1.7 (2.3) | 0.2 (2) | 5 (3) |
| | <i>Pseudognaphalium californicum</i> | 31 (-) | 2.6 (-) | 0.2 (-) | 15 (-) |

***Sambucus nigra* Alliance**
***Sambucus nigra* Association**

Samples Used To Describe Association (n= 2)

Rapid Assessment(s), WR and WSD stands: None
Rapid Assessment(s), OC stands (n=2): 1463KH01, 1463KH02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 42.5 (42.5) | 35-50 (35-50) | - |
| Herb cover (%) | 24.0 (24.0) | 10-38 (10-38) | 0.5-1 (0.5-1) |
| Shrub cover (%) | 18.5 (18.5) | 12-25 (12-25) | 2-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,1), W (1,1)
Macrotopography: middle to upper 1/3 of slope (1,1), middle 1/3 of slope to ridgetop (1,1)
Microtopography: convex (2,2)
Parent material:
Soil texture: clay loam (1,1), loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 1060 (1060) | 1033-1086 (1033-1086) |
| Slope (degrees) | 15 (15.0) | 15-15 (15-15) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 1.5 (1.5) | 1-2 (1-2) |
| Bare ground cover (%) | 61.5 (61.5) | 42-81 (42-81) |
| Litter cover (%) | 35.0 (35.0) | 15-55 (15-55) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---|-----------|-------------|-----------|-----------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 100 (100) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| <i>Baccharis pilularis</i> | 100 (100) | 5 (5) | 3 (3) | 7 (7) |
| <i>Heteromeles arbutifolia</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Sambucus nigra</i> subsp. <i>caerulea</i> | 100 (100) | 12 (12) | 6 (6) | 18 (18) |
| <i>Isocoma menziesii</i> var. <i>vernonioides</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Lonicera subspicata</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>Malosma laurina</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>Marah</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Mimulus aurantiacus</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Opuntia littoralis</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>Salvia apiana</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Salvia mellifera</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Toxicodendron diversilobum</i> | 50 (50) | 1 (1) | 1 (1) | 1 (1) |
| Herb | | | | |
| * <i>Brassica nigra</i> | 100 (100) | 11.5 (11.5) | 6 (6) | 17 (17) |
| * <i>Carduus pycnocephalus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Elymus condensatus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 105

| | | | | |
|--------------------------------------|-----------|-----------|-----------|-----------|
| <i>Galium aparine</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Marrubium vulgare</i> | 100 (100) | 0.6 (0.6) | 0.2 (0.2) | 1 (1) |
| <i>Pseudognaphalium californicum</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Avena barbata</i> | 50 (50) | 3 (3) | 3 (3) | 3 (3) |
| <i>Bowlesia incana</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus diandrus</i> | 50 (50) | 5 (5) | 5 (5) | 5 (5) |
| * <i>Bromus hordeaceus</i> | 50 (50) | 5 (5) | 5 (5) | 5 (5) |
| * <i>Bromus madritensis</i> | 50 (50) | 7 (7) | 7 (7) | 7 (7) |
| * <i>Centaurea melitensis</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| <i>Chenopodium californicum</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Cynara cardunculus</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Eucrypta chrysanthemifolia</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Stachys ajugoides</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| <i>Stipa pulchra</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Urtica dioica</i> | 50 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Toxicodendron diversilobum* Alliance**

***Toxicodendron diversilobum*-*Artemisia californica*/*Leymus condensatus* Association**

Samples Used To Describe Association (n= 5)

Rapid Assessment(s), WR and WSD stands: 1737JD07, 1739JD01, 1737JD05

Rapid Assessment(s), OC stands (n=2): 1218KH01, 1494EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 71.0 (60.0) | 60-85 (60-60) | - |
| Herb cover (%) | 26.8 (46.0) | 6-46 (46-46) | 0.5-2 (0.5-1) |
| Shrub cover (%) | 37.5 (37.5) | 15-60 (15-60) | 0-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 1.0 (1.0) | 1-1 (1-1) | 5-10 (5-10) |

Local Environmental Description (All stands, OC stands)

Aspect: N (2,2), NE (1,0), NW (2,0)

Macrotopography: lower to middle 1/3 of slope (1,0), lower to upper 1/3 of slope (2,0), middle to upper 1/3 of slope (2,2)

Microtopography: convex (1,0), flat (1,0), undulating (3,2)

Parent material: sandstone (1,1)

Soil texture: clay loam (2,1), sandy loam (2,0)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 480 (880) | 197-1516 (243-1516) |
| Slope (degrees) | 22 (25.5) | 12-30 (25-26) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 7.2 (0.5) | 0-25 (0-1) |
| Bare ground cover (%) | 32.0 (47.0) | 13-57 (37-57) |
| Litter cover (%) | 59.0 (50.0) | 40-85 (40-60) |



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 106

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|---------------------------------------|-----------|-------------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 40 (50) | 1.1 (1) | 0.2 (1) | 2 (1) |
| Shrub | | | | | |
| | <i>Toxicodendron diversilobum</i> | 100 (100) | 38.2 (24.5) | 7 (7) | 55 (42) |
| | <i>Rhus integrifolia</i> | 80 (100) | 8.8 (7) | 2 (2) | 12 (12) |
| | <i>Sambucus nigra subsp. caerulea</i> | 80 (100) | 0.4 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>Artemisia californica</i> | 60 (100) | 3.7 (1.5) | 1 (1) | 8 (2) |
| | <i>Baccharis pilularis</i> | 40 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Heteromeles arbutifolia</i> | 40 (100) | 5.5 (5.5) | 5 (5) | 6 (6) |
| Herb | | | | | |
| | <i>Elymus condensatus</i> | 100 (100) | 3.1 (7.1) | 0.2 (0.2) | 14 (14) |
| | <i>Artemisia palmeri</i> | 60 (-) | 19 (-) | 5 (-) | 30 (-) |
| | * <i>Brassica nigra</i> | 60 (50) | 1.4 (3) | 0.2 (3) | 3 (3) |

VEGETATION DOMINATED BY HERBS

Avena (barbata, fatua) Alliance

Avena barbata-Avena fatua Semi-Natural Stands

Samples Used To Describe Association (n= 15)

Rapid Assessment(s), WR and WSD stands: 0519DH01, 1737JD08, 1587LS01, SDRP0237, 1560LS01, 1361LS01, 1697LS01, 1339EB02, 0519RB04, 1361KH03, 0519TK02, 1545LS01, 1596KH01

Rapid Assessment(s), OC stands (n=2): 1041EK02, 1187EK04

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|-------------|
| Total vegetation cover (%) | 48.5 (45.5) | 29-65 (35-56) | - |
| Herb cover (%) | 46.2 (45.5) | 25-65 (35-56) | 0-2 (0-0.5) |
| Shrub cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0-2 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Variable (2,1), SW (5,0), NE (2,0), NW (2,0), SE (3,0), W (1,1)

Macrotopography: bottom (1,0), bottom to lower 1/3 of slope (1,0), bottom to mid 1/3 of slope (3,0), bottom to upper 1/3 of slope (1,0), entire slope (1,0), lower 1/3 of slope (1,0), lower to upper 1/3 of slope (2,0), middle 1/3 of slope (1,1), middle 1/3 of slope to ridgetop (1,1), upper 1/3 of slope to ridgetop (2,0)

Microtopography: undulating (4,0), flat (3,1), concave (2,0), convex (6,1)

Parent material: granitic (2,0), metamorphic (2,0), sandstone (2,2), sandy alluvium (1,0)

Soil texture: clay loam (3,1), loam (2,0), sandy loam (9,0), clay (1,1)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 672 (822) | 115-1289 (607-1037) |
| Slope (degrees) | 17 (16.0) | 0-42 (16-16) |
| Large rock cover (%) | 0.7 (-) | 0-5 (-) |
| Small rock cover (%) | 7.3 (-) | 0-35 (-) |
| Bare ground cover (%) | 35.7 (26.0) | 5-72 (23-29) |
| Litter cover (%) | 41.0 (71.5) | 0-92 (68-75) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|--|----------|-----------|-----------|---------|
| Shrub | | | | |
| <i>Artemisia californica</i> | 60 (50) | 0.8 (0.2) | 0.2 (0.2) | 3 (0.2) |
| <i>Eriogonum fasciculatum</i> var. <i>fasciculatum</i> | 33 (-) | 1.1 (-) | 0.2 (-) | 3 (-) |
| <i>Malosma laurina</i> | 33 (-) | 2.2 (-) | 0.2 (-) | 6 (-) |
| Herb | | | | |
| * <i>Avena barbata</i> | 87 (100) | 22.4 (21) | 6 (16) | 45 (26) |
| * <i>Bromus diandrus</i> | 60 (100) | 5.3 (12) | 0.2 (1) | 23 (23) |
| * <i>Centaurea melitensis</i> | 47 (-) | 6.2 (-) | 0.2 (-) | 20 (-) |
| * <i>Brassica nigra</i> | 40 (100) | 3.6 (1.1) | 0.2 (0.2) | 17 (2) |
| * <i>Avena fatua</i> | 33 (-) | 24.6 (-) | 3 (-) | 40 (-) |
| * <i>Bromus rubens</i> | 33 (-) | 10.4 (-) | 8 (-) | 15 (-) |
| * <i>Hirschfeldia incana</i> | 33 (-) | 1.3 (-) | 0.2 (-) | 5 (-) |
| * <i>Salsola tragus</i> | 27 (-) | 0.9 (-) | 0.2 (-) | 2 (-) |
| <i>Stipa pulchra</i> | 27 (50) | 1.1 (1) | 0.2 (1) | 2 (1) |

Brassica nigra Alliance

Brassica nigra Semi-Natural Stands

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: 1703JD01, SDG0118

Rapid Assessment(s), OC stands (n=2): 1010EK02, 1499KH01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|---------------|-------------------|
| Total vegetation cover (%) | 43.3 (46.0) | 15-66 (45-47) | - |
| Herb cover (%) | 42.0 (44.5) | 13-66 (44-45) | 0.5-2 (0.5-1) |
| Shrub cover (%) | 5.0 (5.0) | 5-5 (5-5) | 0-1 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1), NW (1,0), SW (1,0), W (1,1)

Macrotopography: middle to upper 1/3 of slope (1,1), lower to middle 1/3 of slope (1,0), middle 1/3 of slope (1,1)

Microtopography: convex (1,0), undulating (2,2)

Parent material: basalt (1,0), sandstone (1,1)

Soil texture: clay loam (3,2), sandy loam (1,0)

| | Average | Range |
|------------------------------|----------------|---------------------------------|
| Elevation (ft) | 600 (886) | 128-1119 (653-1119) |
| Slope (degrees) | 19 (14.0) | 3-29 (3-25) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 1.3 (-) | 0-5 (-) |
| Bare ground cover (%) | 32.8 (23.0) | 23-52 (23-23) |
| Litter cover (%) | 52.6 (75.0) | 0.200000002980232-75 (75-75) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|------------------------------------|------------|-------------|------------|------------|
| Shrub | <i>Artemisia californica</i> | 75 (50) | 0.8 (2) | 0.2 (2) | 2 (2) |
| | <i>Opuntia littoralis</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Adolphia californica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Astragalus trichopodus</i> var. | | | | |
| | <i>lonchus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Baccharis pilularis</i> | 25 (50) | 3 (3) | 3 (3) | 3 (3) |
| | <i>Encelia californica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Malacothamnus fasciculatus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Malosma laurina</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | <i>Mirabilis laevis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Salvia mellifera</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | | | | | |
| Herb | <i>*Brassica nigra</i> | 100 (100) | 28.8 (26.5) | 7 (25) | 55 (28) |
| | <i>*Bromus diandrus</i> | 75 (100) | 7.1 (10.5) | 0.2 (10) | 11 (11) |
| | <i>*Carduus pycnocephalus</i> | 75 (100) | 3.5 (0.2) | 0.2 (0.2) | 10 (0.2) |
| | <i>*Avena barbata</i> | 50 (100) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| | <i>*Bromus madritensis</i> | 50 (50) | 2.6 (5) | 0.2 (5) | 5 (5) |

| Species Name | Con | Ave | Min | Max |
|--------------------------------------|----------|-----------|-----------|-----------|
| <i>*Cynara cardunculus</i> | 50 (100) | 3 (3) | 1 (1) | 5 (5) |
| <i>Elymus condensatus</i> | 50 (50) | 0.6 (0.2) | 0.2 (0.2) | 1 (0.2) |
| <i>Antirrhinum nuttallianum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Bromus hordeaceus</i> | 25 (50) | 7 (7) | 7 (7) | 7 (7) |
| <i>Calystegia macrostegia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Centaurea melitensis</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| <i>*Cirsium vulgare</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Conium maculatum</i> | 25 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>Cryptantha</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>*Erodium cicutarium</i> | 25 (50) | 1 (1) | 1 (1) | 1 (1) |
| <i>*Erodium moschatum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Eucrypta chrysanthemifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Festuca myuros</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Foeniculum vulgare</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Galium aparine</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Heliotropium curassavicum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Malva</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Melilotus indicus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Phacelia distans</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Pseudognaphalium californicum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Raphanus sativus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>*Salsola tragus</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Silybum marianum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>*Sonchus asper</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Bromus diandrus* Alliance**

***Bromus diandrus* Semi-Natural Stands**

Samples Used To Describe Association (n= 20)

Rapid Assessment(s), WR and WSD stands: 0421DH02, SDRP0068, 1022DH06, 0423DH02, LAVI.026/BPU 22363-45

Rapid Assessment(s), OC stands (n=15): 1003EK06, 1047EK05, 1066EK04, 1066EK05, 1076MA01, 1079EK01, 1093EK02, 1110EK03, 1113EK01, 1128EK05, 1137EK02, 1197EK05, 1229EK03, 1243EK01, 1289EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 58.4 (57.7) | 35-90 (35-90) | - |
| Herb cover (%) | 56.3 (55.2) | 35-90 (35-90) | 0-1 (0-1) |
| Shrub cover (%) | 1.2 (1.3) | 0-7 (0.2-7) | 0-5 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 110

Local Environmental Description (All stands, OC stands)

Aspect: S (4,4), SW (2,1), Variable (1,1), SE (5,4), NE (2,0), NW (1,1), Flat (2,1), E (2,2), W (1,1)

Macrotopography: middle to upper 1/3 of slope (2,2), upper 1/3 of slope to ridgetop (3,3), upper 1/3 of slope (2,2), lower to middle 1/3 of slope (1,1), bottom (2,0), middle 1/3 of slope (2,2), lower 1/3 of slope (8,5)

Microtopography: concave (1,0), convex (1,1), flat (11,9), undulating (7,5)

Parent material: granitic (4,0), sandstone (13,13), sedimentary (1,1), silty alluvium (1,0)

Soil texture: sandy loam (6,3), silty loam (1,0), loam (2,2), clay loam (6,5), clay (3,3)

| | Average | Range |
|------------------------------|----------------|------------------|
| Elevation (ft) | 710 (662) | 0-1427 (79-1427) |
| Slope (degrees) | 7 (8.1) | 0-22 (0-22) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 0.7 (-) | 0-10 (-) |
| Bare ground cover (%) | 21.0 (21.3) | 0-52 (2-52) |
| Litter cover (%) | 70.8 (76.3) | 0-95 (45-95) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|------------------------------|------------|------------|------------|------------|
| Shrub | | | | |
| <i>Baccharis pilularis</i> | 25 (33) | 1.8 (1.8) | 0.2 (0.2) | 6 (6) |
| <i>Isocoma menziesii</i> | 25 (20) | 0.7 (0.5) | 0.2 (0.2) | 1 (1) |
| <i>Opuntia littoralis</i> | 25 (33) | 0.4 (0.4) | 0.2 (0.2) | 1 (1) |
| Herb | | | | |
| * <i>Bromus diandrus</i> | 100 (100) | 45 (45) | 17 (20) | 83 (83) |
| * <i>Brassica nigra</i> | 60 (80) | 6.5 (6.5) | 1 (1) | 18 (18) |
| * <i>Avena barbata</i> | 40 (40) | 4.7 (6.2) | 0.2 (1) | 10 (10) |
| <i>Croton setiger</i> | 30 (27) | 2.9 (0.4) | 0.2 (0.2) | 10 (1) |
| <i>Eremocarpus setigerus</i> | 30 (-) | 2.9 (-) | 0.2 (-) | 10 (-) |
| * <i>Hirschfeldia incana</i> | 30 (13) | 1.7 (2) | 0.2 (1) | 5 (3) |
| * <i>Bromus hordeaceus</i> | 25 (13) | 2.6 (3.6) | 0.2 (0.2) | 7 (7) |
| * <i>Lactuca serriola</i> | 25 (20) | 0.6 (0.2) | 0.2 (0.2) | 2 (0.2) |

***Cynara cardunculus* Provisional Alliance**

***Cynara cardunculus* Provisional Semi-Natural Stands**

Samples Used To Describe Association (n= 6)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=6): 1015EK01, 1015EK03, 1030EK01, 1088EK03, 1113EK02, 1513EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|-------------------|-------------------|
| Total vegetation cover (%) | 44.8 (44.8) | 24-60 (24-60) | - |
| Herb cover (%) | 44.8 (44.8) | 24-60 (24-60) | 0-1 (0-1) |
| Shrub cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0.5-2 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 111

Local Environmental Description (All stands, OC stands)

Aspect: NE (1,1), NW (2,2), SW (1,1), Variable (2,2)

Macrotopography: middle to upper 1/3 of slope (2,2), upper 1/3 of slope (1,1), lower 1/3 of slope to ridge top (1,1), middle 1/3 of slope (2,2)

Microtopography: concave (1,1), flat (1,1), undulating (4,4)

Parent material: sandstone (5,5)

Soil texture: clay (1,1), clay loam (4,4)

| | Average | Range |
|-----------------------|-------------|---------------------|
| Elevation (ft) | 675 (675) | 253-1014 (253-1014) |
| Slope (degrees) | 15 (15.0) | 4-24 (4-24) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 24.0 (24.0) | 9-58 (9-58) |
| Litter cover (%) | 74.2 (74.2) | 40-90 (40-90) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|------------------------------|-----------|-------------|-----------|-----------|
| Shrub | | | | |
| <i>Rhus integrifolia</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | |
| * <i>Cynara cardunculus</i> | 100 (100) | 31.7 (31.7) | 18 (18) | 43 (43) |
| * <i>Avena barbata</i> | 83 (83) | 2.5 (2.5) | 0.2 (0.2) | 6 (6) |
| * <i>Bromus diandrus</i> | 83 (83) | 4.8 (4.8) | 0.2 (0.2) | 10 (10) |
| * <i>Bromus hordeaceus</i> | 83 (83) | 6 (6) | 0.2 (0.2) | 14 (14) |
| * <i>Brassica nigra</i> | 67 (67) | 1.8 (1.8) | 1 (1) | 2 (2) |
| * <i>Bromus madritensis</i> | 33 (33) | 1.6 (1.6) | 0.2 (0.2) | 3 (3) |
| * <i>Hirschfeldia incana</i> | 33 (33) | 2 (2) | 2 (2) | 2 (2) |
| * <i>Sonchus asper</i> | 33 (33) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Festuca perennis* Alliance**

***Festuca perennis* Semi-Natural Stands**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: 1563MM01, 0505TK22, 1019RB01

Rapid Assessment(s), OC stands (n=1): 1092EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|-------------|
| Total vegetation cover (%) | 59.0 (36.0) | 36-88 (36-36) | - |
| Herb cover (%) | 58.5 (34.0) | 34-88 (34-34) | 0-1 (0-0.5) |
| Shrub cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0-5 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 112

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,0), NE (1,1), NW (1,0), SE (1,0)

Macrotopography: bottom to lower 1/3 of slope (1,0), upper 1/3 of slope (1,1), bottom (2,0)

Microtopography: concave (2,1), flat (2,0)

Parent material: clayey alluvium (1,0), sandstone (1,1), sandy alluvium (1,0)

Soil texture: clay (1,1), clay loam (1,0), loam (1,0), silt (1,0)

| | Average | Range |
|------------------------------|-------------|-----------------|
| Elevation (ft) | 527 (696) | 0-879 (696-696) |
| Slope (degrees) | 7 (26.0) | 0-26 (26-26) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 2.8 (-) | 0-10 (-) |
| Bare ground cover (%) | 21.5 (10.0) | 3-63 (10-10) |
| Litter cover (%) | 73.5 (88.0) | 25-95 (88-88) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--|----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>Quercus agrifolia</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Shrub | | | | | |
| | <i>Artemisia californica</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Baccharis salicifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Baccharis sarothroides</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Grindelia camporum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Isocoma menziesii</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Mimulus aurantiacus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| Herb | | | | | |
| | <i>*Lolium multiflorum</i> | 75 (-) | 51 (-) | 28 (-) | 85 (-) |
| | <i>*Rumex crispus</i> | 75 (-) | 1.4 (-) | 0.2 (-) | 3 (-) |
| | <i>*Bromus hordeaceus</i> | 50 (100) | 0.6 (1) | 0.2 (1) | 1 (1) |
| | <i>Ambrosia psilostachya</i> | 25 (-) | 8 (-) | 8 (-) | 8 (-) |
| | <i>Atriplex prostrata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>*Avena barbata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>*Bromus diandrus</i> | 25 (100) | 7 (7) | 7 (7) | 7 (7) |
| | <i>*Carduus pycnocephalus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cuscuta</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>*Cynara cardunculus</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Daucus pusillus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Dichelostemma capitatum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Distichlis spicata</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>*Erodium botrys</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>*Festuca perennis</i> | 25 (100) | 22 (22) | 22 (22) | 22 (22) |
| | <i>*Foeniculum vulgare</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Frankenia salina</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| | <i>Heliotropium curassavicum</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | <i>*Helminthotheca echioides</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>*Hirschfeldia incana</i> | 25 (-) | 8 (-) | 8 (-) | 8 (-) |
| | <i>*Hordeum murinum subsp. glaucum</i> | 25 (-) | 15 (-) | 15 (-) | 15 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 113

| Species Name | Con | Ave | Min | Max |
|---------------------------------------|----------|-----------|-----------|-----------|
| <i>Juncus acutus subsp. leopoldii</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Juncus dubius</i> | 25 (-) | 12 (-) | 12 (-) | 12 (-) |
| * <i>Lepidium draba</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Lupinus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Malvella leprosa</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Melilotus indicus</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| <i>Ranunculus californicus</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| * <i>Raphanus sativus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Salicornia pacifica</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| <i>Sanicula bipinnatifida</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Sisyrinchium bellum</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Stipa pulchra</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| <i>Xanthium strumarium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

Frankenia salina Alliance

Frankenia salina-Distichlis spicata Association

Samples Used To Describe Association (n= 11)

Rapid Assessment(s), WR and WSD stands: 1020AK06, 0707AK01, 1020AK05, 1020RB03, 1020TK01, 1020TK03, 1023TK06, 1021RB03, 1022DH02, 1022AK04

Rapid Assessment(s), OC stands (n=1): 1471EK03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 62.9 (68.0) | 21-86 (68-68) | - |
| Herb cover (%) | 62.8 (68.0) | 21-86 (68-68) | 0-1 (0-0.5) |
| Shrub cover (%) | - (-) | - (-) | 0-5 (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 1.0 (-) | 1-1 (-) | 10-15 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (7,0), NW (2,0), S (1,1), Variable (1,0)

Macrotopography: lower 1/3 of slope (2,0), bottom (9,1)

Microtopography: flat (7,1), undulating (4,0)

Parent material: clayey alluvium (4,0), sandy alluvium (2,0), silty alluvium (4,0)

Soil texture: sand (1,1), sandy loam (1,0), clay loam (3,0), clay (4,0), silt (2,0)

| | Average | Range |
|------------------------------|-------------|---------------|
| Elevation (ft) | 1 (7) | 0-7 (7-7) |
| Slope (degrees) | 1 (0.0) | 0-8 (0-0) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 0.0 (-) | 0-0.4 (-) |
| Bare ground cover (%) | 13.6 (17.0) | 0-78 (17-17) |
| Litter cover (%) | 83.6 (80.0) | 20-97 (80-80) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|----------------------------------|-----------|-----------|-----------|----------|
| Shrub | | | | | |
| | <i>Cuscuta salina</i> | 27 (100) | 7.1 (0.2) | 0.2 (0.2) | 20 (0.2) |
| Herb | | | | | |
| | <i>Frankenia salina</i> | 100 (100) | 41.9 (38) | 17 (38) | 82 (38) |
| | <i>Distichlis spicata</i> | 82 (100) | 5.2 (0.2) | 0.2 (0.2) | 18 (0.2) |
| | <i>Salicornia pacifica</i> | 82 (100) | 5.5 (22) | 0.2 (22) | 22 (22) |
| | <i>Arthrocnemum subterminale</i> | 27 (-) | 1.7 (-) | 1 (-) | 2 (-) |
| | <i>Atriplex prostrata</i> | 27 (-) | 8.7 (-) | 1 (-) | 22 (-) |
| | <i>Cressa truxillensis</i> | 27 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Heliotropium curassavicum</i> | 27 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| | * <i>Sonchus oleraceus</i> | 27 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

Mediterranean California Naturalized Annual and Perennial Grassland Alliance Mediterranean California Naturalized Annual and Perennial Grassland Association

Samples Used To Describe Association (n= 17)

Rapid Assessment(s), WR and WSD stands: SDRP0146, 1695LS01, SDG0035, 0519TK05, 1684FS01, SDG0047, 0309TK01, SDG0015, SDG0023, SDG0024, SDG0119, SDG0050, 0324DHF5, 1606JD02
Rapid Assessment(s), OC stands (n=3): 1099EK02, 1197EK01, 1471EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 42.6 (45.3) | 15-85 (27-55) | - |
| Herb cover (%) | 39.2 (45.0) | 10-84 (26-55) | 0-1 (0-0.5) |
| Shrub cover (%) | 0.7 (0.7) | 0.2-1 (0.2-1) | 0-2 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (2,0), NW (4,1), S (1,1), SE (2,0), SW (8,1)

Macrotopography: middle to upper 1/3 of slope (1,0), upper 1/3 of slope to ridgetop (1,1), upper 1/3 of slope (1,1), ridge top (1,0), lower to upper 1/3 of slope (1,0), lower to middle 1/3 of slope (2,0), lower 1/3 of slope (4,0), bottom (3,0), middle 1/3 of slope (3,1)

Microtopography: concave (1,0), convex (4,1), flat (3,1), undulating (9,1)

Parent material: granitic (2,0), gravelly alluvium (1,0), igneous (1,0), metamorphic (1,0), sandstone (2,2), shale (1,0)

Soil texture: sandy loam (6,2), silty loam (1,0), clay loam (3,1)

| | Average | Range |
|-----------------------|-------------|-----------------|
| Elevation (ft) | 647 (489) | 0-1378 (75-735) |
| Slope (degrees) | 8 (1.7) | 1-26 (1-2) |
| Large rock cover (%) | 3.5 (-) | 0-37 (-) |
| Small rock cover (%) | 7.6 (1.0) | 0-54 (0-2) |
| Bare ground cover (%) | 21.1 (44.5) | 0-64 (36-53) |
| Litter cover (%) | 21.9 (52.5) | 0-72 (45-60) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 115

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|-------------------------------|----------|-------------|-----------|-----------|
| Shrub | <i>Artemisia californica</i> | 53 (67) | 0.5 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Malosma laurina</i> | 41 (-) | 1.2 (-) | 0.2 (-) | 4 (-) |
| | <i>Eriogonum fasciculatum</i> | 35 (67) | 0.6 (0.2) | 0.2 (0.2) | 2 (0.2) |
| | <i>Salvia apiana</i> | 29 (-) | 1.1 (-) | 0.2 (-) | 2 (-) |
| Herb | <i>*Bromus madritensis</i> | 59 (100) | 4.1 (6.7) | 0.2 (0.2) | 15 (15) |
| | <i>*Bromus hordeaceus</i> | 41 (67) | 6.3 (15) | 1 (10) | 20 (20) |
| | <i>*Erodium botrys</i> | 41 (67) | 32.1 (27.5) | 10 (25) | 60 (30) |
| | <i>*Avena barbata</i> | 35 (67) | 1.9 (0.2) | 0.2 (0.2) | 5 (0.2) |
| | <i>*Bromus diandrus</i> | 35 (67) | 3.9 (7.5) | 0.2 (5) | 10 (10) |
| | <i>Stipa pulchra</i> | 35 (33) | 0.3 (0.2) | 0.2 (0.2) | 1 (0.2) |
| | <i>*Centaurea melitensis</i> | 29 (33) | 6.4 (2) | 0.2 (2) | 16 (2) |
| | <i>Croton setiger</i> | 29 (67) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>*Erodium</i> | 29 (-) | 25 (-) | 7 (-) | 70 (-) |
| | <i>*Hirschfeldia incana</i> | 29 (33) | 1.5 (0.2) | 0.2 (0.2) | 5 (0.2) |

***Muhlenbergia rigens* Alliance**
***Muhlenbergia rigens* Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: 1036MB01, 0311RB22
Rapid Assessment(s), OC stands (n=2): 1140MA01, 1172MA03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|-------------------|-------------|
| Total vegetation cover (%) | 73.3 (71.0) | 70-79 (70-72) | - |
| Herb cover (%) | 73.3 (71.0) | 70-79 (70-72) | 0-1 (0.5-1) |
| Shrub cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 0-2 (1-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NW (1,0), SE (1,0), W (2,2)
Macrotopography: bottom (3,2), middle to upper 1/3 of slope (1,0)
Microtopography: flat (2,1), undulating (1,1), convex (1,0)
Parent material: granitic (3,1)
Soil texture: sand (1,1), clay loam (3,1)

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 2277 (2889) | 1260-3330 (2448-3330) |
| Slope (degrees) | 3 (3.0) | 1-3 (3-3) |
| Large rock cover (%) | 0.8 (0.5) | 0-1 (0-1) |
| Small rock cover (%) | 9.3 (15.0) | 0-30 (0-30) |
| Bare ground cover (%) | 9.8 (12.5) | 3-15 (10-15) |
| Litter cover (%) | 77.0 (67.5) | 55-93 (55-80) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 116

Stand Table

| | Species Name | Con | Ave | Min | Max |
|--------------|--|-----------|-----------|-----------|-----------|
| Shrub | <i>Baccharis salicifolia</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Cercocarpus minutiflorus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Quercus acutidens</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | | | | | |
| Herb | <i>Muhlenbergia rigens</i> | 100 (100) | 39.3 (51) | 15 (35) | 67 (67) |
| | * <i>Sonchus asper</i> | 75 (100) | 1.5 (2.1) | 0.2 (0.2) | 4 (4) |
| | <i>Thalictrum fendleri</i> | 75 (50) | 3.5 (0.2) | 0.2 (0.2) | 10 (0.2) |
| | * <i>Anagallis arvensis</i> | 50 (100) | 4.5 (4.5) | 2 (2) | 7 (7) |
| | * <i>Bromus diandrus</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| | * <i>Carduus pycnocephalus</i> | 50 (-) | 22.6 (-) | 0.2 (-) | 45 (-) |
| | <i>Drymocallis lactea</i> | 50 (100) | 1.1 (1.1) | 0.2 (0.2) | 2 (2) |
| | * <i>Festuca myuros</i> | 50 (50) | 13.5 (7) | 7 (7) | 20 (7) |
| | <i>Sidalcea malviflora</i> | 50 (50) | 2 (2) | 2 (2) | 2 (2) |
| | * <i>Agrostis</i> | 25 (-) | 25 (-) | 25 (-) | 25 (-) |
| | <i>Barbarea orthoceras</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Bromus hordeaceus</i> | 25 (50) | 10 (10) | 10 (10) | 10 (10) |
| | * <i>Bromus madritensis</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | * <i>Bromus rubens</i> | 25 (-) | 5 (-) | 5 (-) | 5 (-) |
| | * <i>Bromus tectorum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | * <i>Centaurea melitensis</i> | 25 (50) | 4 (4) | 4 (4) | 4 (4) |
| | * <i>Cerastium glomeratum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Cirsium occidentale</i> var. <i>occidentale</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Cirsium vulgare</i> | 25 (50) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Claytonia perfoliata</i> | 25 (-) | 6 (-) | 6 (-) | 6 (-) |
| | <i>Deinandra fasciculata</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Erodium botrys</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Euphorbia spathulata</i> | 25 (50) | 1 (1) | 1 (1) | 1 (1) |
| | <i>Galium angustifolium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Galium aparine</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| | * <i>Galium parisiense</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| | <i>Galium porrigens</i> var. <i>porrigens</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Geranium dissectum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Hirschfeldia incana</i> | 25 (50) | 10 (10) | 10 (10) | 10 (10) |
| | * <i>Hypochaeris glabra</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Juncus macrophyllus</i> | 25 (50) | 12 (12) | 12 (12) | 12 (12) |
| | * <i>Lathyrus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Lotus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Nassella</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | * <i>Phalaris aquatica</i> | 25 (50) | 2 (2) | 2 (2) | 2 (2) |
| | <i>Pseudognaphalium californicum</i> | 25 (50) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Sidalcea malviflora</i> subsp. | 25 (-) | 5 (-) | 5 (-) | 5 (-) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 117

| Species Name | Con | Ave | Min | Max |
|--|--------|---------|---------|---------|
| <i>sparsifolia</i> | | | | |
| <i>Sisyrinchium bellum</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Stachys rigida</i> var. <i>rigida</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Phalaris aquatica* Alliance**

***Phalaris aquatica* Semi-Natural Stands**

Samples Used To Describe Association (n= 1)

Rapid Assessment(s), WR and WSD stands: None

Rapid Assessment(s), OC stands (n=1): 1172MA02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|---------------|
| Total vegetation cover (%) | 70.0 (70.0) | 70-70 (70-70) | - |
| Herb cover (%) | 70.0 (70.0) | 70-70 (70-70) | 0.5-1 (0.5-1) |
| Shrub cover (%) | - (-) | - (-) | - (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (1,1)

Macrotopography: bottom (1,1)

Microtopography: concave (1,1)

Parent material:

Soil texture:

| | Average | Range |
|-----------------------|-------------|-----------------------|
| Elevation (ft) | 3310 (3310) | 3310-3310 (3310-3310) |
| Slope (degrees) | 5 (5.0) | 5-5 (5-5) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 5.0 (5.0) | 5-5 (5-5) |
| Litter cover (%) | 90.0 (90.0) | 90-90 (90-90) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-----------------------------|-----------|-----------|-----------|-----------|
| Herb | | | | |
| * <i>Anagallis arvensis</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Barbarea orthoceras</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| * <i>Bromus diandrus</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus hordeaceus</i> | 100 (100) | 10 (10) | 10 (10) | 10 (10) |
| * <i>Cirsium vulgare</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Dryocallis lactea</i> | 100 (100) | 4 (4) | 4 (4) | 4 (4) |
| * <i>Festuca myuros</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Hypochaeris glabra</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Muhlenbergia rigens</i> | 100 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Phalaris aquatica</i> | 100 (100) | 51 (51) | 51 (51) | 51 (51) |
| * <i>Phalaris minor</i> | 100 (100) | 2 (2) | 2 (2) | 2 (2) |
| <i>Sidalcea malviflora</i> | 100 (100) | 1 (1) | 1 (1) | 1 (1) |

***Schoenoplectus acutus* Alliance**
***Schoenoplectus acutus* Association**

Samples Used To Describe Association (n= 2)

Rapid Assessment(s), WR and WSD stands: SDRP0269

Rapid Assessment(s), OC stands (n=1): 1128MA02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|--------------|
| Total vegetation cover (%) | 85.0 (90.0) | 80-90 (90-90) | - |
| Herb cover (%) | 80.0 (85.0) | 75-85 (85-85) | 1-5 (2-5) |
| Shrub cover (%) | 2.0 (2.0) | 2-2 (2-2) | 0-5 (2-5) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 1.5 (2.0) | 1-2 (2-2) | 5-15 (10-15) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (1,1), NE (1,0)

Macrotopography: bottom (1,1), lower 1/3 of slope (1,0)

Microtopography: flat (1,1), convex (1,0)

Parent material: granitic (1,0), sedimentary (1,1)

Soil texture: muck (1,1), sandy loam (1,0)

| | Average | Range |
|-----------------------|-----------|-------------------|
| Elevation (ft) | 358 (381) | 335-381 (381-381) |
| Slope (degrees) | 8 (0.0) | 0-15 (0-0) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 10.0 (-) | 10-10 (-) |
| Litter cover (%) | - (-) | - (-) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|---|-----------|-----------|---------|---------|
| Tree | | | | |
| <i>Salix gooddingii</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Salix laevigata</i> | 50 (100) | 2 (2) | 2 (2) | 2 (2) |
| Shrub | | | | |
| <i>Baccharis salicifolia</i> | 100 (100) | 5 (2) | 2 (2) | 8 (2) |
| <i>Baccharis pilularis</i> | 50 (-) | 9 (-) | 9 (-) | 9 (-) |
| <i>Isocoma menziesii</i> | 50 (-) | 5 (-) | 5 (-) | 5 (-) |
| <i>Opuntia littoralis</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Tamarix</i> | 50 (-) | 2 (-) | 2 (-) | 2 (-) |
| Herb | | | | |
| <i>Schoenoplectus acutus</i> var. <i>occidentalis</i> | 100 (100) | 67.5 (85) | 50 (85) | 85 (85) |
| <i>Ambrosia psilostachya</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Avena</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Brassica nigra</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Bromus diandrus</i> | 50 (-) | 5 (-) | 5 (-) | 5 (-) |
| * <i>Bromus hordeaceus</i> | 50 (-) | 3 (-) | 3 (-) | 3 (-) |
| * <i>Centaurea melitensis</i> | 50 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Distichlis spicata</i> | 50 (-) | 5 (-) | 5 (-) | 5 (-) |



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 119

| Species Name | Con | Ave | Min | Max |
|----------------------|----------|-----------|-----------|-----------|
| <i>Urtica dioica</i> | 50 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |

***Spartina foliosa* Alliance**
***Spartina foliosa* Association**

Samples Used To Describe Association (n= 8)

Rapid Assessment(s), WR and WSD stands: 1023AK01, 1021AK04, 1023RB02, 1021TK02, 1023AK06, 1022TK01, 1021TK07

Rapid Assessment(s), OC stands (n=1): 1472EK02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|-------------|
| Total vegetation cover (%) | 32.6 (60.0) | 11-60 (60-60) | - |
| Herb cover (%) | 32.6 (60.0) | 11-60 (60-60) | 0-1 (0-0.5) |
| Shrub cover (%) | - (-) | - (-) | 0-0 (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: Flat (6,0), SE (2,1)

Macrotopography: bottom (8,1)

Microtopography: flat (8,1)

Parent material: mixed alluvium (1,0), silty alluvium (5,0)

Soil texture: muck (1,0), sandy loam (1,0), silt (2,0), silty loam (1,0), clay (2,0)

| | Average | Range |
|------------------------------|-------------|--------------|
| Elevation (ft) | 4 (33) | 0-33 (33-33) |
| Slope (degrees) | 0 (0.0) | 0-0 (0-0) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | - (-) | - (-) |
| Bare ground cover (%) | 10.0 (42.0) | 0-42 (42-42) |
| Litter cover (%) | 6.6 (30.0) | 0-30 (30-30) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|----------------------------|-----------|-----------|----------|---------|
| Herb | | | | |
| <i>Salicornia pacifica</i> | 100 (100) | 3.6 (12) | 0.2 (12) | 12 (12) |
| <i>Spartina foliosa</i> | 100 (100) | 22.3 (18) | 10 (18) | 55 (18) |
| <i>Batis maritima</i> | 50 (100) | 11.6 (30) | 0.2 (30) | 30 (30) |
| Algae | | | | |
| <i>Algae</i> | 25 (-) | 18.5 (-) | 2 (-) | 35 (-) |

***Stipa lepida* Provisional Alliance**
***Stipa lepida* Provisional Association**

Samples Used To Describe Association (n= 4)

Rapid Assessment(s), WR and WSD stands: SDG0090, 0504AK22, 0310RB25
Rapid Assessment(s), OC stands (n=1): 1137MA02

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|---------------------------------|-------------|---------------|-------------|
| Total vegetation cover (%) | 33.5 (65.0) | 22-65 (65-65) | - |
| Herb cover (%) | 32.8 (65.0) | 21-65 (65-65) | 0-1 (0.5-1) |
| Shrub cover (%) | 2.0 (2.0) | 2-2 (2-2) | 0-2 (0.5-1) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | - (-) | - (-) | - (-) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (2,1), SW (2,0)
Macrotopography: lower 1/3 of slope (3,1), lower to middle 1/3 of slope (1,0)
Microtopography: convex (1,0), flat (1,0), undulating (2,1)
Parent material: conglomerate (1,0), sandstone (1,0), sedimentary (2,1)
Soil texture: sandy loam (1,0), clay (1,0), clay loam (1,0), loam (1,1)

| | Average | Range |
|-----------------------|-------------|-----------------|
| Elevation (ft) | 321 (463) | 0-676 (463-463) |
| Slope (degrees) | 13 (15.0) | 4-16 (15-15) |
| Large rock cover (%) | - (-) | - (-) |
| Small rock cover (%) | 0.8 (-) | 0-3 (-) |
| Bare ground cover (%) | 55.0 (4.0) | 4-89 (4-4) |
| Litter cover (%) | 41.3 (93.0) | 4-93 (93-93) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------|--------------------------------------|-----------|-----------|-----------|-----------|
| Tree | | | | | |
| | <i>Standing Snag</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| Shrub | | | | | |
| | <i>Acemispn glaber</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Artemisia californica</i> | 50 (100) | 1.5 (2) | 1 (2) | 2 (2) |
| | <i>Isocoma menziesii</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| | <i>Baccharis pilularis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Baccharis sarothroides</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Encelia californica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Eriogonum fasciculatum</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| | <i>Grindelia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Malosma laurina</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| | <i>Peritoma arborea var. arborea</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| Herb | | | | | |
| | <i>Stipa lepida</i> | 100 (100) | 9.3 (12) | 6 (12) | 12 (12) |
| | * <i>Avena fatua</i> | 75 (100) | 1.7 (4) | 0.2 (4) | 4 (4) |
| | * <i>Bromus madritensis</i> | 75 (-) | 4.7 (-) | 1 (-) | 12 (-) |
| | * <i>Brachypodium distachyon</i> | 50 (100) | 25 (49) | 1 (49) | 49 (49) |
| | * <i>Brassica nigra</i> | 50 (100) | 2.6 (5) | 0.2 (5) | 5 (5) |

Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 121

| Species Name | Con | Ave | Min | Max |
|-----------------------------------|----------|-----------|-----------|-----------|
| <i>Deinandra fasciculata</i> | 50 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Festuca myuros</i> | 50 (-) | 4.1 (-) | 0.2 (-) | 8 (-) |
| * <i>Hypochaeris glabra</i> | 50 (-) | 0.6 (-) | 0.2 (-) | 1 (-) |
| <i>Acmispon micranthus</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| * <i>Anagallis arvensis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Astragalus gambelianus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Bloomeria crocea</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| * <i>Bromus diandrus</i> | 25 (100) | 7 (7) | 7 (7) | 7 (7) |
| * <i>Bromus hordeaceus</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Calochortus superbus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Calystegia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Cardionema ramosissimum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Carduus pycnocephalus</i> | 25 (100) | 1 (1) | 1 (1) | 1 (1) |
| * <i>Centaurea melitensis</i> | 25 (-) | 2 (-) | 2 (-) | 2 (-) |
| <i>Centaurium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Corethrogyne filaginifolia</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Crassula connata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Cressa truxillensis</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Croton californicus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Croton setiger</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Cynara cardunculus</i> | 25 (100) | 0.2 (0.2) | 0.2 (0.2) | 0.2 (0.2) |
| <i>Daucus pusillus</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Distichlis spicata</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| <i>Dudleya pulverulenta</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Erodium botrys</i> | 25 (-) | 4 (-) | 4 (-) | 4 (-) |
| * <i>Erodium cicutarium</i> | 25 (-) | 9 (-) | 9 (-) | 9 (-) |
| * <i>Foeniculum vulgare</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| <i>Galium angustifolium</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| * <i>Helminthotheca echioides</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Lepidium nitidum</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Linanthus dianthiflorus</i> | 25 (-) | 1 (-) | 1 (-) | 1 (-) |
| * <i>Logfia gallica</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Plantago ovata</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Porophyllum gracile</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Pterostegia drymarioides</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |
| <i>Sisyrinchium bellum</i> | 25 (-) | 3 (-) | 3 (-) | 3 (-) |
| <i>Stipa pulchra</i> | 25 (-) | 0.2 (-) | 0.2 (-) | 0.2 (-) |

***Stipa pulchra* Alliance**
***Stipa pulchra* Association**

Samples Used To Describe Association (n= 41)

Rapid Assessment(s), WR and WSD stands: 1548JD02, 0505RB21, 0505RB24, 0506DH22, SRPL.018, 1690FS01, 0507AK21, 0505RB22, 1761EB01, SDG0089, 1616KG02, 1617MM01, 1158FS01, 0309DH23, WILD.007/BPU 392-361, MURR.011, 1707EB01, SDG0120, 1706EB01, SRPL.004/437, SDRP0094, SDRP0357, SRPL.016, 0624RB02

Rapid Assessment(s), OC stands (n=17): 1003EK01, 1003EK05, 1014EK02, 1015EK04, 1025EK01, 1028EK02, 1047EK02, 1056EK04, 1092EK06, 1110EK01, 1289MA03, 1350EK03, 1364EK01, 1450EK02, 1463EK03, 1465EK01, 3.8jd03

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|----------------|-------------------|-------------------|
| Total vegetation cover (%) | 51.5 (46.2) | 28-90 (29-85) | - |
| Herb cover (%) | 49.3 (42.7) | 27-90 (29-77) | 0-1 (0-1) |
| Shrub cover (%) | 1.8 (2.1) | 0-6 (0.2-6) | 0-5 (0.5-2) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | 0.5-5 (-) |
| Overstory tree cover (%) | 0.2 (0.2) | 0.2-0.2 (0.2-0.2) | 2-10 (2-10) |

Local Environmental Description (All stands, OC stands)

Aspect: NE (11,4), W (1,1), Variable (4,0), SW (3,1), SE (5,3), NW (9,4), N (1,1), Flat (5,1), E (1,1), S (1,1)

Macrotopography: middle 1/3 of slope (14,7), ridge top (2,1), upper 1/3 of slope (4,4), middle to upper 1/3 of slope (2,1), lower to upper 1/3 of slope (1,0), lower to middle 1/3 of slope (1,0), lower 1/3 of slope (8,3), bottom (5,0), bottom to lower 1/3 of slope (3,0)

Microtopography: convex (9,4), flat (17,11), concave (2,0), undulating (13,2)

Parent material: gabbro (1,0), granitic (3,0), igneous (2,0), mixed alluvium (1,0), sandstone (13,12), sandy alluvium (1,0), sedimentary (3,0), volcanic flow (3,0)

Soil texture: silty loam (1,0), sandy loam (8,1), loam (4,0), clay (12,7), clay loam (16,9)

| | Average | Range |
|------------------------------|----------------|--------------------|
| Elevation (ft) | 823 (794) | 62-1998 (151-1476) |
| Slope (degrees) | 9 (10.8) | 0-30 (0-30) |
| Large rock cover (%) | 1.3 (-) | 0-15 (-) |
| Small rock cover (%) | 6.3 (0.4) | 0-34 (0-5) |
| Bare ground cover (%) | 28.5 (28.8) | 0-88 (4-53) |
| Litter cover (%) | 54.6 (68.4) | 0-94 (45-94) |

Stand Table

| Species Name | Con | Ave | Min | Max |
|-------------------------------|------------|-------------|------------|------------|
| Shrub | | | | |
| <i>Isocoma menziesii</i> | 46 (59) | 1 (1) | 0.2 (0.2) | 3 (3) |
| <i>Artemisia californica</i> | 41 (59) | 0.6 (0.5) | 0.2 (0.2) | 2 (2) |
| <i>Eriogonum fasciculatum</i> | 37 (18) | 0.7 (2.1) | 0.2 (0.2) | 5 (5) |
| <i>Opuntia littoralis</i> | 27 (59) | 0.5 (0.5) | 0.2 (0.2) | 1 (1) |
| Herb | | | | |
| <i>Nassella pulchra</i> | 100 (-) | 14.4 (-) | 2 (-) | 50 (-) |
| <i>Stipa pulchra</i> | 100 (100) | 14.4 (15.6) | 2 (2) | 50 (40) |
| * <i>Avena barbata</i> | 61 (76) | 7.8 (8.2) | 0.2 (0.2) | 24 (24) |
| * <i>Bromus diandrus</i> | 51 (76) | 11.3 (12.2) | 0.2 (0.2) | 60 (60) |
| * <i>Bromus hordeaceus</i> | 49 (59) | 5.1 (6.7) | 0.2 (0.2) | 25 (25) |
| * <i>Bromus madritensis</i> | 37 (35) | 4.1 (1.1) | 0.2 (0.2) | 20 (3) |
| * <i>Erodium botrys</i> | 37 (41) | 10.8 (0.9) | 0.2 (0.2) | 70 (2) |
| <i>Sisyrinchium bellum</i> | 32 (35) | 0.9 (1) | 0.2 (0.2) | 5 (5) |

| Species Name | Con | Ave | Min | Max |
|--------------------------------|---------|-----------|-----------|----------|
| <i>Calochortus splendens</i> | 29 (18) | 0.6 (0.2) | 0.2 (0.2) | 2 (0.2) |
| * <i>Centaurea melitensis</i> | 29 (18) | 0.9 (2.1) | 0.2 (0.2) | 5 (5) |
| <i>Dichelostemma capitatum</i> | 29 (47) | 0.4 (0.2) | 0.2 (0.2) | 2 (0.2) |
| * <i>Hirschfeldia incana</i> | 29 (12) | 1.3 (0.2) | 0.2 (0.2) | 10 (0.2) |
| * <i>Cynara cardunculus</i> | 27 (35) | 3.2 (2.3) | 0.2 (0.2) | 18 (10) |

Typha (angustifolia, domingensis, latifolia) Alliance
***Typha domingensis* Association**

Samples Used To Describe Association (n= 10)

Rapid Assessment(s), WR and WSD stands: 1571EB01, 0623DH01, 0708MG02, 0708MG04, 1020RB01, 1020TK05, 1021RB05, 0309AK22, 1021AK01

Rapid Assessment(s), OC stands (n=1): 1517EK01

Local Vegetation Description All stands (OC stands)

| | Average | Range | Height (m) |
|--|-------------|---------------|------------|
| Total vegetation cover (%) | 58.6 (61.0) | 23-84 (61-61) | - |
| Herb cover (%) | 57.1 (61.0) | 21-83 (61-61) | 1-10 (2-5) |
| Shrub cover (%) | - (-) | - (-) | 0-5 (-) |
| Tall shrub / low tree cover (%) | - (-) | - (-) | - (-) |
| Overstory tree cover (%) | 1.8 (-) | 0.2-5 (-) | 2-10 (-) |

Local Environmental Description (All stands, OC stands)

Aspect: SW (2,1), Flat (7,0), NE (1,0)

Macrotopography: bottom to lower 1/3 of slope (1,0), bottom (9,1)

Microtopography: undulating (1,0), concave (1,0), flat (8,1)

Parent material: clayey alluvium (2,0), mixed alluvium (1,0), sandy alluvium (1,0), silty alluvium (2,0), volcanic flow (1,0)

Soil texture: silt (1,0), muck (8,1), sandy loam (1,0)

| | Average | Range |
|------------------------------|-------------|---------------|
| Elevation (ft) | 47 (16) | 0-240 (16-16) |
| Slope (degrees) | 0 (0.0) | 0-1 (0-0) |
| Large rock cover (%) | 1.6 (-) | 0-10 (-) |
| Small rock cover (%) | 2.0 (-) | 0-20 (-) |
| Bare ground cover (%) | 26.4 (31.0) | 0-87 (31-31) |
| Litter cover (%) | 23.8 (35.0) | 0-94 (35-35) |

Stand Table

| | Species Name | Con | Ave | Min | Max |
|-------------|-----------------------------------|-----------|---------|---------|---------|
| Tree | | | | | |
| | <i>Salix gooddingii</i> | 30 (-) | 0.7 (-) | 0.2 (-) | 1 (-) |
| Herb | | | | | |
| | <i>Typha domingensis</i> | 100 (100) | 45 (60) | 5 (60) | 78 (60) |
| | * <i>Apium graveolens</i> | 50 (-) | 2.6 (-) | 0.2 (-) | 6 (-) |
| | * <i>Helminthotheca echioides</i> | 40 (-) | 1.6 (-) | 0.2 (-) | 5 (-) |
| | * <i>Polypogon monspeliensis</i> | 40 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |
| | * <i>Rumex crispus</i> | 40 (-) | 0.9 (-) | 0.2 (-) | 2 (-) |
| | <i>Xanthium strumarium</i> | 40 (-) | 0.4 (-) | 0.2 (-) | 1 (-) |



Vegetation Classification Report
Nature Reserve of Orange County
January 15, 2013
Page 124

| Species Name | Con | Ave | Min | Max |
|---------------------------------------|--------|---------|---------|--------|
| <i>Anemopsis californica</i> | 30 (-) | 0.5 (-) | 0.2 (-) | 1 (-) |
| <i>Distichlis spicata</i> | 30 (-) | 3.5 (-) | 0.2 (-) | 10 (-) |
| <i>Juncus acutus subsp. leopoldii</i> | 30 (-) | 1.1 (-) | 0.2 (-) | 3 (-) |
| <i>Pluchea odorata</i> | 30 (-) | 6.1 (-) | 0.2 (-) | 18 (-) |