

1 Measures

Let E be any set. A collection \mathcal{E} of subsets of E is called a σ -algebra if the following holds:

1. $\emptyset \in \mathcal{E}$.
2. If $A \in \mathcal{E}$, then $A^c = E \setminus A \in \mathcal{E}$.
3. If $(A_n : n \in \mathbb{N})$, $A_n \in \mathcal{E}$, then $\bigcup_n A_n \in \mathcal{E}$.

Examples.

- $\mathcal{E} = \{\emptyset, E\}$
- $\mathcal{E} = \mathcal{P}(E)$, the set of all subsets of E .

Note that $\bigcap_n A_n = (\bigcup_n A_n^c)^c$, any σ -algebra \mathcal{E} is also closed under countable intersection of its elements. Also $B \setminus A = B \cap A^c \in \mathcal{E}$ whenever $A, B \in \mathcal{E}$.

Any set E with a choice of σ -algebra \mathcal{E} is called a *measurable space*, and the elements of \mathcal{E} are called *measurable sets*.

A *measure* μ is a set-function $\mu : \mathcal{E} \rightarrow [0, \infty]$ such that $\mu(\emptyset) = 0$, and for any $(A_n : n \in \mathbb{N})$, $A_n \in \mathcal{E}$ pairwise disjoint ($A_n \cap A_m = \emptyset$ for all $n \neq m$) then

$$\mu\left(\bigcup_n A_n\right) = \sum_n \mu(A_n) \quad (\text{countable additivity of } \mu)$$

If \mathcal{E} is countable, then for any $A \in \mathcal{P}(E)$ and a measure μ

$$\mu(A) = \mu\left(\bigcup_{x \in A} \{x\}\right) = \sum_{x \in A} \mu(\{x\})$$

hence there is a one-to-one correspondence between measures and mass functions on E .

For any collection \mathcal{A} of subsets of E , we define the σ -algebra $\sigma(\mathcal{A})$ generated by \mathcal{A} as

$$\sigma(\mathcal{A}) = \{A \subseteq E : A \in \mathcal{E} \text{ } \forall \sigma\text{-algebras } \mathcal{E} \supseteq \mathcal{A}\}$$

which equals (Example sheet)

$$\sigma(\mathcal{A}) = \bigcap_{\sigma\text{-algebras } \mathcal{E} \supseteq \mathcal{A}} \mathcal{E}$$

To construct good ‘generators’ we define

1. \mathcal{A} is called a *ring over E* if $\emptyset \in \mathcal{A}$; if $A, B \in \mathcal{A}$ then $B \setminus A \in \mathcal{A}$ and $A \cup B \in \mathcal{A}$.

2. \mathcal{A} is called an *algebra over E* if $\emptyset \in \mathcal{A}$; if $A, B \in \mathcal{A}$ then $A^c \in \mathcal{A}$, $A \cup B \in \mathcal{A}$.

Notice that in a ring $A \Delta B = (B \setminus A) \cup (A \setminus B) \in \mathcal{A}$ and $A \cap B = (A \cup B) \setminus (A \Delta B) \in \mathcal{A}$. Also, $B \setminus A = B \cap A^c = (B^c \cup A)^c \in \mathcal{A}$, so an algebra is a ring.

Fact: If $\bigcup_n A_n$, $A_n \in \mathcal{E}$, \mathcal{E} some σ -algebra (or a ring if the union is finite) - then we can find $B_n \in \mathcal{E}$ disjoint such that $\bigcup_n A_n = \bigcup_n B_n$. Indeed, define $\tilde{A}_n = \bigcup_{j \leq n} A_j$, and set $B_n = \tilde{A}_n \setminus \tilde{A}_{n-1}$, then the fact follows. [“disjointification of countable unions”]

Definition. A *set function* on any collection \mathcal{A} of subsets of E (where $\emptyset \in \mathcal{A}$) is a map $\mu : \mathcal{A} \rightarrow [0, \infty]$ such that $\mu(\emptyset) = 0$. We say μ is

1. *increasing* if $\mu(A) \leq \mu(B)$ whenever $A \subseteq B$; $A, B \in \mathcal{A}$
2. *additive* if $\mu(A \cup B) = \mu(A) + \mu(B)$ whenever $A, B \in \mathcal{A}$; $A \cup B \in \mathcal{A}$ and $A \cap B = \emptyset$.
3. *countably additive* if $\mu(\bigcup_n A_n) = \sum_n \mu(A_n)$ for any $(A_n : n \in \mathbb{N})$ where $A_n \in \mathcal{A}$ disjoint and $\bigcup_n A_n \in \mathcal{A}$.
4. *countably sub-additive* if $\mu(\bigcup_n A_n) \leq \sum_n \mu(A_n)$ for all $(A_n : n \in \mathbb{N})$ such that $\bigcup_n A_n \in \mathcal{A}$

Remark: one can show that a measure μ on a σ -algebra satisfies 1-4 above.

Theorem (Caratheodory). *Let μ be a countably additive set function on a ring \mathcal{A} of subsets of E . Then there exists a measure μ^* on $\sigma(\mathcal{A})$ such that $\mu^*|_{\mathcal{A}} = \mu$.*

Proof. For $B \subseteq E$ define the *outer measure* μ^* as

$$\mu^*(B) = \inf \left\{ \sum_{n \in \mathbb{N}} \mu(A_n) : A_n \in \mathcal{A}, B \subseteq \bigcup_n A_n \right\}$$

and set $\mu^*(B) = \infty$ if the set within the infimum is empty.

Define

$$\mathcal{M} = \{A \subseteq E : \mu^*(B) = \mu^*(B \cap A) + \mu^*(B \cap A^c) \forall B \subseteq E\}$$

the “ μ^* -measurable” sets.

Step 1: μ^* is countably sub-additive on $\mathcal{P}(E)$. For any $B \subseteq E$ and $B_n \subseteq E$ such that $B \subseteq \bigcup_n B_n$ we have

$$\mu^*(B) \leq \sum_n \mu^*(B_n) \quad (\dagger)$$

WLOG we assume $\mu^*(B_n) < \infty$ for all n so for all $\varepsilon > 0$, there exists A_{nm} such that $B_n \subseteq \bigcup_m A_{nm}$ and

$$\mu^*(B_n) + \frac{\varepsilon}{2^n} \geq \sum_m \mu(A_{nm})$$

Now since μ^* and since $B \subseteq \bigcup_n B_n \subseteq \bigcup_{n,m} A_{nm}$, hence

$$\mu^*(B) \leq \mu^*\left(\bigcup_{n,m} A_{nm}\right) \leq \sum_{n,m} \mu(A_{nm}) \leq \sum_n \mu^*(B_n) + \underbrace{\sum_n \frac{\varepsilon}{2^n}}_{=\varepsilon}$$

so (\dagger) follows since ε was arbitrary.

Step 2: μ^* extends μ . Let $A \in \mathcal{A}$. Clearly $A = A \cup \emptyset \cup \dots \cup \emptyset$, so by definition of μ^* , $\mu^*(A) \leq \mu(A) + 0 + \dots + 0$. So we need to prove $\mu(A) \leq \mu^*(A)$. Again, assume $\mu^*(A) < \infty$ WLOG, and let $A_n \in \mathcal{A}$ be such that $A \subseteq \bigcup_n A_n$. Then $A = \bigcup_n \underbrace{(A \cap A_n)}_{\in \mathcal{A}}$, and since μ is countably sub-additive on \mathcal{A} , we have

$$\mu(A) = \mu\left(\bigcup_n (A \cap A_n)\right) \leq \sum_n \underbrace{\mu(A \cap A_n)}_{\subseteq A_n} \leq \sum_n \mu(A_n)$$

so since the (A_n) were arbitrary, by taking infima, we have $\mu(A) \leq \mu^*(A)$.

Step 3: $\mathcal{M} \supseteq \mathcal{A}$. Let $A \in \mathcal{A}$, then $B \subseteq (B \cap A) \cup (B \cap A^c) \cup \emptyset \dots \cup \emptyset = \bigcup_n B_n$ so by (\dagger) we have

$$\mu^*(B) \leq \mu^*(B \cap A) + \mu^*(B \cap A^c) + 0 \dots + 0$$

so we need to prove $\mu^*(B) \geq \mu^*(B \cap A) + \mu^*(B \cap A^c)$. Again, WLOG assume $\mu^*(B) < \infty$, and so for all $\varepsilon > 0$ there exist $A_n \in \mathcal{A}$ such that $B \subseteq \bigcup_n A_n$ and

$$\mu^*(B) + \varepsilon \geq \sum_n \mu(A_n) \quad (\circ)$$

now $B \cap A \subseteq \bigcup_n \underbrace{(A_n \cap A)}_{\in \mathcal{A}}$ and $B \cap A^c \subseteq \bigcup_n \underbrace{(A_n \cap A^c)}_{A_n \setminus A \in \mathcal{A}}$. Therefore by definition of inf in μ^* and additivity of μ

$$\begin{aligned} \mu^*(B \cap A) + \mu^*(B \cap A^c) &\leq \sum_n (\mu(A \cap A_n) + \mu(A^c \cap A_n)) \\ &= \sum_n \mu(A_n) \\ &\underbrace{\leq}_{\circ} \mu^*(B) + \varepsilon \end{aligned}$$

since ε was arbitrary, $\mu^*(B) = \mu^*(B \cap A) + \mu^*(B \cap A^c)$, so $A \in \mathcal{M}$.

Step 4: \mathcal{M} is an algebra. Clearly $\emptyset \in \mathcal{M}$, and by the definition of \mathcal{M} its obvious that $A^c \in \mathcal{M}$ whenever $A \in \mathcal{M}$. So let $A_1, A_2 \in \mathcal{M}$

$$\mu^*(B) = \mu^*(B \cap A_1) + \mu^*(B \cap A_1^c), \text{ since } A_1 \in \mathcal{M}$$

$$= \mu^*(B \cap A_1 \cap A_2) + \mu^*(B \cap A_1 \cap A_2^c) + \mu^*(B \cap A_1^c), \text{ since } A_2 \in \mathcal{M}$$

Clearly $A_1 \cap A_2^c = (A_1 \cap A_2^c) \cap A_1$ and $A_1^c = (A_1 \cap A_2)^c \cap A_1^c$ so

$$\begin{aligned} \mu^*(B) &= \mu^*(B \cap A_1 \cap A_2) + \mu^*(B \cap (A_1 \cap A_2)^c \cap A_1) + \mu^*(B \cap (A_1 \cap A_2)^c \cap A_1^c) \\ &= \mu^*(B \cap A_1 \cap A_2) + \mu^*(B \cap (A_1 \cap A_2)^c), \text{ since } A_1 \in \mathcal{M} \end{aligned}$$

so $A_1 \cap A_2 \in \mathcal{M}$, and \mathcal{M} is an algebra.

Step 5: Let $A = \bigcup_n A_n$, $A_n \in \mathcal{M}$, WLOG A_n disjoint (disjointification). Want $A \in \mathcal{M}$ and $\mu^*(A) = \sum_n \mu^*(A_n)$. By (\dagger) we clearly have

$$\mu^*(B) \leq \mu^*(B \cap A) + \mu^*(B \cap A^c) + 0 \dots + 0$$

and

$$\mu^*(A) \leq \sum_n \mu^*(A_n)$$

so we only need two converse inequalities. Similar to before

$$\begin{aligned}
 \mu^*(B) &= \mu^*(B \cap A_1) + \mu^*(B \cap A_1^c) \\
 &= \mu^*(B \cap A_1) + \mu^*(B \cap \underbrace{A_1^c \cap A_2}_{=A_2 \text{ as disjoint}}) + \mu^*(B \cap A_1^c \cap A_2^c) \\
 &= \sum_{n \leq N} \mu^*(B \cap A_n) + \mu^*(B \cap A_1^c \cap \dots \cap A_N^c)
 \end{aligned}$$

since $\bigcup_{n \leq N} A_n \subseteq A$ so $\bigcap_{n \leq N} A_n^c \supseteq A^c$, taking limits

$$\mu^*(B) \geq \sum_{n=1}^{\infty} \mu^*(B \cap A_n) + \mu^*(B \cap A^c)$$

and by (†)

$$\mu^*(B) \geq \mu^*(B \cap A) + \mu^*(B \cap A^c)$$

so $A \in \mathcal{M}$. Applying the previous with $B = A$, we see

$$\mu^*(A) \geq \sum_{n=1}^{\infty} \mu^*(A \cap A_n) + \mu^*(A \cap A^c) = \sum_n \mu^*(A_n)$$

□

Definition. A collection \mathcal{A} of subsets of E is called a π -system if $\emptyset \in \mathcal{A}$ and if $A, B \in \mathcal{A}$, then $A \cap B \in \mathcal{A}$.

Definition. \mathcal{A} is called a d -system if $E \in \mathcal{A}$, and if $B_1, B_2 \in \mathcal{A}$ such that $B_1 \subseteq B_2$, then $B_2 \setminus B_1 \in \mathcal{A}$, and if $A_n \in \mathcal{A}$, $A_n \uparrow \bigcup_n A_n = A$, then $A \in \mathcal{A}$.

One shows (Example sheet) that a d -system which is also a π -system is a σ -algebra.

Lemma (Dynkin). *Let \mathcal{A} be a π -system. Then any d -system that contains \mathcal{A} also contains $\sigma(\mathcal{A})$.*

Proof. Define

$$\mathcal{D} = \bigcap_{\mathcal{D}' \supseteq \mathcal{A} \text{ a } d\text{-system}} \mathcal{D}'$$

which is again a d -system (Example sheet). We show that \mathcal{D} is a π -system, hence a σ -algebra containing \mathcal{A} . Define

$$\mathcal{D}' = \{B \in \mathcal{D} : B \cap A \in \mathcal{D} \forall A \in \mathcal{A}\}$$

which contains \mathcal{A} as \mathcal{A} is a π -system. Next we show \mathcal{D}' is a d -system. Clearly $E \cap A = A \in \mathcal{A} \subseteq \mathcal{D}$, so $E \in \mathcal{D}'$. Next let $B_1, B_2 \in \mathcal{D}'$ such that $B_1 \subseteq B_2$ then $(B_2 \setminus B_1) \cap A = \underbrace{(B_2 \cap A)}_{\in \mathcal{D}} \setminus \underbrace{(B_1 \cap A)}_{\in \mathcal{D}} \in \mathcal{D}$ and so $B_2 \setminus B_1 \in \mathcal{D}'$.

Next take $B_n \uparrow B$, $B_n \in \mathcal{D}'$ then $\underbrace{B_n \cap A}_{\in \mathcal{D}} \uparrow B \cap A \in \mathcal{D}$ so $B \in \mathcal{D}'$.

Hence \mathcal{D}' is a d -system containing \mathcal{A} , so by minimality of \mathcal{D} , $\mathcal{D} \subseteq \mathcal{D}'$. Conversely, by construction $\mathcal{D}' \subseteq \mathcal{D}$, so $\mathcal{D}' = \mathcal{D}$.

Next define

$$\mathcal{D}'' = \{B \in \mathcal{D} : B \cap A \in \mathcal{D} \forall A \in \mathcal{D}\}$$

which by the preceding step ($\mathcal{D}' = \mathcal{D}$) contains \mathcal{A} . Just as before, one shows that $\mathcal{D}'' = \mathcal{D}$ and so \mathcal{D} is a π -system (as \mathcal{D}'' is by construction). \square

Theorem (Uniqueness of extension). *Let μ_1, μ_2 be measures on (E, \mathcal{E}) such that $\mu_1(E) = \mu_2(E) < \infty$, and suppose $\mu_1 = \mu_2$ on a π -system \mathcal{A} such that $\mathcal{E} \subseteq \sigma(\mathcal{A})$. Then $\mu_1 = \mu_2$ on \mathcal{E} .*

Proof. Define

$$\mathcal{D} = \{A : \mu_1(A) = \mu_2(A)\}$$

which contains \mathcal{A} by hypothesis. We show that \mathcal{D} is a d -system, and hence by Dynkin's Lemma, contains $\sigma(\mathcal{A})$, so the theorem follows.

To see this, note first that $E \in \mathcal{D}$ by hypothesis. Next, by additivity and finiteness of μ_1, μ_2 , for $B_1 \subseteq B_2$, $B_1, B_2 \in \mathcal{D}$.

$$\mu_1(B_2 \setminus B_1) = \mu_1(B_2) - \mu_1(B_1) = \mu_2(B_2) - \mu_2(B_1) = \mu_2(B_2 \setminus B_1)$$

so $B_2 \setminus B_1 \in \mathcal{D}$. Finally take $B_n \uparrow B$, $B_n \in \mathcal{D}$. This implies $B \setminus B_n \downarrow \emptyset$ and (by Example sheet) $\mu_i(B \setminus B_n) \rightarrow \mu_i(\emptyset) = 0$ for $i = 1, 2$. This implies for $\mu_i(B) < \infty$ that $\mu_i(B_n) \rightarrow \mu_i(B)$ as $n \rightarrow \infty$ for both $i = 1, 2$. But then

$$\mu_1(B) = \lim_{n \rightarrow \infty} \mu_1(B_n) = \lim_{n \rightarrow \infty} \mu_2(B_n) = \mu_2(B)$$

and so $B \in \mathcal{D}$, and thus \mathcal{D} is a d-system. \square

Remark: the above theorem applies to finite measures μ such that $\mu(E) < \infty$. The above theorem extends (as we will see) to σ -finite measures μ for which $E = \bigcup_{n \in \mathbb{N}} E_n$ such that $\mu(E_n) < \infty$.

Borel- σ -algebras

Definition. Let E be a topological space (Hausdorff, or metric space). The σ -algebra generated by $\mathcal{A} = \{B \subseteq E : B \text{ is open}\}$ is called the *Borel- σ -algebra*, denoted by $\mathcal{B}(E)$, or just \mathcal{B} when $E = \mathbb{R}$. Elements of $\mathcal{B}(E)$ are the Borel subsets of E . A measure μ on $(E, \mathcal{B}(E))$ is called a *Borel measure on E* . A *Radon* measure μ is a Borel measure such that $\mu(K) < \infty$ for all $K \subseteq E$ compact (closed in Hausdorff spaces, hence measurable).

Construction of Lebesgue measure

We will (eventually) construct a unique Borel measure μ on \mathbb{R}^d such that

$$\mu\left(\prod_{i=1}^d [a_i, b_i]\right) = \prod_{i=1}^d |b_i - a_i|, \quad a_i < b_i, \quad i = 1, \dots, d$$

We will do $d = 1$ first.

Theorem. *There exists a unique Borel measure (called the Lebesgue measure) μ on \mathbb{R} such that*

$$\mu([a, b]) = b - a, \quad \forall a < b$$

Proof. Consider subsets of \mathbb{R} of the form

$$A = (a_1, b_1] \cup \dots \cup (a_n, b_n]$$

which form a ring and π -system of Borel sets. \square