

1 Lebesgue Integration Theory

1.1 Review of measure theory

Definition. Given a set E , a σ -algebra on E is a collection \mathcal{E} of subsets of E such that:

- (i) $E \in \mathcal{E}$;
 - (ii) $A \in \mathcal{E} \Rightarrow A^c = E \setminus A \in \mathcal{A}$;
 - (iii) $A_n \in \mathcal{E}, n \in \mathbb{N} \Rightarrow \bigcup_{n \in \mathbb{N}} A_n \in \mathcal{E}$.
- (E, \mathcal{E}) is called a *measurable space*, and any $A \in \mathcal{E}$ is called a *measurable set*.

Given a collection \mathcal{A} of subsets of E , $\sigma(\mathcal{A})$ is the smallest σ -algebra containing \mathcal{A} .

Definition. A *measure* on (E, \mathcal{E}) is a function $\mu : \mathcal{E} \rightarrow [0, \infty]$ such that

- (i) $\mu(\emptyset) = 0$;
- (ii) $A_n \in \mathcal{E}, n \in \mathbb{N} \text{ disjoint} \Rightarrow \mu\left(\bigcup_{n \in \mathbb{N}} A_n\right) = \sum_{n \in \mathbb{N}} \mu(A_n)$.

(E, \mathcal{E}, μ) is called a *measure space*.

Definition (Borel measure). If (E, τ) is a topological space, then $\sigma(\tau)$ is called a *Borel algebra*, denoted $\mathcal{B}(E)$, and a measure on $(E, \mathcal{B}(E))$ is called a *Borel measure*.

Example. $E = \mathbb{R}^n$, μ the Lebesgue measure satisfying $\mu((a_1, b_1) \times \dots \times (a_n, b_n)) = (b_1 - a_1) \dots (b_n - a_n)$.

Notation: we write $\mu(dx) = dx$ and $\mu(A) = |A|$ when μ is the Lebesgue measure.

Definition (Measurable function). Let (E, \mathcal{E}) and (F, \mathcal{F}) be measurable spaces. Then $f : E \rightarrow F$ is *measurable* if $f^{-1}(A) \in \mathcal{E}$ whenever $A \in \mathcal{F}$. If (E, \mathcal{E}) and (F, \mathcal{F}) are Borel algebras, a measurable function is called a *Borel function*. Special case: $(F, \mathcal{F}) = ([0, \infty], \mathcal{B}([0, \infty]))$, then $f : E \rightarrow F$ is called a *non-negative measurable function*.

Fact. The class of measurable functions is closed under addition, multiplication and taking (pointwise) limits.

Definition. $f : E \rightarrow F$ ($F = [0, \infty]$ or \mathbb{R}^n or \mathbb{C}^n) is a *simple function* if $f = \sum_{k=1}^K a_k \mathbb{1}_{A_k}$ for some $K \in \mathbb{N}$, $a_k \in F$, $A_k \in \mathcal{E}$. For a simple function, the integral is

$$\int f d\mu = \int f(x) d\mu(x) = \sum_{k=1}^K a_k \mu(A_k) \quad (0 \cdot \infty := 0).$$

For a non-negative measurable f , we define

$$\int f d\mu = \sup \left\{ \int g d\mu : g \text{ simple}, 0 \leq g \leq f \right\}.$$

Definition. A measurable function $f : E \rightarrow \mathbb{R}$ is said to be *integrable* if $\int |f| d\mu < \infty$. Write $f = f_+ - f_-$ with f_{\pm} non-negative, measurable, $\int f_{\pm} d\mu < \infty$, and then $\int f d\mu = \int f_+ d\mu - \int f_- d\mu$. For $f : E \rightarrow \mathbb{R}^n$, this is applied in each component.

Theorem (Monotone convergence theorem). *Let (E, \mathcal{E}, μ) be a measure space, and let (f_n) be a (pointwise) increasing sequence of non-negative functions on E converging to f . Then*

$$\lim_{n \rightarrow \infty} \int_E f_n d\mu = \int_E f d\mu.$$

Theorem (Dominated convergence theorem). *Let (f_n) be a sequence of measurable functions on a measure space (E, \mathcal{E}, μ) such that:*

- (i) $f_n \rightarrow f$ pointwise almost everywhere;
- (ii) $|f_n| \leq g$ almost everywhere for some integrable g .

Then

$$\lim_{n \rightarrow \infty} \int_E f_n d\mu = \int_E f d\mu.$$

1.2 L^p spaces

Definition. Let (E, \mathcal{E}, μ) be a measure space. For $p \in [1, \infty)$ and $f : E \rightarrow \mathbb{R}$ define

$$\|f\|_{L^p} = \left(\int_E |f|^p d\mu \right)^{1/p}$$

and

$$\|f\|_{L^\infty} = \text{esssup}|f| = \inf\{K : |f| \leq K \text{ a.e.}\}.$$

The space L^p , $p \in [1, \infty]$ is defined by

$$L^p = L^p(E, \mathcal{E}, \mu) = \{f : E \rightarrow \mathbb{R} \text{ measurable} : \|f\|_{L^p} < \infty\} / \sim.$$

Where $f \sim g$ if $f = g$ a.e.

Theorem (Riesz-Fisher theorem). L^p is a Banach space for all $p \in [1, \infty]$.

Notation: when $E = \mathbb{R}^n$, μ the Lebesgue measure, write $L^p(E, \mu) = L^p(\mathbb{R}^n)$.

Fact. For $p \in [1, \infty)$, the simple functions f with $\mu(\{x : f(x) \neq 0\}) < \infty$ are dense in L^p . For $p = \infty$ we can drop the condition on the measure of the support.

Definition. For $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$, the *convolution* $f * g$ is defined by

$$f * g(x) = \int_{\mathbb{R}^n} f(y)g(x-y)dy,$$

provided the integral exists. Note that $f * g = g * f$, convolution is associative, and $\mu(f * g) = \mu(f)\mu(g)$.

Theorem. $C_c^\infty(\mathbb{R}^n)$ is dense in $L^p(\mathbb{R}^n)$ for $p \in [1, \infty)$.

Remark. This theorem is false for $p = \infty$.

Notation: a multiindex is $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Set $|\alpha| = \alpha_1 + \dots + \alpha_n$; $\alpha! = \alpha_1! \dots \alpha_n!$; $X^\alpha = X_1^{\alpha_1} \dots X_n^{\alpha_n}$ for $X \in \mathbb{R}^n$; $\nabla^\alpha f = D^\alpha f = \frac{\partial^\alpha f}{\partial x^\alpha} = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}}$.

Definition. We say $f \in L_{\text{loc}}^p(\mathbb{R}^n)$ if $f \mathbb{1}_K \in L^p(\mathbb{R}^n)$ for any $K \subseteq \mathbb{R}^n$ compact.

Proposition. Let $f \in L_{\text{loc}}^1(\mathbb{R}^n)$, $g \in C_c^k(\mathbb{R}^n)$, some $k \geq 0$. Then $f * g \in C^k(\mathbb{R}^n)$ and $\nabla^\alpha(f * g) = f * (\nabla^\alpha g)$ for all $|\alpha| \leq k$.

Proof. First we check for $k = 0$. Set $T_z f(x) = f(x - z)$, $z \in \mathbb{R}^n$. Then $T_z(f * g) = f * (T_z g)$. Also $T_z g(x) \rightarrow g(x)$ for all x as $z \rightarrow 0$ (continuity of g). Furthermore $|T_z g(x)| \leq \|g\|_{L^\infty} \mathbb{1}_{B_R(0)}(x)$ if $|x| + 1 \leq R$, $|z| < 1$ (we can just take R large enough so it holds everywhere since g has compact support). Then $|f(y)T_z g(x - y)| \leq C|f(y)|\mathbb{1}_{B_R(0)}(x - y)$, for $C := \|g\|_{L^\infty}$.

Since $f \in L_{\text{loc}}^1(\mathbb{R}^n)$, $|f(y)|\mathbb{1}_{B_R(0)}(x - y)$ is integrable in y , so by the dominated convergence theorem,

$$T_z(f * g) = (f * T_z g)(x) = \int_{\mathbb{R}^n} f(y)T_z g(x - y)dy \xrightarrow{z \rightarrow 0} \int_{\mathbb{R}^n} f(y)g(x - y)dy = (f * g)(x).$$

And so $f * g \in C^0$. Now let $k = 1$. Let $\nabla_i^h g(x) = \frac{g(x + h e_i) - g(x)}{h}$, where e_i is the i th unit vector. Then $\nabla_i^h g(x) \rightarrow \nabla_i g(x)$ as $h \rightarrow 0$.

By the mean value theorem, there exists $t \in [-h, h]$ such that

$$\nabla_i^h g(x) = \nabla_i g(x + t e_i) \Rightarrow |\nabla_i^h g(x)| \leq \|\nabla_i g\|_{L^\infty} \mathbb{1}_{B_R(0)}(x).$$

Again by the dominated convergence theorem, $\nabla_i^h(f * g) = f * (\nabla_i^h g) \rightarrow f * \nabla_i g$. Thus $f * g \in C^1$. The case $k > 1$ is similar, with induction. \square

Proposition (Minkowski's integral inequality). Let $p \in [1, \infty)$ and $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ Borel. Then

$$\left[\int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} f(x, y) dx \right|^p dy \right]^{1/p} \leq \int_{\mathbb{R}^n} \left[\int_{\mathbb{R}^n} |f(x, y)|^p dy \right]^{1/p} dx.$$

Proof. Example sheet 1. \square

Proposition. Let $p \in [1, \infty)$, $g \in L^p(\mathbb{R}^n)$. Then

$$\|T_z g - g\|_{L^p} \rightarrow 0 \text{ as } |z| \rightarrow 0.$$

Remark. This is not true for $p = \infty$. Let $\theta(x) = \mathbb{1}_{x \geq 0}$. Then $\|T_z \theta - \theta\|_{L^\infty} = 1$ if $z \neq 0$.

Proof. Consider first $g = \mathbb{1}_R$, R a rectangle. Then the result is clear. Hence it also follows for any finite union of rectangles. If B is a Borel set, $|B| < \infty$, then for every $\varepsilon > 0$, there exists a finite union of rectangles R such that

$$\|\mathbb{1}_B - \mathbb{1}_R\|_{L^p} = |B \triangle R|^{1/p} < \varepsilon.$$

Hence

$$\|T_z \mathbb{1}_B - \mathbb{1}_B\|_{L^p} \leq \underbrace{\|T_z \mathbb{1}_B - T_z \mathbb{1}_R\|_{L^p}}_{=\|\mathbb{1}_B - \mathbb{1}_R\|_{L^p} < \varepsilon} + \underbrace{\|T_z \mathbb{1}_R - \mathbb{1}_R\|_{L^p}}_{< \varepsilon \text{ for } |z| \text{ small}} + \underbrace{\|\mathbb{1}_R - \mathbb{1}_B\|_{L^p}}_{< \varepsilon}.$$

Thus the result holds for $g = \mathbb{1}_B$, $B \in \mathcal{B}(\mathbb{R}^n)$. Thus the result holds for simple functions g . Finally, for any $g \in L^p$, there is a \tilde{g} simple such that $\|g - \tilde{g}\|_{L^p} < \varepsilon$. Then

$$\|T_z g - g\|_{L^p} \leq \underbrace{\|T_z g - T_z \tilde{g}\|_{L^p}}_{=\|g - \tilde{g}\|_{L^p} < \varepsilon} + \underbrace{\|T_z \tilde{g} - \tilde{g}\|_{L^p}}_{< \varepsilon \text{ for } |z| \text{ small}} + \underbrace{\|g - \tilde{g}\|_{L^p}}_{< \varepsilon}.$$

□