

Overview

- Likelihood principle (11 lectures)
- Bayesian inference (2 lectures)
- Decision theory (3 lectures)
- Multivariate analysis (2 lectures)
- Nonparametric inference & Monte Carlo techniques (6 lectures)

Books:

- Theory of point estimation - Lehmann & Casella
- “Asymptotic Statistics” - van der Vaart
- “Statistical Inference” - Casella & Berger
- “Intro to Multivariate Statistical Analysis” - Anderson

Introduction

Goal: Make inference about unknown probability distributions based on access to random samples.

Consider a real valued random variable X on a probability space Ω with distribution function

$$F(t) = \mathbb{P}(\omega \in \Omega : X(\omega) \leq t) \quad \forall t \in \mathbb{R}$$

When X is discrete, $F(t) = \sum_{x \leq t} f(x)$, where f is the pmf of X .

When X is continuous, $F(t) = \int_{-\infty}^t f(s)ds$, where f is the pdf of X .

For all the results in this course, we assume either pdf or pmf exists.

Often, the distribution of X is parameterised by an unknown value θ . The goal is to infer something about θ based on (iid) samples X_1, \dots, X_n .

Definition. A *statistical model* for a sample from X is any family of probability distributions $\{P_\theta : \theta \in \Theta\}$ for the law of X . When P_θ has a pmf (pdf) $f(\cdot, \theta)$, this is also written as $\{f(\cdot, \theta) : \theta \in \Theta\}$. The index set Θ is the *parameter space*.

Example.

- (i) $\mathcal{N}(\theta, 1); \theta \in \Theta = \mathbb{R}$.
- (ii) $\mathcal{N}(\mu, \sigma^2), \theta = (\mu, \sigma^2) \in \Theta = \mathbb{R} \times (0, \infty)$.
- (iii) $\text{Exp}(\theta); \theta \in \Theta = (0, \infty)$.

(iv) $\mathcal{N}(\theta, 1)$; $\theta \in \Theta = [-1, 1]$.

Remark: for a variable X with distribution P , the model $\{P_\theta : \theta \in \Theta\}$ is *correctly specified* if there exists $\theta \in \Theta$ such that $P = P_\theta$. For instance, if $X \sim \mathcal{N}(2, 1)$, the model in (i) is correctly specified, but the model in (iv) is not.

In the case of a correctly specified model, we often use θ_0 to denote the “true value” of the parameter. We also say $\{X_1, \dots, X_n\}$ are iid from a model $\{P_\theta : \theta \in \Theta\}$ in the case of a correctly specified model.

Statistical goals:

- Estimation: construct $\hat{\theta} = \hat{\theta}(X_1, \dots, X_n)$ such that $\hat{\theta}$ is close to θ_0 when $X_i \sim P_{\theta_0}$.
- Hypothesis testing: determine whether the null hypothesis $H_0 : \theta = \theta_0$ or the alternative hypothesis $H_1 : \theta \neq \theta_0$ is true, using a test $\psi_n = \psi(X_1, \dots, X_n)$ such that $\psi_n = 0$ when H_0 is true and $\psi_n = 1$ when H_1 is true, with high probability.
- Inference: find confidence intervals (confidence sets) $\mathcal{C}_n = \mathcal{C}(X_1, \dots, X_n)$ such that for some $0 < \alpha < 1$ we have $\mathbb{P}_\theta(\theta \in \mathcal{C}_n) \geq 1 - \alpha$, for all $\theta \in \Theta$, where α is the significance level.

1 The Likelihood Principle

Suppose X_1, \dots, X_n are iid from a Poisson model $\{\text{Poi}(\theta) : \theta \geq 0\}$ with numerical values $X_i = x_i$, for all $1 \leq i \leq n$. The joint distribution of the sample is

$$f(x_1, \dots, x_n; \theta) = \mathbb{P}_\theta(X_1 = x_1, \dots, X_n = x_n) = \prod_{i=1}^n \left(e^{-\theta} \frac{\theta^{x_i}}{x_i!} \right) = e^{-n\theta} \prod_{i=1}^n \frac{\theta^{x_i}}{x_i!} = L_n(\theta)$$

We can think of $L_n(\theta)$ as a random function from Θ to \mathbb{R} , where the randomness comes from $\{X_i\}_{i=1}^n$. This is the probability of occurrence of the observed sample $(X_1 = x_1, \dots, X_n = x_n)$, as a function of the unknown parameter θ .

The idea of the likelihood principle is to find θ which maximises $L_n(\theta)$, or equivalently $l_n(\theta) = \log L_n(\theta)$. In the example, we have

$$l_n(\theta) = -n\theta + \log(\theta) \sum_{i=1}^n x_i - \sum_{i=1}^n \log(x_i!)$$

Setting $l'_n(\theta) = 0$ gives

$$-n + \frac{1}{\theta} \sum_{i=1}^n x_i = 0$$

and the solution is $\hat{\theta}_{\text{mle}} = \frac{1}{n} \sum_{i=1}^n x_i$, which is the sample mean. One can also check that $l_n''(\theta) < 0$ for all $\theta > 0$. When all X_i 's are 0, one can check that maximising $l_n(\theta)$ is equivalent to maximising $-n\theta$, so $\hat{\theta}_{\text{mle}} = 0$ in this case.

Maximum likelihood estimator

Suppose $\{f(\cdot, \theta) : \theta \in \Theta\}$ is a statistical model of pdfs/pmfs for the distribution of a random variable X , and X_1, \dots, X_n are iid copies of X .

Define the *likelihood function*

$$L_n(\theta) = \prod_{i=1}^n f(x_i, \theta)$$

the *log likelihood function*

$$l_n(\theta) = \log L_n(\theta) = \sum_{i=1}^n \log f(x_i, \theta)$$

and the *normalised log likelihood function*

$$\bar{l}_n(\theta) = \frac{1}{n} l_n(\theta) = \frac{1}{n} \sum_{i=1}^n \log f(x_i, \theta)$$

Definition. The *maximum likelihood estimator* is any element $\hat{\theta} = \hat{\theta}_{\text{mle}} = \hat{\theta}_{\text{mle}}(X_1, \dots, X_n) \in \Theta$ for which $L_n(\hat{\theta}) = \max_{\theta \in \Theta} L_n(\theta)$.

Remark: the definition of MLE can be generalised to non-iid data, provided a joint pdf/pmf of (X_1, \dots, X_n) can be specified.

Example.

- (i) For $X_i \sim \text{Poi}(\theta)$, $\theta \geq 0$, we calculated $\hat{\theta}_{\text{mle}} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}_n$.
- (ii) For $X_i \sim \mathcal{N}(\mu, \sigma^2)$, $\theta = (\mu, \sigma^2) \in \mathbb{R} \times (0, \infty)$, we have $\hat{\mu}_{\text{mle}} = \bar{X}_n$ and $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2$ (see Example sheet).
- (iii) In the Gaussian linear model $Y = X\theta + \varepsilon$, with a known $X \in \mathbb{R}^{n \times p}$, unknown $\theta \in \mathbb{R}^p$, and $\varepsilon \sim \mathcal{N}(0, I_n)$, the observations (Y_1, \dots, Y_n) are not iid, but a joint distribution $f(Y_1, \dots, Y_n; \theta)$ can still be specified. The MLE is the least squares estimator (see Example sheet).

Definition. For $\Theta \subseteq \mathbb{R}^p$ and l_n differentiable at θ , the *score function* S_n is

$$S_n(\theta) = \begin{pmatrix} \frac{\partial}{\partial \theta_1} l_n(\theta) \\ \vdots \\ \frac{\partial}{\partial \theta_p} l_n(\theta) \end{pmatrix}$$

Solving for a root of $S_n(\theta)$ is a common heuristic for maximising $l_n(\theta)$. In many cases, it is a necessary and sufficient condition.

Note: derivatives are taken with respect to θ , not the x_i 's.

Information geometry

Recall that if X is a random variable with distribution P_θ on some space $\mathcal{X} \subseteq \mathbb{R}^d$, and $g : \mathcal{X} \rightarrow \mathbb{R}$ is a function, then

$$E_\theta[g(X)] = \int_{\mathcal{X}} g(x) dP_\theta(x) = \int_{\mathcal{X}} g(x) f(x, \theta) dx$$

if X has a pdf $f(x, \theta)$, and

$$\mathbb{E}_\theta[g(X)] = \sum_{x \in \mathcal{X}} g(x) f(x, \theta)$$

if X has a pmf $f(x, \theta)$

Theorem 1.1. Consider a model $\{f(\cdot, \theta) : \theta \in \Theta\}$, where $f(\cdot, \theta)$ is a pdf/pmf and $f(x, \theta) > 0$ for all x, θ . Also suppose the model is correctly specified, with θ_0 equal to the true parameter, and $\mathbb{E}_{\theta_0}[|\log(f(X, \theta))|] < \infty$ for all $\theta \in \Theta$. Then the function defined by $l(\theta) = \mathbb{E}_{\theta_0}[\log(f(X, \theta))]$ is maximised at θ_0 .

Proof. Consider the case when X has a pdf (discrete case is analogous). For all $\theta \in \Theta$, we have

$$\begin{aligned} l(\theta) - l(\theta_0) &= \mathbb{E}_{\theta_0}[\log(f(X, \theta))] - \mathbb{E}_{\theta_0}[\log(f(X, \theta_0))] \\ &= \mathbb{E}_{\theta_0} \left[\log \left(\frac{f(X, \theta)}{f(X, \theta_0)} \right) \right] \end{aligned}$$

Jensen's inequality: $\mathbb{E}[\varphi(Z)] \leq \varphi(\mathbb{E}[Z])$ for any random variable Z and concave function φ .

Since \log is concave,

$$\begin{aligned} l(\theta) - l(\theta_0) &\leq \log \left(\mathbb{E}_{\theta_0} \left[\frac{f(X, \theta)}{f(X, \theta_0)} \right] \right) \\ &= \log \left(\int_{\mathcal{X}} \frac{f(x, \theta)}{f(x, \theta_0)} f(x, \theta_0) dx \right) = \log 1 = 0 \end{aligned} \quad (*)$$

□

Remark: under the assumption of “strict identifiability of the model parameterisation”, i.e.,

$$f(\cdot, \theta) = f(\cdot, \theta') \iff \theta = \theta'$$

the inequality (*) is strict, since equality occurs in Jensen only when φ is linear or Z is constant.

Remark: the quantity $l(\theta_0) - l(\theta)$ computed above can be written as

$$\text{KL}(P_{\theta_0}, P_\theta) = \int_{\mathcal{X}} f(x, \theta_0) \log \left(\frac{f(x, \theta_0)}{f(x, \theta)} \right) dx$$

and is the Kullback-Leibler divergence in information theory. It is a “distance” between distributions. Maximising $l(\theta)$ is equivalent to minimising KL.

Fisher information

We consider the gradient and Hessian of the likelihood function.

Theorem 1.2. *For a parametric model $\{f(\cdot, \theta) : \theta \in \Theta\}$, “regular enough” so integration and differentiation can be interchanged, we have $\mathbb{E}_\theta[\nabla_\theta \log(f(X, \theta))] = 0$ for all $\theta \in \text{interior}(\Theta)$.*

Proof. We write the expectation

$$\begin{aligned}\mathbb{E}_\theta[\nabla_\theta \log(f(X, \theta))] &= \int_{\mathcal{X}} (\nabla_\theta \log f(x, \theta)) f(x, \theta) dx \\ &= \int_{\mathcal{X}} \frac{\nabla_\theta f(x, \theta)}{f(x, \theta)} f(x, \theta) dx \\ &= \nabla_\theta \left(\int_{\mathcal{X}} f(x, \theta) dx \right) = \nabla_\theta(1) = 0\end{aligned}$$

□

Remark: in particular, when $\theta_0 \in \text{interior}(\Theta)$, then $\mathbb{E}_{\theta_0}[\nabla_\theta \log(f(X, \theta))] = 0$.

Definition. For a parameter space $\Theta \subseteq \mathbb{R}^p$, the *Fisher information* matrix is defined by

$$I(\theta) = \mathbb{E}_\theta \left[(\nabla_\theta \log f(X, \theta)) (\nabla_\theta \log f(X, \theta))^T \right], \quad \forall \theta \in \text{interior}(\Theta)$$

in other words,

$$I_{ij}(\theta) = \mathbb{E}_\theta \left[\frac{\partial}{\partial \theta_i} \log f(X, \theta) \frac{\partial}{\partial \theta_j} \log f(X, \theta) \right]$$

Remark: in 1 dimension, we have

$$I(\theta) = \mathbb{E}_\theta \left[\left(\frac{d}{d\theta} \log f(X, \theta) \right)^2 \right] = \text{Var}_\theta \left[\frac{d}{d\theta} \log f(X, \theta) \right]$$

Thus I_{θ_0} describes random variations of $S_n(\theta_0)$ about its mean. This in turn will help quantify the precision of $\hat{\theta}$, a zero of $S_n(\hat{\theta}) = 0$, about θ_0 .

Theorem 1.3. *Under the same regularity assumptions as the previous theorem*

$$I(\theta) = -\mathbb{E}_\theta \left[\nabla_\theta^2 \log(f(X, \theta)) \right], \quad \forall \theta \in \text{interior}(\Theta)$$

i.e.,

$$I_{ij}(\theta) = -\mathbb{E}_\theta \left[\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log f(X, \theta) \right]$$

Proof. We write

$$\nabla_{\theta}^2 \log f(X, \theta) = \nabla_{\theta} \left(\frac{\nabla_{\theta} f(X, \theta)}{f(X, \theta)} \right) = \frac{\nabla_{\theta}^2 f(X, \theta)}{f(X, \theta)} - \frac{\nabla_{\theta} f(X, \theta) \nabla_{\theta} f(X, \theta)^T}{f(X, \theta)^2}$$

note that

$$\mathbb{E} \left[\frac{\nabla_{\theta}^2 f(X, \theta)}{f(X, \theta)} \right] = \int_{\mathcal{X}} \nabla_{\theta}^2 f(X, \theta) dx = \nabla_{\theta}^2 \int_{\mathcal{X}} f(X, \theta) dx = 0$$

Therefore

$$\begin{aligned} -\mathbb{E}_{\theta} [\nabla_{\theta}^2 \log f(X, \theta)] &= \mathbb{E}_{\theta} \left[\frac{\nabla_{\theta} f(X, \theta) \nabla_{\theta} f(X, \theta)^T}{f^2(X, \theta)} \right] \\ &= \mathbb{E} \left[\frac{\nabla_{\theta} f(X, \theta)}{f(X, \theta)} \left(\frac{\nabla_{\theta} f(X, \theta)}{f(X, \theta)} \right)^T \right] \\ &= \mathbb{E}_{\theta} [(\nabla_{\theta} \log f(X, \theta))(\nabla_{\theta} \log f(X, \theta))^T] \\ &= I(\theta) \end{aligned}$$

□

Remark: continuing the previous remark, in 1 dimension

$$\text{Var}_{\theta} \left[\frac{d}{d\theta} \log f(X, \theta) \right] = I(\theta) = -\mathbb{E}_{\theta} \left[\frac{d^2}{d\theta^2} \log f(X, \theta) \right]$$

this relates the variance of the score function and the curvature of l , both of which are relevant to describing the quality of the MLE $\hat{\theta}$ as an approximation to θ_0 .