

# 1 Kernel Machines

Consider a linear model

$$Y_i = x_i^T \beta^0 + \varepsilon_i, \quad i = 1, \dots, n, \quad x_i \in \mathbb{R}^p \text{ fixed}$$

where  $\mathbb{E}\varepsilon = 0$ ,  $\text{Var}(\varepsilon) = \sigma^2 I_n$ . We have

$$\begin{aligned} \hat{\beta}^{\text{ols}} &= \underset{\beta \in \mathbb{R}^p}{\text{argmin}} \sum_{i=1}^n (Y_i - x_i^T \beta)^2 \\ &= \underset{\beta \in \mathbb{R}^p}{\text{argmin}} \|Y - X\beta\|^2 \\ &= (X^T X)^{-1} X^T Y. \end{aligned}$$

Classical theory:

- $\hat{\beta}^{\text{ols}}$  unbiased,

$$\text{Var}(\hat{\beta}^{\text{ols}}) = \sigma^2 (X^T X)^{-1} = i^{-1}(\beta^0)$$

Where  $i$  is the Fisher information.

- Cramér-Rao lower bound: if an estimator  $\tilde{\beta}$  is unbiased then

$$\text{Var}(\tilde{\beta}) - i^{-1}(\beta^0) \underset{\text{positive semi-definite}}{\geq} 0.$$

- If  $\varepsilon \sim \mathcal{N}(0, \sigma^2 I_n)$ , then  $\hat{\beta}^{\text{ols}}$  is the MLE of  $\beta^0$ . Furthermore  $\sqrt{n}(\hat{\beta}^{\text{ols}} - \beta^0) \sim \mathcal{N}(0, n\sigma^2(X^T X)^{-1})$ . From this we can derive confidence intervals, hypothesis test, etc.

In a general model with parameter  $\theta \in \mathbb{R}^p$ ,  $n$  independent observations, under regularity, we have asymptotic normality, i.e.  $\sqrt{n}(\hat{\theta}^{\text{MLE}} - \theta^0) \xrightarrow{d} \mathcal{N}(0, I^{-1}(\theta^0))$  (with  $p$  fixed).

Question: what happens when  $p$  is large relative to  $n$ ?

- If  $p > n$ ,  $\hat{\beta}^{\text{ols}}$  is not even defined.
- If  $p \approx n$ ,  $\text{Var}(\hat{\beta}^{\text{ols}})$  explodes since  $X^T X$  is near singular.
- More generally, if  $p, n \rightarrow \infty$  then asymptotic normality can break down.

Recall the bias-variance decomposition:

$$\begin{aligned} \text{mse}(\tilde{\beta}) &= \mathbb{E}_{\beta^0, \sigma^2} [(\tilde{\beta} - \beta^0)(\tilde{\beta} - \beta^0)] \\ &= \mathbb{E}_{\beta^0, \sigma^2} \left\| \tilde{\beta} - \mathbb{E}\tilde{\beta} + \mathbb{E}\tilde{\beta} - \beta^0 \right\|^2 \\ &= \text{Var}(\tilde{\beta}) + \left\| \mathbb{E}(\tilde{\beta}) - \beta^0 \right\|^2. \end{aligned}$$

We introduce bias to reduce the variance.

## 1.1 Ridge regression

Define

$$(\hat{\mu}_\lambda^R, \hat{\beta}_\lambda^R) = \operatorname{argmin}_{(\mu, \beta) \in \mathbb{R} \times \mathbb{R}^p} \left[ \|Y - \mu \mathbf{1} - X\beta\|^2 + \underbrace{\lambda \|\beta\|^2}_{\text{penalty for large } \beta} \right].$$

$\lambda$  is called a *regularisation* or *tuning* parameter. We shall assume the columns of  $X$  have been standardised (mean 0, variance 1).

After standardisation, we can show that

$$\hat{\mu}_\lambda^R = \frac{1}{n} \sum_{i=1}^n Y_i = \bar{Y}.$$

Hence, if we replace  $Y$  with  $Y - \mathbf{1}\bar{Y}$  we can write

$$\begin{aligned} \hat{\beta}_\lambda^R &= \operatorname{argmin}_{\beta \in \mathbb{R}^p} [\|Y - X\beta\|^2 + \lambda \|\beta\|^2] \\ &= \underbrace{(X^T X + \lambda I_p)^{-1}}_{\text{always invertible}} X^T Y. \end{aligned}$$

**Theorem 1.1.** For  $\lambda > 0$  sufficiently small,

$$\mathbb{E}\|\hat{\beta}^{\text{ols}} - \beta^0\|^2 - \mathbb{E}\|\hat{\beta}_\lambda^R - \beta^0\|^2 > 0. \quad (*)$$

*Proof.* We have

$$Y = X\beta^0 + \varepsilon.$$

The bias of  $\hat{\beta}_\lambda^R$  is

$$\begin{aligned} \mathbb{E}(\hat{\beta}_\lambda^R - \beta^0) &= (X^T X + \lambda I)^{-1} X^T X \beta^0 - \beta^0 \\ &= (X^T X + \lambda I)^{-1} (X^T X + \lambda I - \lambda I) \beta^0 - \beta^0 \\ &= -\lambda (X^T X + \lambda I)^{-1} \beta^0. \end{aligned}$$

While we have variance

$$\begin{aligned} \operatorname{Var}(\hat{\beta}_\lambda^R) &= \mathbb{E} \|(X^T X + \lambda I)^{-1} X^T \varepsilon\|^2 \\ &= \sigma^2 [(X^T X + \lambda I)^{-1} X^T X (X^T X + \lambda I)^{-1}]. \end{aligned}$$

Then  $(*)$  becomes

$$\begin{aligned} &\mathbb{E}\|\hat{\beta}^{\text{ols}} - \beta^0\|^2 - \mathbb{E}\|\hat{\beta}_\lambda^R - \beta^0\|^2 \\ &= \sigma^2 (X^T X)^{-1} - \sigma^2 (X^T X + \lambda I) X^T X (X^T X + \lambda I)^{-1} \\ &\quad - \lambda^2 (X^T X + \lambda I)^{-1} \beta^0 (\beta^0)^T (X^T X + \lambda I)^{-1} \\ &= \vdots \quad \quad \quad (\text{use SVD } X = UDU^T) \\ &= \lambda (X^T X + \lambda I)^{-1} [\sigma^2 \{2I_p + \lambda (X^T X)^{-1}\} - \lambda \beta^0 (\beta^0)^T] (X^T X + \lambda I)^{-1}. \end{aligned}$$

We want to show this is positive definite. This is equivalent to

$$\begin{aligned}\sigma^2 [2I + \lambda(X^T X)^{-1}] - \lambda\beta^0(\beta^0)^T &> 0 \\ \iff 2\sigma^2 I - \lambda\beta^0(\beta^0)^T &> 0 \\ \iff 2\sigma^2 \|z\|^2 - \lambda(z^T \beta^0)^2 &> 0 \quad \forall z \in \mathbb{R}^p.\end{aligned}\tag{\dagger}$$

We also have  $(z^T \beta^0)^2 \leq \|z\|^2 \|\beta^0\|^2$  by Cauchy-Schwartz. Hence  $(\dagger)$  holds for all  $\lambda < \frac{2\sigma^2}{\|\beta^0\|^2}$ .  $\square$