

Prevalence of pathological tooth wear in patients with chronic alcoholism

N. D. Robb, BDS, FDS

The Dental School, Framlington Place, Newcastle upon Tyne NE2 4BW

B. G. N. Smith, PhD, MSc, BDS, FDS

Department of Conservative Dentistry, UMDS, Guy's Hospital, London SE1 9RT

Although chronic alcoholism is a very common condition, with potentially harmful consequences for the sufferer, there has been little emphasis in the literature on its effects on the teeth. This study of 37 alcoholic patients showed that their teeth had significantly more wear than age and sex matched controls. The tooth wear was most marked in males, and those whose alcohol consumption was continuous rather than in the form of episodic binges. The wear appeared to be erosive in nature, and in 40% of the sample it affected the palatal surfaces of the upper anterior teeth. It is therefore suggested that general dental practitioners should bear in mind the possibility of chronic alcoholism in cases of unexplained dental erosion.

Chronic alcoholism is a serious condition, with many potentially life-threatening complications.¹ These include liver cirrhosis, neurological disorders, such as Wernicke-Korsakoff syndrome and peripheral neuropathy, interactions with other drugs, malnutrition, aggravation of pre-existing cardiovascular disease and mental disorders, such as alcohol psychoses and affective disorders.¹⁻³ Alcohol causes severe irritation of the mucosa of the stomach and duodenum, which is worse in alcoholics than non-alcoholics.⁴

Chronic alcoholism is thought to affect about 10% of the adult population.^{2,5} The cost implications, both in terms of the medical management and the time lost from employment are large, having been estimated at £43 billion in the USA in 1975.⁶

Changes in the oral tissues can help in the diagnosis of many disorders.⁷ This is also true of alcoholism, where enlargement of the parotid glands⁸ and attrition (tooth wear caused by physical contact of one tooth with another)⁹ have been reported. Patients with a history of excessive alcohol consumption have been shown to have pathological levels of tooth wear which looks more like erosion (the loss of enamel and dentine due to chemical attack, other than acids produced by bacteria) rather than attrition.¹⁰ The current study was undertaken to assess the prevalence of pathological tooth wear in a population of chronic alcoholic patients.

Materials and methods

In-patients admitted for the treatment of chronic alcoholism were seen at the Maudsley Hospital, London and Warlingham Park Hospital, Surrey. The psychiatric diagnosis had already been established by the physicians in charge of the patients' care. Criteria for inclusion in the study were a willingness to participate and a minimum of 12 scoreable permanent teeth.

The patients were all seen by the same examiner, and detailed histories of the factors known to be involved in the aetiology of tooth wear were taken under the headings shown in Table I. A 'Tooth Wear Index'¹¹ was recorded, scoring

Table I Factors in the history of tooth wear

Diet
Digestive disorders
Oral hygiene practices
Habits (eg bruxism, nail biting)
Past medical and dental history
Present medical and dental history

cervical, buccal, occlusal, and lingual surfaces of all standing permanent teeth, giving a maximum of 128 surfaces per dentition. Missing, restored and unscoreable surfaces were also recorded, as was Angles Classification, quality of posterior support, and the ease of movement in lateral and protrusive excursions.

The raw data were analysed using a specially written computer programme to produce pathological tooth wear levels and also mean pathological wear profiles for different groups of sites.¹¹

A control group of healthy individuals were used for comparison. The control group were selected from a separate study of 1000 dental attenders from 102 dental practices in the South East of England. The control for each alcoholic subject was selected on a best match basis for age, sex, social class, according to occupation and number of standing teeth. Individuals with a history of alcoholism or prolonged excessive alcohol consumption were excluded prior to the matching process. Similar histories and examinations were carried out by the same examiner, and the data handled in the same way. This method was used in a previous study of individuals suffering from anorexia and bulimia nervosa.¹²

The total number of sites scored and the number rated as pathological were calculated. The differences between the groups were analysed using a test to determine the significance of the differences between proportions.

Results

The mean ages and number of sites scored per patient in the alcoholic and control groups are shown in Table II. The groups are comparable by these criteria.

Br Dent J 1990; 169: 267
© British Dental Journal 1990
Accepted for publication 22.5.90

Table II A comparison of the experimental and control groups

	Mean age in years (range)	Mean number of surfaces scored (range)	Mean number pathologically worn (range)	Total % pathol wear	Number of individuals with upper palatal pathol wear (%)	Number of individuals with no pathological wear (%)
Alcoholic	38.4 (23-65)	91.7 (43-128)	9.01 (0-51)	9.85	15 (40.5)	3 (8.1)
Control	39.6 (25-63)	98.1 (56-128)	1.8 (0-6)	1.82 ^a	3 ^b (8.1)	13 ^b (35.1)

^a $P < 0.001$ ^b $P < 0.01$ **Table III** Number of subjects divided by the percentage of pathologically worn sites

Percentage of sites affected by pathological tooth wear	Control group	Alcoholic group
0	13	3
0.1-5.0	21	16
5.1-10.0	3	5
10.1-15.0	0	4
15.1-20.0	0	1
20.1-25.0	0	3
>30	0	5
	37	37

Table IV Difference in wear between males and females

	Number of subjects	Total sites scored	Number of sites pathologically worn	% Pathologically worn
Alcoholic male	25	2228	248	11.11
female	12	1164	86	7.39 ^a
Control male	25	2426	51	2.10
female	12	1205	15	1.30 ^b

^a $P < 0.001$ ^bN/S**Table V** Difference in wear between continuous and binge drinkers, and those with and without morning retching and vomiting

	Number of subjects	Total sites recorded	Number of sites pathologically worn	% Pathologically worn
Continuous drinkers	19	1719	195	11.34 ^c
Binge	16 (a)	1524	95	6.23
Vomiters	22	2133	149	6.98
Non-vomiters	12 (b)	880	52	5.91 ^d

^aTwo subjects not recorded^bThree subjects not recorded^c $P < 0.001$ ^dN/S

The prevalence of pathological tooth wear

Table II shows the differences between the experimental and the control groups. The alcoholic group had significantly more pathologically worn sites than the control group ($P < 0.001$). Table II also shows the number of patients with pathological wear of the palatal surfaces of the upper anterior teeth, one of the features of a 'regurgitation erosion' pattern of wear,¹³ and the number of individuals with no pathological tooth wear. The pathological wear was not spread evenly between the subjects in the alcoholic group, as can be seen from Table III. It was not possible, from the histories given, to explain why some patients were affected more than others.

When the experimental group was divided by sex (Table IV) the males had significantly more pathologically worn sites than the females. This difference was not found to be significant in the control group.

The sample of alcoholics was also divided into those who drank regularly and those who binge drank, ie drinking large amounts of alcohol in phases lasting up to 1-2 weeks and then have dry periods in between, and those who complained of nausea and vomiting in the morning and those who did not (Table V).

The typical clinical appearance of the tooth wear in this group has already been described,¹⁰ and a case is shown (fig. 1).

Discussion

The majority of these patients showed a predominance of smooth surface wear, suggesting that erosion is the most likely cause, although the exact nature of this process has yet to be determined. It is, however, possible that it was caused either

**Fig. 1** Palatal view showing typical erosive wear on the unrestored palatal surfaces of 3.2.1 (13, 12, 11).

eam,
200n
mes
5 day

ists' Terms
ee require
ists to writ
lovir" whe
ribing Zov
the NHS

Zoviran

Zoviran

Zoviran

Zoviran

Zoviran

Power
of o
Tha

Power
of o
Tha

Power
of o
Tha

THE W
OUND
LEADER IN A

the acid in the beverages or, more likely, from a subclinical regurgitation due to the chronic gastritis which is known to be produced by the ingestion of large amounts of alcohol. The latter is more probable, as 40% of the alcoholic patients had a large amount of wear palatally in the upper incisor region, a pattern which is typical of patients who are known to vomit and regurgitate regularly.

Chronic alcoholism is a condition which still carries a large social stigma, and most patients are secretive about it. It is therefore possible that the general dental practitioner may be the first person to suspect that an individual is suffering from chronic alcoholism on the evidence of unexplained dental erosion.

Conclusions

The experimental group showed significantly more pathological tooth wear than the controls, and there was significantly more pathological wear in continuous drinkers than in binge drinkers, and in male alcoholics than in female alcoholics.

There was no difference between those alcoholics who reported feelings of nausea and vomiting in the morning and those who did not, although the clinical appearance of the wear was the same as regurgitation erosion. In view of the high prevalence of alcohol abuse in the general population, the possibility of chronic alcoholism should be borne in mind when seeing patients with tooth wear problems, particularly when it affects the palatal surfaces of the upper anterior teeth. Further work is needed to establish why some individuals are affected to a much greater extent than others.

Acknowledgements

We would like to thank Dr J. J. Gayford and the staff at Warlingham Park Hospital, Professor G. Edwards and his staff at the Maudsley Hospital, and Mrs E. Geidrys-Leeper for their help in this study. Financial support was given by the Special Trustees at Guy's Hospital and the Endowment fund at St. George's Hospital.

References

- 1 Echaridt M J, Hartford T C, Kaelber C T, Parker E S, Rosenthal L, Ryback R S, Salmoiraght G S, Vanderveen E, Warren K R. Health hazards associated with alcohol consumption. *J Am Med Assoc* 1981; **246**: 648-666.
- 2 Christen A G. Dentistry and the alcoholic patient. *Dent Clin N Am* 1983; **27**: 341-361.
- 3 Schuckit M A. Overview of alcoholism. *J Am Dent Assoc* 1979; **99**: 489-493.
- 4 Gottfried E G, Karsten M A, Leiber C S. Alcohol induced gastric and duodenal lesions in man. *Am J Gastroenterol* 1978; **70**: 587-592.
- 5 Brickley M, Shepherd J P. Alcohol intoxication and dependence in adult dental out-patients. *J Dent Res* 1989; **68**: 591 (Abstr no. 261).
- 6 Berry R E Jnr, Boland J P. *The economic cost of alcohol abuse*. New York: Free Press, 1977.
- 7 Cawson R A. *Essentials of dental surgery and pathology*. 4th ed. pp358-389. Edinburgh: Churchill Livingstone, 1981.
- 8 Mandel L, Baurmash H. Parotid enlargement due to alcoholism. *J Am Dent Assoc* 1971; **82**: 369.
- 9 King W H, Tucker K M. Dental problems of alcoholic and non alcoholic psychiatric patients. *Q J Stud Alcohol* 1973; **34**: 1208-1211.
- 10 Smith B G N, Robb N D. Dental erosion in patients with chronic alcoholism. *J Dent* 1989; **17**: 219-221.
- 11 Smith B G N, Knight J K. An index for measuring the wear of teeth. *Br Dent J* 1984; **156**: 435-438.
- 12 Robb N D, Smith B G N, Geidrys-Leeper E. Prevalence of pathological tooth wear in patients with anorexia and bulimia nervosa. *J Dent Res* 1988; **67**: 661 (Abstr no. 175).
- 13 Smith B G N, Knight J K. A comparison of patterns of wear with aetiological factors. *Br Dent J* 1984; **157**: 16-19.

Guidelines for Authors

Manuscripts should be typewritten, double spaced, on one side only of A4 size paper, with large margins. Page 1 of the manuscript should give only: title of article, author's name, qualifications and address. The top copy and one other copy should be sent to the Editor, 64 Wimpole Street, London W1M 8AL. Authors should also keep a copy. All authors must sign their consent to publication.

Abstracts may be up to 200 words in length and should be able to stand alone.

Tables should be typed on separate pages, with one table per page. Each table should have a title.

Illustrations should be clearly identified on the back using a soft pencil or adhesive label. Legends should be typed on a separate sheet of paper. Lettering in illustrations should be put on the original. Figures, charts, graphs and drawings must be neatly drawn in indian ink, large enough to withstand reduction. All illustrations and tables should be referred to in the text. Computer-generated artwork is *not* suitable for publication, and should be redrawn in ink.

Illustrations in colour can only be reproduced by special arrangement.

Two copies of all illustrations should be provided.

SI units must be used throughout.

References should be in the Vancouver style, numbered in the order in which they appear (Naylor²⁻⁵ reported . . .). At the end of the article the full list of references should be given. Titles of books should be followed by the place of publication, the publisher, and the year. The author is responsible for accuracy of the reference list.

Manuscripts must be solely the work of the author/authors stated, must not have been previously published elsewhere and must not be under consideration by another periodical.

Ethics. Articles involving clinical research should, in general, have received ethical committee approval.

Acknowledgements. Group these in a paragraph at the end of the text. Where the research project was supported by industry, this should be acknowledged in the covering letter to the Editor on submission of the manuscript.

The Editor reserves the right to edit manuscripts for clarity of expression and to conform to acceptable style and the limits of the space available.

The entire text matter in the *British Dental Journal* is copyright and permission for reproduction must be obtained by application in writing to the Editor.