

An Inferentialist Account of Identity and Modality

Greg Restall



EXPRESS/PHILMATH SEMINAR · 16 FEBRUARY 2021

[https://consequently.org/presentation/2021/
an-inferentialist-account-of-identity-and-modality](https://consequently.org/presentation/2021/an-inferentialist-account-of-identity-and-modality)

My Aim

To show how *defining rules*

My Aim

To show how *defining rules* in a *hypersequent* setting

My Aim

To show how *defining rules* in a *hypersequent* setting can give a uniform proof-theoretic semantics of *identity*

My Aim

To show how *defining rules* in a *hypersequent* setting can give a uniform proof-theoretic semantics of *identity* and *modality*,

My Aim

To show how *defining rules* in a *hypersequent* setting can give a uniform proof-theoretic semantics of *identity* and *modality*, allowing — equally naturally — for

(1) modal operators for which identity statements are *necessary* (if true),

To show how *defining rules* in a *hypersequent* setting can give a uniform proof-theoretic semantics of *identity* and *modality*, allowing — equally naturally — for (1) modal operators for which identity statements are *necessary* (if true), and (2) modal operators for which identity statements can be *contingently* true.

My Plan

My Approach

Defining Identity

Defining Necessity

Combining Them

The Upshot

MY APPROACH

Normative Pragmatic, Semantically Anti-Realist and Dialogical

- ▶ Define concepts by way of rules for their *use*. (That's *normative pragmatism*.)

Normative Pragmatic, Semantically Anti-Realist and Dialogical

- ▶ Define concepts by way of rules for their *use*. (That's *normative pragmatism*.)
- ▶ Their semantics is not given in the first instance in terms of reference relations to objects. (That's *semantic anti-realism*.)

Normative Pragmatic, Semantically Anti-Realist and Dialogical

- ▶ Define concepts by way of rules for their *use*. (That's *normative pragmatism*.)
- ▶ Their semantics is not given in the first instance in terms of reference relations to objects. (That's *semantic anti-realism*.)
- ▶ The rules are grounded in our *communicative* and *dialogical practice*, involving assertion, denial and inference.



See Catarina Dutilh Novaes,
The Dialogical Roots of Deduction, 2020.

Defining Rules for Logical Concepts

\wedge Conjunction

$$\frac{X, A, B \succ Y}{X, A \wedge B \succ Y} \wedge Df$$

Asserting a conjunction has the same force as asserting its conjuncts.

Defining Rules for Logical Concepts

- Λ Conjunction
- Conditional

$$\frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df$$

Denying a material conditional has the same force as asserting its antecedent and denying its consequent.

Or equivalently, to *prove* a material conditional *suppose* its antecedent and *prove* its consequent.

Defining Rules for Logical Concepts

\wedge Conjunction

\rightarrow Conditional

\forall Universal Quantifier

$$\frac{X \succ A[n], Y}{X \succ \forall x A, Y} \forall Df$$

To prove that everything has some feature, prove that n has that feature, making no assumptions about n .

Or equivalently, denying that everything satisfies A has the same force as denying that n satisfies A for some fresh name ‘ n ’.

How are these *definitions*?

These defining rules show how to add the defined concept to a language that does not possess it . . .

How are these *definitions*?

These defining rules show how to add the defined concept to a language that does not possess it . . .

. . . not by finding a paraphrase in the original language, but by ‘making explicit’ something that was implicit.

How are these *definitions*?

These defining rules show how to add the defined concept to a language that does not possess it . . .

. . . not by finding a paraphrase in the original language, but by ‘making explicit’ something that was implicit.

In controlled circumstances, we can prove that concepts added by defining rules like these are *conservative* and *uniquely defined*, just like definitions by paraphrase.

Why Define?

In our everyday vocabulary, we make moves *like these* with our informal concepts ‘and’, ‘not’, ‘if’ and ‘all’.

Why Define?

In our everyday vocabulary, we make moves *like these* with our informal concepts ‘and’, ‘not’, ‘if’ and ‘all’.

The logician’s *sharply defined* concepts \wedge , \rightarrow , \forall , etc. are *simple, well-behaved, useful* and *can be followed*.

Why Define?

In our everyday vocabulary, we make moves *like these* with our informal concepts ‘and’, ‘not’, ‘if’ and ‘all’.

The logician’s *sharply defined* concepts \wedge , \rightarrow , \forall , etc. are *simple, well-behaved, useful* and *can be followed*.

Defining rules suffice for us to *coordinate* and to *communicate*, especially in those contexts where precision is valuable.

Why Define?

In our everyday vocabulary, we make moves *like these* with our informal concepts ‘and’, ‘not’, ‘if’ and ‘all’.

The logician’s *sharply defined* concepts \wedge , \rightarrow , \forall , etc. are *simple, well-behaved, useful* and *can be followed*.

Defining rules suffice for us to *coordinate* and to *communicate*, especially in those contexts where precision is valuable.

These defining rules give rise to Gentzen-style left/right rules in the presence of *Id* and *Cut*, and the resulting rules allow for the usual *cut elimination* argument.

Generality and Existence 1

THE REVIEW OF SYMBOLIC LOGIC
Volume 12, Number 1, March 2019

GENERALITY AND EXISTENCE 1: QUANTIFICATION AND FREE LOGIC

GREG RESTALL

School of Historical and Philosophical Studies, University of Melbourne

Abstract. In this paper, I motivate a cut free sequent calculus for classical logic with first order quantification, allowing for singular terms free of existential import. Along the way, I motivate a criterion for rules designed to answer Prior's question about what distinguishes rules for logical concepts, like conjunction from apparently similar rules for putative concepts like Prior's *tonk*, and I show that the rules for the quantifiers—and the existence predicate—satisfy that condition.

§1. Sequents and defining rules. Let's take it for granted for the moment that learning a language involves—at least in part—learning how assertions and denials expressed in that language bear on one another. The basic connection, of course, is that to assert *A* and to deny *A* clash. When we learn conjunction, we learn that there is a clash involved in asserting *A*, asserting *B* and denying *A* \wedge *B*. Similarly, when we learn disjunction, we learn that there is a clash involved in asserting *A* \vee *B*, denying *A* and denying *B*.

One way to systematically take account of the kinds of clashes involved in these acts of assertion and denial is through the language of the sequent calculus. Given collections Γ and Δ of sentences from our language \mathcal{L} , a sequent $\Gamma \succ \Delta$ makes the claim that there is a clash involved in asserting each element of Γ and denying each element of Δ . The structural rules of the sequent calculus can be understood in the following way [16].

Identity:

$$A \succ A.$$

There is a clash involved in asserting *A* and denying *A*. *Weakening*¹:

$$\frac{\Gamma \succ \Delta}{\Gamma, A \succ \Delta} [KL] \qquad \frac{\Gamma \succ \Delta}{\Gamma \succ A, \Delta} [KR].$$

DEFINING IDENTITY

Identity and Harmony

Identity and harmony

STEPHEN READ

1. Harmony

The inferentialist account of logic says that the meaning of a logical operator is given by the rules for its application. Prior (1960–61) showed that a simple and straightforward interpretation of this account of logicality reduces to absurdity. For if ‘tonk’ has the meaning given by the rules Prior proposed for it, contradiction follows. Accordingly, a more subtle interpretation of inferentialism is needed. Such a proposal was put forward initially by Gentzen (1934) and elaborated by, e.g., Prawitz (1977). The meaning of a logical expression is given by the rules for the assertion of statements containing that expression (as designated component); these are its introduction-rules. The meaning so given justifies further rules for drawing inferences from such assertions; these are its elimination-rules:

The introductions represent, as it were, the ‘definitions’ of the symbols concerned, and the eliminations are no more, in the final analysis, than the consequence of these definitions. (Gentzen 1934: 80)

For example, if the only ground for assertion of ‘ p tonk q ’ is given by Prior’s rule:

$$\frac{p}{p \text{ tonk } q} \text{ tonk-I}$$

then Prior mis-stated the elimination-rule. It should read

$$\frac{(p) \quad p \text{ tonk } q}{r} \text{ tonk-E}$$



A Defining Rule for Identity

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

A Defining Rule for Identity

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

Denying $a = b$ has the same significance as taking there to be *some* feature F that holds of a but not b , or *vice versa*.

A Defining Rule for Identity

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

Denying $a = b$ has the same significance as taking there to be *some* feature F that holds of a but not b , or *vice versa*.

Or equivalently, to prove that $a = b$, prove Fb from the assumption Fa (and *vice versa*), where the predicate F is *arbitrary*.

A Defining Rule for Identity

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

Denying $a = b$ has the same significance as taking there to be *some* feature F that holds of a but not b , or *vice versa*.

Or equivalently, to prove that $a = b$, prove Fb from the assumption Fa (and *vice versa*), where the predicate F is *arbitrary*.

Identity is a kind of *indistinguishability*.

An Example Derivation

$$\frac{\begin{array}{c} \mathsf{Fa} \succ \mathsf{Fa} \quad \mathsf{Fa} \succ \mathsf{Fa} \\ \hline \succ a = a \end{array}}{\succ (\lambda x. x = a) a} =_{Df\downarrow}$$
$$\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, \mathsf{Fa} \succ \mathsf{Fb} \end{array}}{a = b, (\lambda x. x = a) a \succ (\lambda x. x = a) b} =_{Df\uparrow}$$
$$\frac{\begin{array}{c} a = b \succ (\lambda x. x = a) b \\ \hline a = b \succ b = a \end{array}}{a = b \succ b = a} Spec_{(\lambda x. x = a)}^{\mathsf{F}}$$

Cut

An Example Derivation

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

$$\frac{\begin{array}{c} Fa \succ Fa \quad Fa \succ Fa \\ \hline \succ a = a \end{array}}{\succ (\lambda x. x = a) a} =Df\downarrow$$

$$\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, Fa \succ Fb \end{array}}{a = b, (\lambda x. x = a) a \succ (\lambda x. x = a) b} =Df\uparrow$$

*Spec*_($\lambda x. x = a$)^F *Cut*

$$\frac{a = b \succ (\lambda x. x = a) b}{a = b \succ b = a} \lambda Df\uparrow$$

An Example Derivation

$$\frac{X \succ Y}{X[F/P] \succ Y[F/P]} Spec_P^F$$

$$\frac{\begin{array}{c} Fa \succ Fa & Fa \succ Fa \\ \hline \succ a = a \end{array}}{\succ (\lambda x. x = a) a} = Df \downarrow$$

$$\frac{\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, Fa \succ Fb \end{array}}{a = b, (\lambda x. x = a) a \succ (\lambda x. x = a) b} Spec_{(\lambda x. x = a)}^F}{\hline} Cut$$

$$\frac{\hline}{a = b \succ b = a} \lambda Df \uparrow$$

$$a = b \succ (\lambda x. x = a) b$$

$$a = b \succ b = a$$

An Example Derivation

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

$$\frac{\frac{Fa \succ Fa \quad Fa \succ Fa}{\frac{\succ a = a}{\succ (\lambda x. x = a)a}} =Df\downarrow \quad \frac{a = b \succ a = b}{\frac{a = b, Fa \succ Fb}{a = b, (\lambda x. x = a)a \succ (\lambda x. x = a)b}} =Df\uparrow}{a = b, (\lambda x. x = a)a \succ (\lambda x. x = a)b} Spec_{(\lambda x. x = a)}^F Cut$$
$$\frac{a = b \succ (\lambda x. x = a)b}{a = b \succ b = a} \lambda Df\uparrow$$

An Example Derivation

$$\frac{X \succ A[x/a], Y}{X \succ (\lambda x.A)a, Y} \lambda Df$$

$$\frac{\begin{array}{c} Fa \succ Fa \\ Fa \succ Fa \end{array}}{\begin{array}{c} \succ a = a \\ \succ (\lambda x.x = a)a \end{array}} =Df\downarrow$$

$$\frac{\begin{array}{c} a = b \succ a = b \\ a = b, Fa \succ Fb \end{array}}{a = b, (\lambda x.x = a)a \succ (\lambda x.x = a)b} =Df\uparrow \text{ Spec}_{(\lambda x.x = a)}^F$$

$$\frac{a = b \succ (\lambda x.x = a)b}{a = b \succ b = a} \lambda Df\uparrow$$

An Example Derivation

$$\frac{\begin{array}{c} \mathsf{Fa} \succ \mathsf{Fa} \quad \mathsf{Fa} \succ \mathsf{Fa} \\ \hline \succ a = a \end{array}}{\succ (\lambda x. x = a) a} =_{Df\downarrow}$$
$$\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, \mathsf{Fa} \succ \mathsf{Fb} \end{array}}{a = b, (\lambda x. x = a) a \succ (\lambda x. x = a) b} =_{Df\uparrow}$$
$$\frac{a = b \succ (\lambda x. x = a) b}{a = b \succ b = a} Spec_{(\lambda x. x = a)}^{\mathsf{F}}$$
$$Cut$$

An Example Derivation

$$\frac{X \succ A[x/a], Y}{X \succ (\lambda x.A)a, Y} \lambda Df$$

$$\frac{\begin{array}{c} Fa \succ Fa & Fa \succ Fa \\ \hline \succ a = a & \end{array}}{\begin{array}{c} \succ (\lambda x.x = a)a \\ \hline \end{array}} \lambda Df\downarrow$$

$$\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, Fa \succ Fb \end{array}}{a = b, (\lambda x.x = a)a \succ (\lambda x.x = a)b} \frac{=Df\uparrow}{Spec_{(\lambda x.x = a)}^F} Cut$$

$$\frac{a = b \succ (\lambda x.x = a)b}{a = b \succ b = a} \lambda Df\uparrow$$

An Example Derivation

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df \quad \frac{X \succ A[x/a], Y}{X \succ (\lambda x.A)a, Y} \lambda Df \quad \frac{X \succ Y}{X[F/P] \succ Y[F/P]} Spec_P^F$$

$$\frac{\begin{array}{c} Fa \succ Fa \quad Fa \succ Fa \\ \succ a = a \end{array}}{\begin{array}{c} \succ a = a \\ \succ (\lambda x.x = a)a \end{array}} =Df\downarrow \quad \frac{\begin{array}{c} a = b \succ a = b \\ a = b, Fa \succ Fb \end{array}}{a = b, (\lambda x.x = a)a \succ (\lambda x.x = a)b} =Df\uparrow$$

$Spec_{(\lambda x.x = a)}^F$ Cut

$$\frac{a = b \succ (\lambda x.x = a)b}{a = b \succ b = a} \lambda Df\uparrow$$

From Df to L/R : $=R$ is $=Df \downarrow$

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =_{Df \downarrow}$$

From Df to L/R : These are the $=L$ rules

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1}$$

$$\frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

From Df to L/R : These are the $=L$ rules

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1}$$

$$\frac{\frac{\frac{X \succ Pa, Y}{\frac{\frac{a = b \succ a = b}{a = b, Fa \succ Fb} =_{Df\uparrow}}{\frac{a = b, Pa \succ Pb}{a = b, X \succ Pb, Y} Spec_P^F} Cut}{a = b, X \succ Pb, Y} Cut \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} Cut$$

From Df to L/R : These are the $=L$ rules

$$\frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

$$\frac{\begin{array}{c} a = b \succ a = b \\ \hline a = b, Fb \succ Fa \end{array} =_{Df\uparrow} \quad a = b, Pb \succ Pa \quad Spec_P^F}{\begin{array}{c} a = b, X \succ Pa, Y \\ \hline a = b, X, X' \succ Y, Y' \end{array} Cut \quad Cut}$$

From Df to L/R : These are the $=L$ rules

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1}$$

$$\frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

Going Without Some Rules

$$\frac{X \succ Y}{X[F/P] \succ Y[F/P]} Spec_P^F$$

$$\frac{X \succ A, Y \quad X', A \succ Y'}{X, X' \succ Y, Y'} Cut$$

Going Without Some Rules

$$\frac{X \succ Y}{X[F/P] \succ Y[F/P]} Spec_P^F$$

- *Spec* is *height-preserving admissible* in the system with L/R rules + *Cut*.

Going Without Some Rules

$$\frac{X \succ A, Y \quad X', A \succ Y'}{X, X' \succ Y, Y'} \textit{Cut}$$

- ▶ *Spec* is *height-preserving admissible* in the the system with L/R rules + *Cut*.
- ▶ *Cut* is eliminable in the system with L/R rules.

Eliminating *Cut*

$$\frac{\vdots \delta_1 \quad \vdots \delta_2 \quad \vdots \delta' \quad \vdots \delta''}{\begin{array}{c} X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y \\ X \succ a = b, Y \end{array}} =_R \quad \frac{X' \succ Pa, Y' \quad X'', Pb \succ Y''}{\begin{array}{c} a = b, X', X'' \succ Y', Y'' \\ Cut \end{array}} =_{L_1}$$
$$X, X', X'' \succ Y, Y', Y''$$

Eliminating Cut

$$\frac{\vdots \delta_1 \quad \vdots \delta_2 \quad \vdots \delta' \quad \vdots \delta''}{\begin{array}{c} X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y \\ X \succ a = b, Y \end{array}} =_R \quad \frac{\begin{array}{c} X' \succ Pa, Y' \quad X'', Pb \succ Y'' \\ a = b, X', X'' \succ Y', Y'' \end{array}}{X, X', X'' \succ Y, Y', Y''} =_{L_1} Cut$$

becomes

$$\frac{\vdots \delta' \quad \vdots \delta_1[F/P] \quad \vdots \delta''}{\begin{array}{c} X' \succ Pa, Y' \quad X, Pa \succ Pb, Y \\ X, X' \succ Pb, Y, Y' \end{array}} Cut \quad \frac{X'', Pb \succ Y''}{X, X', X'' \succ Y, Y', Y''} Cut$$

But eliminating *Cut* hardly seems worth it!

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =_R$$

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1} \quad \frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

But eliminating *Cut* hardly seems worth it!

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =_R$$

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1} \quad \frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

Violations of the subformula property *everywhere*.

But eliminating *Cut* hardly seems worth it!

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =_R$$

$$\frac{X \succ Pa, Y \quad X', Pb \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_1} \quad \frac{X \succ Pb, Y \quad X', Pa \succ Y'}{a = b, X, X' \succ Y, Y'} =_{L_2}$$

Violations of the subformula property *everywhere*.

For *analytic* derivations, we need different rules.

From Rules to Axioms: From $=R$ to Refl

$$\frac{}{\succ a = a} \text{Refl}$$

From Rules to Axioms: From $=R$ to Refl

$$\frac{}{\succ a = a} \text{Refl}$$

$$\frac{\vdash a \succ a \quad \vdash a \succ a}{\succ a = a} =R$$

From *Refl* to $=R$

Replace this:

$$\frac{\vdots \delta_1 \quad \vdots \delta_2}{\begin{array}{c} X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y \\ \hline X \succ a = b, Y \end{array}} =R$$

From Refl to $=R$

Replace this:

$$\frac{\vdots \delta_1 \quad \vdots \delta_2}{\begin{array}{c} X, F a \succ F b, Y \\ X, F b \succ F a, Y \end{array}} =R$$
$$X \succ a = b, Y$$

With this:

$$\frac{\frac{\frac{\vdash a = a}{\succ a = a}^{\text{Refl}} \quad \vdots \delta_1[F/\lambda x.(a = x)]}{\succ \lambda x.(a = x)a}^{\lambda R} \quad X, \lambda x.(a = x)a \succ \lambda x.(a = x)b, Y}{\frac{X \succ \lambda x.(a = x)b, Y}{X \succ a = b, Y}}^{\lambda R} \quad \text{Cut}$$

From $=L$ to $=Lax$ and back.

$$\frac{}{a = b, Pa \succ Pb} =_{Lax_1} \quad \frac{}{a = b, Pb \succ Pa} =_{Lax_2}$$

From $=L$ to $=Lax$ and back.

$$\frac{}{a = b, Pa \succ Pb} =_{Lax_1} \quad \frac{}{a = b, Pb \succ Pa} =_{Lax_2}$$

$$\frac{Pa \succ Pa \quad Pb \succ Pb}{a = b, Pa \succ Pb} =_{L_1} \quad \frac{Pb \succ Pb \quad Pa \succ Pa}{a = b, Pb \succ Pa} =_{L_2}$$

From $=L$ to $=Lax$ and back.

$$\frac{}{a = b, Pa \succ Pb} =L.ax_1 \quad \frac{}{a = b, Pb \succ Pa} =L.ax_2$$

$$\frac{\begin{array}{c} X \succ Pa, Y \\ \frac{\begin{array}{c} a = b, Pa \succ Pb \\ \frac{}{a = b, X \succ Pb, Y} =L.ax_1 \end{array}}{a = b, X, X' \succ Y, Y'} Cut \\ X', Pb \succ Y' \end{array}}{a = b, X, X' \succ Y, Y'} Cut$$

$$\frac{\begin{array}{c} X \succ Pb, Y \\ \frac{\begin{array}{c} a = b, Pb \succ Pa \\ \frac{}{a = b, X \succ Pa, Y} =L.ax_2 \end{array}}{a = b, X, X' \succ Y, Y'} Cut \\ X', Pa \succ Y' \end{array}}{a = b, X, X' \succ Y, Y'} Cut$$

We can restrict $=L.ax$ to primitive predicates

$$\frac{\frac{\frac{a = b, Pa \succ Pb}{a = b, Pa \wedge Qa \succ Pb} =L.ax_1}{a = b, Pa \wedge Qa \succ Pb} \wedge_L \quad \frac{a = b, Qa \succ Qb}{a = b, Pa \wedge Qa \succ Qb} =L.ax_1}{a = b, Pa \wedge Qa \succ Pa \wedge Qb} \wedge_R \\ \frac{a = b, Pa \wedge Qa \succ Pa \wedge Qb}{a = b, \lambda x.(Px \wedge Qx)a \succ \lambda x.(Px \wedge Qx)b} \lambda$$

We can restrict $=L.ax$ to primitive predicates

$$\frac{\frac{\frac{\overline{a = b, Pb \succ Pa}}{a = b, Pb, \neg Pa \succ} \neg_L}{a = b, \neg Pa \succ \neg Pb} \neg_R}{a = b, \lambda x.(\neg Px)a \succ \lambda x.(\neg Px)b} \lambda$$

We can restrict $=Lax$ to primitive predicates

$$\frac{\frac{\frac{\overline{a = b, Pac \succ Pbc}}{=Lax_2} \quad \neg L}{a = b, \forall y Pay \succ Pbc} \quad \neg R}{a = b, \forall y Pay \succ \forall y Pby} \quad \lambda}{a = b, \lambda x. (\forall y Pxy) a \succ \lambda x. (\forall y Pxy) b}$$

Now eliminate *Cut*

$$\frac{\frac{}{a = b, Pa \succ Pb} =L.ax_1 \quad \frac{}{c = b, Pb \succ Pc} =L.ax_2}{a = b, c = b, Pa \succ Pc} Cut$$

Now eliminate *Cut*

$$\frac{\frac{}{a = b, Pa \succ Pb} =L.ax_1 \quad \frac{}{c = b, Pb \succ Pc} =L.ax_2}{a = b, c = b, Pa \succ Pc} Cut$$

becomes

$$\frac{}{a = b, c = b, Pa \succ Pc} =L??$$

Now eliminate *Cut*

It suffices to close the *axioms* under *Cut*.

$$\frac{}{I_b^a, P_a \succ P_b} =_{L.ax^*}$$

Where I_b^a is any multiset
of identities *linking a to b*,
and P is any primitive predicate.

Now eliminate *Cut*

It suffices to close the *axioms* under *Cut*.

$$\frac{}{I_b^a, P_a \succ P_b} =_{L.ax^*}$$

Where I_b^a is any multiset
of identities *linking a to b*,
and P is any primitive predicate.

- (a) The *empty* multiset links a to a.
- (b) $a = b$ links a to b and b to a.
- (c) If X links a to b and Y links
b to c then X, Y links a to c.

Now eliminate Cut

It suffices to close the *axioms* under *Cut*.

$$\frac{}{I_b^a, P_a \succ P_b} =_{L.ax^*}$$

Where I_b^a is any multiset
of identities *linking a to b*,
and P is any primitive predicate.

- (a) The *empty* multiset links a to a.
- (b) $a = b$ links a to b and b to a.
- (c) If X links a to b and Y links
b to c then X, Y links a to c.

(We can leave ' P_a ' out if it is $a = a$.)

Kinds of Identity Rules

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

- $=Df$ introduces the identity predicate in *arbitrary* positions, defining the denial of $a = b$ in terms of *distinguishing* a and b .

Kinds of Identity Rules

$$\frac{}{\succ a = a} \text{Refl} \qquad \frac{}{I_b^a, P_a \succ P_b} =_{L.ax_*}$$

- $=_{Df}$ introduces the identity predicate in *arbitrary* positions, defining the denial of $a = b$ in terms of *distinguishing* a and b .
- Refl and $=_{L.ax_*}$ are *semantic constraints connecting* primitive predicates.

Kinds of Identity Rules

$$\frac{X, Fa \succ Fb, Y \quad X, Fb \succ Fa, Y}{X \succ a = b, Y} =Df$$

$$\frac{}{\succ a = a} Refl \qquad \frac{}{I_b^a, Pa \succ Pb} =L.ax_*$$

- $=Df$ introduces the identity predicate in *arbitrary* positions, defining the denial of $a = b$ in terms of *distinguishing* a and b .
- $Refl$ and $=L.ax_*$ are *semantic constraints connecting* primitive predicates.
- They are *equivalent* as far as derivability goes.

Identity depends on *predication*

To apply identity rules, we need to agree
on what counts as a *singular term*
and what counts as a *predicate*.

DEFINING NECESSITY

Modal reasoning trades on *supposition* and *context shift*

Oswald shot Kennedy. But suppose he *hadn't*.

Modal reasoning trades on *supposition* and *context shift*

Oswald shot Kennedy. But suppose he *hadn't*.

I think that Oswald shot Kennedy.
But let's suppose that you're right and he didn't.

Modal reasoning trades on *supposition* and *context shift*

Oswald shot Kennedy. But suppose he *hadn't*.

I think that Oswald shot Kennedy.
But let's suppose that *you're* right and he *didn't*.

Oswald shot Kennedy | Oswald didn't shoot Kennedy

Two notions of necessity



1. SUBJUNCTIVE/METAPHYSICAL — how things *could have been* (had things gone differently) — If Oswald *hadn't* shot Kennedy ...

Two notions of necessity



1. SUBJUNCTIVE/METAPHYSICAL — how things *could have been* (had things gone differently) — If Oswald *hadn't* shot Kennedy ...
2. INDICATIVE/EPISTEMIC — how things *could actually be* (given some body of evidence) — If Oswald *didn't* shoot Kennedy ...

Two notions of necessity



1. SUBJUNCTIVE/METAPHYSICAL — how things *could have been* (had things gone differently) — If Oswald *hadn't* shot Kennedy ...
 2. INDICATIVE/EPISTEMIC — how things *could actually be* (given some body of evidence) — If Oswald *didn't* shoot Kennedy ...
- + Possible worlds semantics is useful in modelling *both* modal notions.

Two notions of necessity



1. SUBJUNCTIVE/METAPHYSICAL — how things *could have been* (had things gone differently) — If Oswald *hadn't* shot Kennedy ...
2. INDICATIVE/EPISTEMIC — how things *could actually be* (given some body of evidence) — If Oswald *didn't* shoot Kennedy ...
 - + Possible worlds semantics is useful in modelling *both* modal notions.
 - + Let's consider both these options in the development of a *proof system*.

An Analogy

A *modal operator* stands to a kind of *context shift*
as a *quantifier* stands to a category of *terms*.

Hypersequents

$$\Box A \succ \mid \succ A$$

A Defining Rule for Necessity

$$\frac{\succ A | X \succ Y | \mathcal{S}}{X \succ \Box A, Y | \mathcal{S}} \Box Df$$

A Defining Rule for Necessity

$$\frac{\succ A \mid X \succ Y \mid \mathcal{S}}{X \succ \Box A, Y \mid \mathcal{S}} \Box Df$$

Denying that A is necessary has the force of denying A in some context.

A Defining Rule for Necessity

$$\frac{\succ A \mid X \succ Y \mid \mathcal{S}}{X \succ \Box A, Y \mid \mathcal{S}} \Box Df$$

Denying that A is necessary has the force of denying A in some context.

Or, to prove that A is necessary, you prove A in an arbitrary context.

A Defining Rule for Necessity

$$\frac{\succ A \mid X \succ Y \mid \mathcal{S}}{X \succ \Box A, Y \mid \mathcal{S}} \Box Df$$

Denying that A is necessary has the force of denying A in some context.

Or, to prove that A is necessary, you prove A in an arbitrary context.

So, necessity expresses a kind of *preservation across contexts*.

A Defining Rule for Necessity

$$\frac{\succ A \mid X \succ Y \mid S}{X \succ \Box A, Y \mid S} \Box Df$$

Denying that A is necessary has the force of denying A in some context.

Or, to prove that A is necessary, you prove A in an arbitrary context.

So, necessity expresses a kind of *preservation across contexts*.

Indicative shifts define a kind of ‘epistemic’ necessity (\Box_E).

Subjunctive shifts define a kind of ‘metaphysical’ necessity (\Box_M).

A Defining Rule for Necessity

$$\frac{\succ A \mid X \succ Y \mid S}{X \succ \Box A, Y \mid S} \Box Df$$

Denying that A is necessary has the force of denying A in some context.

Or, to prove that A is necessary, you prove A in an arbitrary context.

So, necessity expresses a kind of *preservation across contexts*.

Indicative shifts define a kind of ‘epistemic’ necessity (\Box_E).

Subjunctive shifts define a kind of ‘metaphysical’ necessity (\Box_M).

The *form* of the rule is the same in either case, though the *content* differs.

An example derivation

$$\frac{\frac{\frac{\Box p \succ \Box p}{\succ p \mid \Box p \succ} \Box Df \uparrow \quad \frac{\neg \Box p \succ \neg \Box p}{\Box p, \neg \Box p \succ} \neg Df \uparrow}{\succ \Box p \mid \Box p \succ} \Box Df \downarrow \quad \frac{}{\Box p, \neg \Box p \succ} \neg Df \uparrow}{\frac{}{\neg \Box p \succ \mid \Box p \succ} \neg Df \downarrow \quad \frac{\neg \Box p \succ \mid \succ \neg \Box p}{\neg \Box p \succ \Box \neg \Box p} \Box Df \downarrow} Cut$$

Both kinds of shift are important to creatures like us

- We *disagree*. We have reason to come to shared positions.

Both kinds of shift are important to creatures like us

- ▶ We *disagree*. We have reason to come to shared positions.
- ▶ We *plan*. We have reason to consider options (prospectively) or to replay scenarios (retrospectively).

Both kinds of shift are important to creatures like us

- ▶ We *disagree*. We have reason to come to shared positions.
- ▶ We *plan*. We have reason to consider options (prospectively) or to replay scenarios (retrospectively).
- ▶ Our everyday practice is messy and informal. *Formalising* an aspect of it helps us isolate distinctive features of that practice.

Both kinds of shift are important to creatures like us

- ▶ We *disagree*. We have reason to come to shared positions.
- ▶ We *plan*. We have reason to consider options (prospectively) or to replay scenarios (retrospectively).
- ▶ Our everyday practice is messy and informal. *Formalising* an aspect of it helps us isolate distinctive features of that practice.
- ▶ None of this requires an antecedent commitment to an ontology of *worlds*.

Combining Indicative and Subjunctive Shifts

You can combine the two kinds of context shift
into one hypersequent system easily enough

(see my “A Cut-Free Sequent System for Two-Dimensional Modal Logic”)
but we don’t need to worry about those details here.

From $\Box Df$ to $\Box L/\Box R$

$$\frac{\succ A | X \succ Y | S}{X \succ \Box A, Y | S} \Box Df$$

From $\Box Df$ to $\Box L/\Box R$

$$\frac{\succ A | X \succ Y | S}{X \succ \Box A, Y | S} \Box R$$

From $\Box Df$ to $\Box L/\Box R$

$$\frac{\succ A | X \succ Y | S}{X \succ \Box A, Y | S} \Box R$$

$$\frac{\frac{\Box A \succ \Box A}{\succ A | \Box A \succ} \Box Df \uparrow \quad X, A \succ Y | S}{\Box A \succ | X \succ Y | S} Cut$$

From $\Box Df$ to $\Box L/\Box R$

$$\frac{\succ A | X \succ Y | S}{X \succ \Box A, Y | S} \Box R$$

$$\frac{X, A \succ Y | S}{\Box A \succ | X \succ Y | S} \Box L$$


COMBINING THEM

Doesn't identity become necessary, *automatically*?

$$\frac{\frac{\frac{\vdash a = a}{\vdash a = a} Refl}{\vdash \Box(a = a)} \Box I}{\vdash \lambda x. \Box(a = x) a} \lambda Df \downarrow \quad \frac{}{a = b, \lambda x. \Box(a = x) a \succ \lambda x. \Box(a = x) b} =L.ax_1$$

$$\frac{}{a = b \succ \lambda x. \Box(a = x) b} \lambda Df \uparrow \quad \frac{a = b \succ \lambda x. \Box(a = x) b}{a = b \succ \Box(a = b)} \lambda Df \uparrow$$

Willamson *defends* the necessity of identity



Williamson on the necessity of identity

... to mess with the modal or temporal logic of identity in order to avoid ontological inflation would be a lapse of methodological good taste, or good sense, for it means giving more weight to ontology than to the vastly better developed and more successful discipline of logic.

More specifically, the classical modal or temporal logic is a strong, simple, and elegant theory. To weaken, complicate, and uglify it without overwhelming reason to do so merely in order to block the derivation of the necessity or permanence of identity would be as retrograde and wrong-headed a step in logic and metaphysics as natural scientists would consider a comparable sacrifice of those virtues in a physical theory.

— Timothy Williamson, *Modal Logic as Metaphysics*, pp. 26, 27.


Modal Logic with Contingent Identity




Giovanna Corsi “Counterpart Semantics” 2002




Maria Aloni “Individual Concepts in Modal Predicate Logic” 2005




Hesperus



Phosphorous



Hesperus



Phosphorous

$$h = p \mid h \neq p$$

Identity and Context Shift

$$a = b \succ | Fa \succ Fb$$

Is it consistent to grant that $a = b$ in one context,
while distinguishing a from b in another?

Identity and Context Shift

$$a = b \succ | Fa \succ Fb$$

Is it consistent to grant that $a = b$ in one context,
while distinguishing a from b in another?

Indicative shifts (*disagreement*): This seems consistent.
(We *learned* that $h = p$. The possibility that $h \neq p$ was open to us.)

Identity and Context Shift

$$a = b \succ | Fa \succ Fb$$

Is it consistent to grant that $a = b$ in one context,
while distinguishing a from b in another?

- Indicative shifts (*disagreement*): This seems consistent.
(We *learned* that $h = p$. The possibility that $h \neq p$ was open to us.)
- Subjunctive shifts (*planning*): This is much less plausible.
($h = p$. Were I to travel to h , then of necessity, I am going to p .)

More from Williamson

... we are not interested in epistemic readings of 'it is possible that'.

— Timothy Williamson, *Modal Logic as Metaphysics*, p. 11, fn. 17.

Subjunctive Context Shifts: *Necessary Identity Rules*

$$\frac{X \succ Y | X', F a \succ F b, Y' | \mathcal{S} \quad X \succ Y | X', F b \succ F a, Y' | \mathcal{S}}{X \succ a = b, Y | X' \succ Y' | \mathcal{S}} = Df$$

Subjunctive Context Shifts: Necessary Identity Rules

$$\frac{X \succ Y | X', Fa \succ Fb, Y' | \mathcal{S} \quad X \succ Y | X', Fb \succ Fa, Y' | \mathcal{S}}{X \succ a = b, Y | X' \succ Y' | \mathcal{S}} = Df$$

$$\frac{}{\succ a = a} Refl \quad \frac{I_b^a | Pa \succ Pb}{I_b^a | Pa \succ Pb} = L.ax_*$$

I_b^a is a multiset of identities, linking a and b , distributed across the zones of the hypersequent, in the LHS of each sequent.

(And again, we can leave Pa out if it is $a = a$.)

Subjunctive Context Shifts: Necessary Identity Rules

$$\frac{X \succ Y | X', Fa \succ Fb, Y' | \mathcal{S} \quad X \succ Y | X', Fb \succ Fa, Y' | \mathcal{S}}{X \succ a = b, Y | X' \succ Y' | \mathcal{S}} =Df$$

$$\frac{}{\succ a = a} Refl \quad \frac{}{I_b^a | Pa \succ Pb} =L.ax_*$$

I_b^a is a multiset of identities, linking a and b , distributed across the zones of the hypersequent, in the LHS of each sequent.

(And again, we can leave Pa out if it is $a = a$.)

To deny $a = b$ is to take a and b to be *distinguishable*.

A Cut-Free Derivation

$$\frac{\overline{a = b \succ \mid \succ a = b}}{a = b \succ \Box a = b} =Lax_* \Box R$$

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$
- ▶ DERIVE THIS: $a = b, Fa \succ Fb$

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$
- ▶ DERIVE THIS: $a = b, Fa \succ Fb$
- ▶ This requires distinguishing kinds of predicates, because we *also* want to reject $a = b, \lambda x. \Box_E(a = x)a \succ \lambda x. \Box_E(a = x)b$.

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$
- ▶ DERIVE THIS: $a = b, Fa \succ Fb$
- ▶ This requires distinguishing kinds of predicates, because we *also* want to reject $a = b, \lambda x. \Box_E(a = x)a \succ \lambda x. \Box_E(a = x)b$.
 - ▶ To distinguish a from b we want to find some *feature* a has that b lacks (or *vice versa*).

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$
- ▶ DERIVE THIS: $a = b, Fa \succ Fb$
- ▶ This requires distinguishing kinds of predicates, because we *also* want to reject $a = b, \lambda x. \Box_E(a = x)a \succ \lambda x. \Box_E(a = x)b$.
 - ▶ To distinguish a from b we want to find some *feature* a has that b lacks (or *vice versa*). However, epistemic modalities do not always generate features.

Epistemic Shifts: Contingent Identity

- ▶ BLOCK THIS: $a = b \succ | Fa \succ Fb$
- ▶ DERIVE THIS: $a = b, Fa \succ Fb$
- ▶ This requires distinguishing kinds of predicates, because we *also* want to reject $a = b, \lambda x. \Box_E(a = x)a \succ \lambda x. \Box_E(a = x)b$.
 - ▶ To distinguish a from b we want to find some *feature* a has that b lacks (or *vice versa*). However, epistemic modalities do not always generate features.
 - ▶ After all, it's no argument against $h = p$ that it's epistemically necessary that $h = h$ and not epistemically necessary that $h = p$.

Two Kinds of Predicates

F : *feature predicates*

G : *general predicates*

Two Kinds of Predicates

F : *feature predicates*

G : *general predicates*

- ▶ Feature predicates are closed under the classical connectives and quantifiers and λ .

Two Kinds of Predicates

F : *feature predicates*

G : *general predicates*

- ▶ Feature predicates are closed under the classical connectives and quantifiers and λ .
- ▶ Applying λ into \Box_E contexts creates a *general predicate*.

Two Kinds of Predicates

F : *feature predicates*

G : *general predicates*

- ▶ Feature predicates are closed under the classical connectives and quantifiers and λ .
- ▶ Applying λ into \Box_E contexts creates a *general predicate*.
- ▶ There are two kinds of *Spec* rule: one for feature predicates, and one for general predicates.

Contingent Identity Rules

$$\frac{X, Fa \succ Fb, Y | \mathcal{S} \quad X, Fb \succ Fa, Y | \mathcal{S}}{a = b, X \succ Y | \mathcal{S}} = Df$$

Contingent Identity Rules

$$\frac{X, Fa \succ Fb, Y | \mathcal{S} \quad X, Fb \succ Fa, Y | \mathcal{S}}{a = b, X \succ Y | \mathcal{S}} = Df$$

$$\frac{}{\succ a = a} Refl \quad \frac{I_b^a, Pa \succ Pb}{\quad} = L.ax_*$$

I_b^a is a multiset of identities, linking a and b .

P is a *feature predicate*.

(And again, we can leave Pa out if it is $a = a$.)

Contingent Identity Rules

$$\frac{X, Fa \succ Fb, Y | \mathcal{S} \quad X, Fb \succ Fa, Y | \mathcal{S}}{a = b, X \succ Y | \mathcal{S}} = Df$$

$$\frac{}{\succ a = a} Refl \quad \frac{I_b^a, Pa \succ Pb}{\quad} = L.ax_*$$

I_b^a is a multiset of identities, linking a and b .

P is a *feature predicate*.

(And again, we can leave Pa out if it is $a = a$.)

To deny $a = b$ is to take there to be some way to distinguish a and b .

Two Dimensions—with *actuality*

$$\begin{array}{c} X_1^1 \succ_{@} Y_1^1 \quad | \quad X_2^1 \succ Y_2^1 \quad | \quad \cdots \quad | \quad X_{m_1}^1 \succ Y_{m_1}^1 \quad || \\ X_1^2 \succ_{@} Y_1^2 \quad | \quad X_2^2 \succ Y_2^2 \quad | \quad \cdots \quad | \quad X_{m_2}^2 \succ Y_{m_2}^2 \quad || \\ \vdots \qquad \qquad \vdots \qquad \qquad \vdots \\ X_1^n \succ_{@} Y_1^n \quad | \quad X_2^n \succ Y_2^n \quad | \quad \cdots \quad | \quad X_{m_n}^n \succ Y_{m_n}^n \end{array}$$

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*.

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*. (If \Box is introduced twice, given the same hypersequent separator, those operators are equivalent.)

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*. (If \Box is introduced twice, given the same hypersequent separator, those operators are equivalent.)
- ▶ The necessary identity rules are conservative and uniquely defining, relative to the choice of *predicates* in the language.

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*. (If \Box is introduced twice, given the same hypersequent separator, those operators are equivalent.)
- ▶ The necessary identity rules are conservative and uniquely defining, relative to the choice of *predicates* in the language. (If two identity predicates are added using the same form of rules, they are equivalent, provided that both identity predicates count as *predicates* for the purposes of both sets of rules.)

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*. (If \Box is introduced twice, given the same hypersequent separator, those operators are equivalent.)
- ▶ The necessary identity rules are conservative and uniquely defining, relative to the choice of *predicates* in the language. (If two identity predicates are added using the same form of rules, they are equivalent, provided that both identity predicates count as *predicates* for the purposes of both sets of rules.)
- ▶ The contingent identity rules are conservative and uniquely defining, relative to the choice of *feature predicates* in the language.

Features of these rules

- ▶ The modal rules are conservative and uniquely defining, relative to the choice of *context shift*. (If \Box is introduced twice, given the same hypersequent separator, those operators are equivalent.)
- ▶ The necessary identity rules are conservative and uniquely defining, relative to the choice of *predicates* in the language. (If two identity predicates are added using the same form of rules, they are equivalent, provided that both identity predicates count as *predicates* for the purposes of both sets of rules.)
- ▶ The contingent identity rules are conservative and uniquely defining, relative to the choice of *feature predicates* in the language. (If two identity predicates are added using the same form of rules, they are equivalent, provided that both identity predicates count as *feature predicates* for the purposes of both sets of rules.)

Models

SLOGAN: The *limit* of the process of filling out an underivable (hyper)sequent describes a *model*.

Models

SLOGAN: The *limit* of the process of filling out an underivable (hyper)sequent describes a *model*.

$$\frac{X \succ A, Y | \mathcal{S} \quad X, A \succ Y | \mathcal{S}}{X \succ Y | \mathcal{S}} \text{Cut}' \qquad \frac{\succ A | X \succ \Box A, Y | \mathcal{S}}{X \succ \Box A, Y | \mathcal{S}} \Box R'$$

Models

SLOGAN: The *limit* of the process of filling out an underivable (hyper)sequent describes a *model*.

$$\frac{X \succ A, Y | \mathcal{S} \quad X, A \succ Y | \mathcal{S}}{X \succ Y | \mathcal{S}} \text{Cut}' \qquad \frac{\succ A | X \succ \Box A, Y | \mathcal{S}}{X \succ \Box A, Y | \mathcal{S}} \Box R'$$

The zones in correspond to *worlds*,
the formulas in the LHS are *true*
and those in the RHS are *false*.

Necessary Identity Models

- ▶ ‘ \equiv ’ determines an equivalence relation on terms.

Necessary Identity Models

- ▶ ‘ \equiv ’ determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.

Necessary Identity Models

- ▶ ' \equiv ' determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.
- ▶ ' \equiv ' determines a congruence relation for each predicate at each world.

Necessary Identity Models

- ▶ ‘ \equiv ’ determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.
- ▶ ‘ \equiv ’ determines a congruence relation for each predicate at each world.
- ▶ Take the quotient of the terms by identity to form your ‘domain’.

Necessary Identity Models

- ▶ ‘ \equiv ’ determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.
- ▶ ‘ \equiv ’ determines a congruence relation for each predicate at each world.
- ▶ Take the quotient of the terms by identity to form your ‘domain’.
- ▶ The expected truth conditions hold for the logical operators.

Necessary Identity Models

- ▶ ‘ \equiv ’ determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.
- ▶ ‘ \equiv ’ determines a congruence relation for each predicate at each world.
- ▶ Take the quotient of the terms by identity to form your ‘domain’.
- ▶ The expected truth conditions hold for the logical operators.
- ▶ These are regular models for constant domain s5 with necessary identity.

Necessary Identity Models


- ▶ ‘ \equiv ’ determines an equivalence relation on terms.
- ▶ Its extension is constant across each world.
- ▶ ‘ \equiv ’ determines a congruence relation for each predicate at each world.
- ▶ Take the quotient of the terms by identity to form your ‘domain’.
- ▶ The expected truth conditions hold for the logical operators.
- ▶ These are regular models for constant domain s5 with necessary identity.
- ▶ (The usual rules for quantifiers work, too.)

Contingent Identity Models

$$a = b, c = d \succ b = c \mid b = c \succ a = d \mid a = b, b = c \succ a = d$$


Contingent Identity Models

$$a = b, c = d \succ b = c \mid b = c \succ a = d \mid a = b, b = c \succ a = d$$




Contingent Identity Models

$$a = b, c = d \succ b = c \mid b = c \succ a = d \mid a = b, b = c \succ a = d$$




Contingent Identity Models

$F_a, F_b \quad F_c, F_d \quad F_a, F_d \quad F_b, F_c \quad F_a, F_b, F_c \quad F_d$
 $a = b, c = d \succ b = c \mid b = c \succ a = d \mid a = b, b = c \succ a = d$




Contingent Identity Models



F : feature predicate
 G : general predicate

Contingent Identity Models



THE UPSHOT

Identities: metaphysically necessary, epistemically contingent

- We would expect to have to *resolve* disagreements about identities. Identities are the kinds of things we can *learn* by engagement with the world around us, and they are *open* to us until we learn them.

Identities: metaphysically necessary, epistemically contingent

- ▶ We would expect to have to *resolve* disagreements about identities. Identities are the kinds of things we can *learn* by engagement with the world around us, and they are *open* to us until we learn them.
- ▶ On the other hand, once we have learned an identity, that should constrain our view of how things *could go*, or how things *could have gone*.

Identities: metaphysically necessary, epistemically contingent

- ▶ We would expect to have to *resolve* disagreements about identities. Identities are the kinds of things we can *learn* by engagement with the world around us, and they are *open* to us until we learn them.
- ▶ On the other hand, once we have learned an identity, that should constrain our view of how things *could go*, or how things *could have gone*.
- ▶ So, to define identity we require preservation of features, but not preservation by predicates that induce *epistemic* shifts.

Identities: metaphysically necessary, epistemically contingent

- ▶ We would expect to have to *resolve* disagreements about identities. Identities are the kinds of things we can *learn* by engagement with the world around us, and they are *open* to us until we learn them.
- ▶ On the other hand, once we have learned an identity, that should constrain our view of how things *could go*, or how things *could have gone*.
- ▶ So, to define identity we require preservation of features, but not preservation by predicates that induce *epistemic* shifts.
- ▶ So, a ‘property’ of the form $\lambda x. \Box_E (a = x)$ is not the kind of thing that is preserved by an identity. (Even if it holds of a , and if $a = b$, it need not hold of b .)

Identities: metaphysically necessary, epistemically contingent

- ▶ We would expect to have to *resolve* disagreements about identities. Identities are the kinds of things we can *learn* by engagement with the world around us, and they are *open* to us until we learn them.
- ▶ On the other hand, once we have learned an identity, that should constrain our view of how things *could go*, or how things *could have gone*.
- ▶ So, to define identity we require preservation of features, but not preservation by predicates that induce *epistemic* shifts.
- ▶ So, a ‘property’ of the form $\lambda x. \Box_E (a = x)$ is not the kind of thing that is preserved by an identity. (Even if it holds of a , and if $a = b$, it need not hold of b .)
- ▶ These distinctions can be explained and motivated in terms of our dialogical practice, and its aims. The result is a uniform treatment of epistemic and metaphysical modality and their distinctive interactions with identity.

There is More to Do

- ▶ Quantifiers

There is More to Do

- ▶ Quantifiers
- ▶ Sortals, Conceptual Covers

There is More to Do

- ▶ Quantifiers
- ▶ Sortals, Conceptual Covers
- ▶ Existence

There is More to Do

- ▶ Quantifiers
- ▶ Sortals, Conceptual Covers
- ▶ Existence
- ▶ Comparisons with other systems

References

CATARINA DUTILH NOVAES

The Dialogical Roots of Deduction

Cambridge University Press, 2021

STEPHEN READ

“Identity and Harmony”

Analysis, 64(2):113–115, 2004

GREG RESTALL

“Generality and Existence 1:

Quantification and Free Logic”

Review of Symbolic Logic, 12:1–29, 2019

TIMOTHY WILLIAMSON

Modal Logic as Metaphysics

Oxford University Press, 2013

GREG RESTALL

“A Cut-Free Sequent System for

Two-Dimensional Modal Logic, and
why it matters”

Annals of Pure and Applied Logic,

163(11):1611–1623, 2012

MARIA ALONI

“Individual Concepts in Modal
Predicate Logic”

Journal of Philosophical Logic,

34(1):1–64, 2005

GIOVANNA CORSI

“Counterpart Semantics: A
Foundational Study on Quantified
Modal Logics”

ILLC Technical Report PP-2002-20

ANIL GUPTA

The Logic of Common Nouns: An

Investigation in Quantified Modal Logic

Yale University Press, 1980

NUEL BELNAP AND THOMAS MÜLLER

“CIFOL: Case-Intensional First Order
Logic”

Journal of Philosophical Logic,

43(2):393–437, 2013

THANK YOU!