

# Proof Theory: Logical and Philosophical Aspects

## Class 5: Semantics and beyond

*Greg Restall and Shawn Standefer*



THE UNIVERSITY OF  
MELBOURNE

NASSLLI · JULY 2016 · RUTGERS

To introduce *proof theory*, with a focus in its applications in philosophy, linguistics and computer science.

Examine the connections between proof theory and semantics, both formal *model theory*, and more general philosophical considerations concerning meaning.

Speech Acts and Norms

Proofs and Models

Beyond

# SPEECH ACTS AND NORMS

# Normative Pragmatics

An idea found in Brandom's *Making It Explicit* is that the *meaning* of linguistic items should first be understood in terms of their *use*

The linguistic (conceptual) practices of communities set up *norms* governing their behavior

These practices have features that we can make explicit through the introduction of new vocabulary

# Rules as Definitions

The rules that govern a connective are taken to *define* the new connective

This appears to make it really easy to introduce new logical terms

Specify a set of rules governing a connective, and you've got a new connective

# Rules as Definitions

The rules that govern a connective are taken to *define* the new connective

This appears to make it really easy to introduce new logical terms

Specify a set of rules governing a connective, and you've got a new connective

But, there's a problem



# Tonk

Arthur Prior pointed out that if a set of rules is enough to define a connective, then *tonk* is legitimate

$$\frac{X, A \vdash C}{X, A \odot B \vdash C} \text{ [tonkL]}$$

$$\frac{X \vdash B}{X \vdash A \odot B} \text{ [tonkR]}$$

# Tonk

Arthur Prior pointed out that if a set of rules is enough to define a connective, then *tonk* is legitimate

$$\frac{X, A \vdash C}{X, A \odot B \vdash C} \text{ [tonkL]}$$

$$\frac{X \vdash B}{X \vdash A \odot B} \text{ [tonkR]}$$

$$\frac{\frac{B \vdash B}{B \vdash A \odot B} \text{ [tonkR]} \quad \frac{A \vdash A}{A \odot B \vdash A} \text{ [tonkL]}}{B \vdash A} \text{ [Cut]}$$

## Responding to tonk

Nuel Belnap responded to Prior's article, saying that additional conditions need to be satisfied in order to define a connective

Connectives aren't introduced out of thin air, there is a *context of deducibility*, e.g. the full set of Gentzen's structural rules

In order to be a definition, an extension has to be *conservative*, while tonk manifestly is *not*

In order to be a definition, an addition has to be *uniquely specified*

These ideas have been taken up and developed by Dummett and others in discussions of *harmony*

# Assertion and Denial

Many philosophers and logicians take *assertion* to be the primary speech act, which is used to define others

Others argue that *denial* should be understood as a primitive act on its own

We take logic, in particular valid sequents, as presenting normative relations between assertions and denials

$X \vdash Y$  tells us that one should not assert everything in  $X$  while denying everything in  $Y$

# Positions

---

$$X \vdash Y$$

# Positions

$X \not\models Y$

$$[X : Y]$$

Invalid sequents can be viewed as *positions* in a discourse

# Structural Rules

---

What do the structural rules say in terms of assertion and denial?



$$A \vdash A$$

Asserting  $A$  clashes with denying  $A$

## Structural Rules

$$\frac{X, Y \vdash Z}{X, A, Y \vdash Z} \text{ [KL]}$$

$$\frac{X \vdash Y, Z}{X \vdash Y, A, Z} \text{ [KR]}$$

If asserting  $X, Y$  clashes with denying  $Z$ , then asserting more stuff still clashes

## Structural Rules

$$\frac{X, A, AY \vdash Z}{X, A, Y \vdash Z} \text{ [WL]}$$

$$\frac{X \vdash Y, A, A, Z}{X \vdash Y, A, Z} \text{ [WR]}$$

If asserting or denying  $A$  twice results in a clash, then asserting or denying  $A$  just once results in a clash

## Structural Rules

$$\frac{X, A, B, Y \vdash Z}{X, B, A, Y \vdash Z} \text{ [CL]}$$

$$\frac{X \vdash Y, A, B, Z}{X \vdash Y, B, A, Z} \text{ [CR]}$$

If some assertions and denials clash, then asserting and denying the same things in a different order still clashes

## Structural Rules

$$\frac{X \vdash Y, A \quad A, X \vdash Y}{X \vdash Y} [\text{Cut}]$$

If asserting  $X$  and denying  $A$  and  $Y$  clashes, and asserting  $X$  and  $A$  while denying  $Y$  clashes, then asserting  $X$  and denying  $Y$

Contrapositively, if asserting  $X$  and denying  $Y$  does not clash, then either asserting  $X$  and  $A$  while denying  $Y$  does not clash or asserting  $X$  while denying  $Y$  and  $A$  does not clash

# Declaratives Are Not Enough

Belnap argued that a systematic logical treatment of language should give equal weight to imperatives and interrogatives

## 1. THE DECLARATIVE FALLACY

My thesis is simple: systematic theorists should not only stop neglecting interrogatives and imperatives, but should begin to give them equal weight with declaratives. A study of the grammar, semantics, and pragmatics of all three types of sentence is needed for every single serious program in philosophy that involves giving important attention to language.<sup>1</sup>

Attempting to understand all linguistic behavior in terms of assertions commits the *Declarative Fallacy*

The hope is that the view of sequents and logic can be extended to other speech acts

# PROOFS AND MODELS

# Models as Ideal Positions

How might *truth* enter this picture?

Models are ways of systematically elaborating finite positions into ideal, infinite positions that settle every proposition

In the propositional case, valuations are generated by ideal positions



$$[X : Y]$$

The members of  $X$  are *true* and the members of  $Y$  are *false*

$$[X : Y]$$

The members of  $X$  are *true* and the members of  $Y$  are *false*  
(relative to  $[X : Y]$ ).

## Example

$$[p \vee q, r : \neg p]$$

## Example

$$[p \vee q, r : \neg p]$$

---

$$p \vee q, r$$

---

## Example

$$[p \vee q, r : \neg p]$$

---

$$p \vee q, r \quad \textit{true}$$

---

## Example

$$[p \vee q, r : \neg p]$$

---

$$p \vee q, r \quad \textit{true}$$
$$\neg p$$

---

## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |

---

## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           |              |

---



## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           | ???          |

---

## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           | ???          |

---

DEFINITION:  $A$  is *true* at  $[X : Y]$  iff  $X \vdash A, Y$ .

DEFINITION:  $A$  is *false* at  $[X : Y]$  iff  $X, A \vdash Y$ .

## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           | <i>true</i>  |

---

DEFINITION:  $A$  is *true* at  $[X : Y]$  iff  $X \vdash A, Y$ .

DEFINITION:  $A$  is *false* at  $[X : Y]$  iff  $X, A \vdash Y$ .

## Example

$[p \vee q, r : \neg p]$

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           | <i>true</i>  |
| $p \wedge r$  |              |

DEFINITION:  $A$  is *true* at  $[X : Y]$  iff  $X \vdash A, Y$ .

DEFINITION:  $A$  is *false* at  $[X : Y]$  iff  $X, A \vdash Y$ .

## Example

$[p \vee q, r : \neg p]$

---

|               |              |
|---------------|--------------|
| $p \vee q, r$ | <i>true</i>  |
| $\neg p$      | <i>false</i> |
| $p$           | <i>true</i>  |
| $p \wedge r$  | <i>true</i>  |

---

DEFINITION:  $A$  is *true* at  $[X : Y]$  iff  $X \vdash A, Y$ .

DEFINITION:  $A$  is *false* at  $[X : Y]$  iff  $X, A \vdash Y$ .

# Classical Logic

$A \wedge B$  is true at  $[X : Y]$  iff  $A$  and  $B$  are true at  $[X : Y]$ .

$A \vee B$  is false at  $[X : Y]$  iff  $A$  and  $B$  are false at  $[X : Y]$ .

$\neg A$  is true at  $[X : Y]$  iff  $A$  is false at  $[X : Y]$ .

$\neg A$  is false at  $[X : Y]$  iff  $A$  is true at  $[X : Y]$ .

# Classical Logic

$A \wedge B$  is true at  $[X : Y]$  iff  $A$  and  $B$  are true at  $[X : Y]$ .

$A \vee B$  is false at  $[X : Y]$  iff  $A$  and  $B$  are false at  $[X : Y]$ .

$\neg A$  is true at  $[X : Y]$  iff  $A$  is false at  $[X : Y]$ .

$\neg A$  is false at  $[X : Y]$  iff  $A$  is true at  $[X : Y]$ .

---

However,  $p \wedge q$  is false at  $[ : p \wedge q]$

# Classical Logic

$A \wedge B$  is true at  $[X : Y]$  iff  $A$  and  $B$  are true at  $[X : Y]$ .

$A \vee B$  is false at  $[X : Y]$  iff  $A$  and  $B$  are false at  $[X : Y]$ .

$\neg A$  is true at  $[X : Y]$  iff  $A$  is false at  $[X : Y]$ .

$\neg A$  is false at  $[X : Y]$  iff  $A$  is true at  $[X : Y]$ .

---

However,  $p \wedge q$  is false at  $[ : p \wedge q]$  but neither  $p$  nor  $q$  is false at  $[ : p \wedge q]$  since neither  $p \vdash p \wedge q$  nor  $q \vdash p \wedge q$ .



# Classical Logic

$A \wedge B$  is true at  $[X : Y]$  iff  $A$  and  $B$  are true at  $[X : Y]$ .

$A \vee B$  is false at  $[X : Y]$  iff  $A$  and  $B$  are false at  $[X : Y]$ .

$\neg A$  is true at  $[X : Y]$  iff  $A$  is false at  $[X : Y]$ .

$\neg A$  is false at  $[X : Y]$  iff  $A$  is true at  $[X : Y]$ .

---

However,  $p \wedge q$  is false at  $[ : p \wedge q]$  but neither  $p$  nor  $q$  is false at  $[ : p \wedge q]$  since neither  $p \vdash p \wedge q$  nor  $q \vdash p \wedge q$ .

Similarly,  $r$  is neither true nor false at  $[p : q]$ .

## Extensions

**FACT:** If  $A$  is neither true nor false in  $[X : Y]$   
then both  $[X, A : Y]$  and  $[X : A, Y]$  is invalid,  
and each sequent settles  $A$  — one as *true* and the other as *false*.

## Extensions

**FACT:** If  $A$  is neither true nor false in  $[X : Y]$   
then both  $[X, A : Y]$  and  $[X : A, Y]$  is invalid,  
and each sequent settles  $A$  — one as *true* and the other as *false*.

So, if  $[X : Y]$  doesn't settle the truth of a statement  $A$ , then we can throw  $A$  in on either side, to get a more comprehensive sequent which *does* settle it.

## Extensions

**FACT:** If  $A$  is neither true nor false in  $[X : Y]$   
then both  $[X, A : Y]$  and  $[X : A, Y]$  is invalid,  
and each sequent settles  $A$  — one as *true* and the other as *false*.

So, if  $[X : Y]$  doesn't settle the truth of a statement  $A$ , then we can throw  $A$  in on either side, to get a more comprehensive sequent which *does* settle it.

In general, if  $X \not\vdash Y$  then either  $X, A \not\vdash Y$  or  $X \not\vdash A, Y$ .

## Extensions

**FACT:** If  $A$  is neither true nor false in  $[X : Y]$   
then both  $[X, A : Y]$  and  $[X : A, Y]$  is invalid,  
and each sequent settles  $A$  — one as *true* and the other as *false*.

So, if  $[X : Y]$  doesn't settle the truth of a statement  $A$ , then we can throw  $A$  in on either side, to get a more comprehensive sequent which *does* settle it.

In general, if  $X \not\vdash Y$  then either  $X, A \not\vdash Y$  or  $X \not\vdash A, Y$ .

$$\frac{X \vdash Y, A \quad A, X \vdash Y}{X \vdash Y} \text{ [Cut]}$$

# Maximal Sequents

---

# Maximal Sequents

$[X : Y]$  is finitary, where  $X$  and  $Y$  are sets (or multisets or lists ...).

# Maximal Sequents

$[X : Y]$  is finitary, where  $X$  and  $Y$  are sets (or multisets or lists ...).

A *maximal* sequent is the limit of the process of throwing in each sentence in either the left or the right hand side. You can think of it as:



# Maximal Sequents

$[X : Y]$  is finitary, where  $X$  and  $Y$  are sets (or multisets or lists ...).

A *maximal* sequent is the limit of the process of throwing in each sentence in either the left or the right hand side. You can think of it as:

- ▶ A pair  $[\mathcal{X} : \mathcal{Y}]$  of *infinite* sets, such that  $\mathcal{X} \not\vdash \mathcal{Y}$  and  $\mathcal{X} \cup \mathcal{Y}$  is the whole language.

# Maximal Sequents

$[X : Y]$  is finitary, where  $X$  and  $Y$  are sets (or multisets or lists ...).

A *maximal* sequent is the limit of the process of throwing in each sentence in either the left or the right hand side. You can think of it as:

- ▶ A pair  $[\mathcal{X} : \mathcal{Y}]$  of *infinite* sets, such that  $\mathcal{X} \not\vdash \mathcal{Y}$  and  $\mathcal{X} \cup \mathcal{Y}$  is the whole language.

FACT: Every maximal sequent makes each sentence either *true* or *false*.

# Maximal Sequents

$[X : Y]$  is finitary, where  $X$  and  $Y$  are sets (or multisets or lists ...).

A *maximal* sequent is the limit of the process of throwing in each sentence in either the left or the right hand side. You can think of it as:

- ▶ A pair  $[\mathcal{X} : \mathcal{Y}]$  of *infinite* sets, such that  $\mathcal{X} \not\vdash \mathcal{Y}$  and  $\mathcal{X} \cup \mathcal{Y}$  is the whole language.

FACT: Every maximal sequent makes each sentence either *true* or *false*.

FACT: If  $X \not\vdash Y$ , there's a *maximal*  $[\mathcal{X} : \mathcal{Y}]$  extending  $[X : Y]$ .

Assign truth values relative to *maximal* positions.

Assign truth values relative to *maximal* positions.

In a slogan, *truth value = location in a maximal position*,

## Variations

The ideal position construction handles classical logic

With some small adjustments, it can be used to provide models for intuitionistic logic

The system LJ is single-conclusion, but there is a intuitionistic sequent system that has multiple conclusions

The construction with these two systems yield *Kripke models* and *Beth models*

The hypersequent system for S5 can be used to give a similar construction

Each component of a hypersequent describes a possible world

## S5 hypersequents

$$\frac{\mathcal{H}[X \vdash Y \mid X', A \vdash Y']}{\mathcal{H}[X, \Box A \vdash Y \mid X' \vdash Y']} [\Box L]$$

$$\frac{\mathcal{H}[X \vdash Y \mid \vdash A]}{\mathcal{H}[X \vdash \Box A, Y]} [\Box R]$$

$$\frac{\mathcal{H}[X \vdash Y \mid A \vdash]}{\mathcal{H}[\Diamond A, X \vdash Y]} [\Diamond L]$$

$$\frac{\mathcal{H}[X \vdash Y \mid X' \vdash A, Y']}{\mathcal{H}[X \vdash \Diamond A, Y \mid X' \vdash Y']} [\Diamond R]$$



## Extending positions

Invalid *sequents*  $[X : Y]$

Invalid *hypersequents*  $[[X : Y], [X' : Y'], \dots]$

Invalid *sequents*  $[X : Y]$

Invalid *hypersequents*  $[[X : Y], [X' : Y'], \dots]$

Say one set of pairs  $\mathcal{H}$  extends another  $\mathcal{G}$ ,  $\mathcal{G} \preceq \mathcal{H}$ , just in case for each component  $[X : Y]$  in  $\mathcal{G}$ , there is a component  $[U : V]$  in  $\mathcal{H}$  such that  $X \subseteq U$  and  $Y \subseteq V$

Example:  $\{[p : q], [s : r]\}$  is extended by both  $\{[p, s : r, q, t]\}$  and by  $\{[p, t : q], [s : r, p]\}$

# Where are the truth values *now*?

---

## Where are the truth values *now*?

Maximal *positions*  $[\mathcal{X} : \mathcal{Y}]$

Maximal *modal positions*  $[[\mathcal{X} : \mathcal{Y}], [\mathcal{X}' : \mathcal{Y}'], \dots]$

## Where are the truth values *now*?

Maximal *positions*  $[\mathcal{X} : \mathcal{Y}]$

Maximal *modal positions*  $[[\mathcal{X} : \mathcal{Y}], [\mathcal{X}' : \mathcal{Y}'], \dots]$

A set of pairs  $\mathcal{H}$  is a *modal position* iff there is no valid hypersequent  
 $X_1 \vdash Y_1 \mid \dots \mid X_n \vdash Y_n$  extended by  $\mathcal{H}$

A modal position  $\mathcal{H}$  is *maximal* iff there is no modal position  $\mathcal{J}$  such that  $\mathcal{H} \prec \mathcal{J}$

## Building maximal modal positions

The process of expanding a modal position can add formulas to a *component* as well as adding *more components*

Some maximal modal positions, however, will contain finitely many components

The construction builds connected chunks of the S5 canonical model, taking the accessibility relation to be an equivalence relation rather than the universal relation

# Building maximal modal positions

As in the classical case, the Cut rule adds new formulas to individual components

The modal rules can extend a position with new components

## Building maximal modal positions

$$\frac{\mathcal{H}[X \vdash Y \mid X', A \vdash Y']}{\mathcal{H}[X, \Box A \vdash Y \mid X' \vdash Y']} [\Box L]$$

$$\frac{\mathcal{H}[X \vdash Y \mid \vdash A]}{\mathcal{H}[X \vdash \Box A, Y]} [\Box R]$$

If  $[[X : \Box A, Y], [X_i : Y_i]]$  isn't derivable, then  $[[X : \Box A, Y], [\vdash A], [X_i : Y_i]]$  can't be either

If the latter were derivable then the former would be by  $[\Box R]$

Similarly but for  $[\Box L]$ , if, e.g.  $[[X, \Box A : Y], [X' : Y'], [X_i : Y_i]]$  isn't derivable, then  $[[X, \Box A : Y], [X', A : Y'], [X_i : Y_i]]$  can't be



## Necessity in maximal modal positions

For a maximal modal position  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\}$ ,  
 $\Box A$  is true at  $[\mathcal{X}_i : \mathcal{Y}_i]$  iff  $A$  is true at each  $[\mathcal{X}_j : \mathcal{Y}_j]$ ,  $j \in I$

( $\Rightarrow$ ) If  $\Box A$  is true at  $[\mathcal{X}_i : \mathcal{Y}_i]$  and  $A$  were not true at some component  $[\mathcal{X} : \mathcal{Y}]$ , then since  $[\mathcal{X} : \mathcal{Y}]$  is a maximal position, we would have  $A \in \mathcal{Y}$  but  $\Box A \notin \mathcal{X}$ .  $\Box A \vdash \vdash A$  is a valid sequent (by  $[\Box L]$  from the axiom  $\vdash \vdash A \vdash A$ ), so  $[[\mathcal{X}_i : \mathcal{Y}_i], [\mathcal{X} : \mathcal{Y}]]$  would not be a position, as  $\Box A \in \mathcal{X}_i$  and  $A \in \mathcal{Y}$ , so  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\}$  isn't a position. As it is, whenever  $\Box A \in [\mathcal{X}_i : \mathcal{Y}_i]$ ,  $A$  is true at every component.

## Necessity in maximal modal positions

For a maximal modal position  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\}$ ,  
 $\Box A$  is true at  $[\mathcal{X}_i : \mathcal{Y}_i]$  iff  $A$  is true at each  $[\mathcal{X}_j : \mathcal{Y}_j]$ ,  $j \in I$

( $\Leftarrow$ ) Suppose  $\Box A$  isn't true at  $[\mathcal{X}_i : \mathcal{Y}_i]$ . So we have  $\Box A \in \mathcal{Y}_i$ . Take  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\} \cup [ : A]$ . This is a position. Suppose that it is not. Then there is a derivable hypersequent  $\vdash A \mid X \vdash Y \mid \mathcal{H}$ , where  $X \subseteq \mathcal{X}_i$ ,  $Y \subseteq \mathcal{Y}_i$  and  $\mathcal{H}$  is extended by the other components of the modal position. If that were the case, then by  $[\Box R]$ , we could derive  $X \vdash \Box A, Y \mid \mathcal{H}$ , but that is extended by the original modal position. It is, then, not valid. So,  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\} \cup [ : A]$  is a position, so it is extended by a maximal modal position, which must be  $\{[\mathcal{X}_i : \mathcal{Y}_i] : i \in I\}$ , as that is not extended by any modal positions. Therefore, for some  $j \in I$ ,  $A \in \mathcal{Y}_j$ .

## A modal position extended by a finite, maximal modal position

$\vdash \Box p \vee \Box \neg p$  is not valid

So  $[ : \Box p \vee \Box \neg p ]$  is a position

Using the rules, one obtains

$[[ : ], [ : ], [ : \Box p \vee \Box \neg p ]]$

One can then choose extensions in such a way that no additional components are needed

## A modal position extended by a finite, maximal modal position

$\vdash \Box p \vee \Box \neg p$  is not valid

So  $[ : \Box p \vee \Box \neg p ]$  is a position

Using the rules, one obtains

$[[ : ], [ : ], [ : \Box p, \Box \neg p, \Box p \vee \Box \neg p ]]$

One can then choose extensions in such a way that no additional components are needed

## A modal position extended by a finite, maximal modal position

$\vdash \Box p \vee \Box \neg p$  is not valid

So  $[ : \Box p \vee \Box \neg p ]$  is a position

Using the rules, one obtains

$[[ : p], [ : \neg p], [ : \Box p, \Box \neg p, \Box p \vee \Box \neg p]]$

One can then choose extensions in such a way that no additional components are needed

## A modal position extended by a finite, maximal modal position

$\vdash \Box p \vee \Box \neg p$  is not valid

So  $[ : \Box p \vee \Box \neg p]$  is a position

Using the rules, one obtains

$[[ : p], [p : \neg p], [ : \Box p, \Box \neg p, \Box p \vee \Box \neg p]]$

One can then choose extensions in such a way that no additional components are needed

# Maximality Facts

Each modal position can be extended to a maximal modal position

Each component of a maximal modal position is a maximal position

Each maximal modal position corresponds to a simple Kripke model:  $\Box A$  is true at  $\{[\mathcal{X}_i; \mathcal{Y}_i] : i \in I\}$  iff  $A$  is true in *every* position in the modal position

BEYOND



## Further directions

There are many directions one could go from here

One could add *quantifiers* and *predicates*

One could add axioms to obtain *theories*

# Quantifiers

$$\frac{A(a), X \vdash Y}{\forall x A(x), X \vdash Y} [\forall L]$$

$$\frac{A(y), X \vdash Y}{\exists x A(x), X \vdash Y} [\exists L]$$

$$\frac{X \vdash Y, A(y)}{X \vdash Y, \forall x A(x)} [\forall R]$$

$$\frac{X \vdash Y, A(a)}{X \vdash Y, \exists x A(x)} [\exists R]$$

In  $[\forall R]$  and  $[\exists L]$ ,  $y$  cannot occur freely in the lower sequent

These rules permit an Elimination Theorem

# Identity

$$\vdash a = a$$

$$\frac{A(s), X \vdash Y}{s = t, A(t), X \vdash Y} [= L]$$

$$\frac{X \vdash Y, A(s)}{s = t, X \vdash Y, A(t)} [= L]$$

## Identity, alternative rules

Alternative rules make it easier to eliminate Cut

$$\frac{a = a, X \vdash Y}{X \vdash Y} [= R]$$

$$\frac{s = t, A(t), A(s), X \vdash Y}{s = t, A(t), X \vdash Y} [= L]$$

One can use the Dragalin-style proof to show that Cut can be eliminated

# Truth

$$\frac{A, X \vdash Y}{T\langle A \rangle, X \vdash Y} \text{ [TL]}$$

$$\frac{X \vdash Y, A}{X \vdash Y, T\langle A \rangle} \text{ [TR]}$$

These rules are inconsistent in classical logic, so one will need to go non-classical to hang onto them

They take complex formulas to atomic formulas, which leads to complications for showing that Cut can be eliminated

# Arithmetic

Take a language with  $=, 0, ', +, \times$

$$\vdash x + 0 = x$$

$$\vdash x + y' = (x + y)'$$

$$\vdash x \times 0 = 0$$

$$\vdash x \times y' = (x \times y) + x$$

$$x' = y' \vdash x = y$$

$$0 = x' \vdash$$

$$\frac{X \vdash A(0), Y \quad X, A(x) \vdash A(x'), Y}{X \vdash A(x), Y}$$

$$\frac{X, A(x') \vdash A(x), Y \quad A(0), X \vdash Y}{A(x), X \vdash Y}$$

# Inferentialism



NUEL BELNAP

“Tonk, Plonk and Plink.”

*Analysis*, 22(6): 130–134, 1962.



NUEL BELNAP

“Declaratives Are Not Enough.”

*Philosophical Studies*, 59(1): 1–30, 1990.



JAROSLAV PEREGRIN

“An Inferentialist Approach to Semantics: Time for a New Kind of Structuralism?”

*Philosophy Compass*, 3(6): 1208–1223, 2008.



ARTHUR PRIOR

“The Runabout Inference Ticket.”

*Analysis* 21(2): 38–39, 1960.

# Positions and Models



GREG RESTALL

“Multiple Conclusions” pp. 189–205 in *Logic, Methodology and Philosophy of Science: Proceedings of the Twelfth International Congress*, edited by P. Hájek, L. Valdés-Villanueva and D. Westerståhl  
KCL Publications, 2005.



GREG RESTALL

“Truth Values and Proof Theory.”  
*Studia Logica*, 92(2):241–264, 2009.



RAYMOND SMULLYAN

*First-Order Logic*.  
Springer-Verlag, 1968.



GAISI TAKEUTI

*Proof Theory*, 2nd edition.  
Elsevier, 1987.



# And Beyond



**MICHAEL KREMER**

“Kripke and the Logic of Truth.”

*Journal of Philosophical Logic*, 17: 225–278, 1988. DOI: 10.1007/BF00247954



**SARA NEGRI AND JAN VON PLATO**

*Structural Proof Theory*

Cambridge University Press, 2002.



**GREG RESTALL**

“Assertion, Denial and Non-Classical Theories.”

In *Paraconsistency: Logic and Applications*, edited by Koji Tanaka, Francesco Berto, Edwin Mares and Francesco Paoli, pp. 81–99, 2013.



**ANNE TROELSTRA AND HELMUT SCHWICHTENBERG**

*Basic Proof Theory*, 2nd ed.

Cambridge University Press, 2000.

# THANK YOU!

<https://consequently.org/class/2016/PTPLA-NASSLLI/>

@consequently / @standefer on Twitter