Defining Rules, Proofs and Counterexamples

Greg Restall



VII WORKSHOP ON PHILOSOPHICAL LOGIC · BUENOS AIRES · AUGUST 3, 2018

My Aim

To present an account of defining rules, with the aim of explaining these rules they play a central role in analytic proofs.

My Aim

Along the way, I'll explain how Kreisel's squeezing argument helps us understand the connection between an informal notion of validity and the notions formalised in our accounts of proofs and models, and the relationship between proof-theoretic and model-theoretic analyses of logical consequence.

Outline

Positions and Bounds

Definitions

What Proofs Are & What They Do

Counterexamples & Kreisel's Squeeze

POSITIONS AND BOUNDS

Positions ...

Assertions and Denials

[X : Y]

Assertions and denials are moves in a practice.

Assertions and denials are moves in a practice.

I can deny what you assert.

Assertions and denials are moves in a practice.

I can deny what you assert.

I can retract an assertion or a denial.

Assertions and denials are moves in a practice.

I can deny what you assert.

I can retract an assertion or a denial.

I can 'try on' assertion or denial hypothetically.

Assertions and denials are moves in a practice.

I can deny what you assert.

I can retract an assertion or a denial.

I can 'try on' assertion or denial hypothetically.

They are connected to other speech acts, too, like imperatives, interrogatives, recognitives, observatives, etc.

Norms for Assertion and Denial

Assertions and denials take a *stand* (*pro* or *con*) on something.

DENIAL clashes with assertion. ASSERTION clashes with denial.

Ask: p?

Ask: p?

YES

Ask: p?

YES

Assert p

Ask: p?

YES NO

Assert p

Ask: p?

YES NO

Assert p Deny p

Ask: p?

YES NO

Assert p Deny p

These two answers clash.

Not all 'no's have the same force.

Not all 'no's have the same force.

Greg: Is Jen in the study?

Not all 'no's have the same force.

Greg: Is Jen in the study?

Lesley: No.

Not all 'no's have the same force.

Greg: Is Jen in the study?

Lesley: No.

Lesley: She's outside.

Not all 'no's have the same force.

Greg: Is Jen in the study? *Greg*: Jen is in the study.

Lesley: No.

Lesley: She's outside.

Not all 'no's have the same force.

Greg: Is Jen in the study? *Greg*: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside.

Not all 'no's have the same force.

Greg: Is Jen in the study? Greg: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside. Lesley: She's either in the study or outside.

Not all 'no's have the same force.

Greg: Is Jen in the study? Greg: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside. Lesley: She's either in the study or outside.

Strong denial

Not all 'no's have the same force.

Greg: Is Jen in the study? Greg: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside. Lesley: She's either in the study or outside.

Strong denial Weak denial

Not all 'no's have the same force.

Greg: Is Jen in the study? Greg: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside. Lesley: She's either in the study or outside.

Strong denial Weak denial

Adds p to the common ground on the negative side

Not all 'no's have the same force.

Greg: Is Jen in the study? *Greg*: Jen is in the study.

Lesley: No. Lesley: No.

Lesley: She's outside. Lesley: She's either in the study or outside.

Strong denial Weak denial

Adds p to the common ground

Retracts p from the common ground
on the positive side

on the negative side on the positive side

Weak Assertion?

Perhaps p.

Weak Assertion?

Perhaps p.

Retracts p from the common ground on the negative side?

► IDENTITY: [A : A] is out of bounds.

- ► IDENTITY: [A : A] is out of bounds.
- ▶ WEAKENING: If [X : Y] is out of bounds, then [X, A : Y] and [X : A, Y] are also out of bounds.

- ► IDENTITY: [A : A] is out of bounds.
- ▶ WEAKENING: If [X : Y] is out of bounds, then [X, A : Y] and [X : A, Y] are also out of bounds.
- ▶ CUT: If [X, A : Y] and [X : A, Y] are out of bounds, then so is [X : Y].

- ► IDENTITY: [A : A] is out of bounds.
- ▶ WEAKENING: If [X : Y] is out of bounds, then [X, A : Y] and [X : A, Y] are also out of bounds.
- ▶ CUT: If [X, A : Y] and [X : A, Y] are out of bounds, then so is [X : Y].
- ► A position that is OUT OF BOUNDS is overcommitted.

On Cut

Suppose [X : Y] is not out of bounds.

Suppose [X : Y] is not out of bounds.

Suppose [X, A : Y] is out of bounds.

Suppose [X : Y] is not out of bounds.

Suppose [X, A : Y] is out of bounds.

Ask the question: A?

Suppose [X : Y] is not out of bounds.

Suppose [X, A : Y] is out of bounds.

Ask the question: A?

The answer *no* is forced, as a *yes* answer is excluded (given our other commitments).

Structural Rules

$$A \succ A \quad Id \qquad \frac{X \succ A, Y \quad X', A \succ Y'}{X, X' \succ Y, Y'} \quad Cut$$

$$\frac{X \succ Y}{X \succ A, Y} \quad K \qquad \frac{X \succ Y}{X, A \succ Y} \quad K$$

$$\frac{X \succ A, A, Y}{X \succ A, Y} \quad W \qquad \frac{X, A, A \succ Y}{X, A \succ Y} \quad W$$

Fs
$$s >_F t$$
 $s \geqslant_F t$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u >_F u$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u \rightarrow s >_F u$ weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \rightarrow s \geqslant_F u$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u \succ s >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u >_F u >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

weak reflexivity: $\succ s \geqslant_F s$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u >_F u >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

contraries: $s >_F t$, $t \geqslant_F s >$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u >_F u >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

weak reflexivity: $\succ s \geqslant_F s$

contraries: $s >_F t$, $t \geqslant_F s >$

subcontraries: $> s >_F t$, $t \geqslant_F s$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

weak reflexivity: $\succ s \geqslant_F s$

contraries: $s >_F t, t \geqslant_F s >$

subcontraries: $\succ s >_F t$, $t \geqslant_F s$

strength: $s >_F t >_F t$

Fs
$$s >_F t$$
 $s \geqslant_F t$

strong transitivity: $s >_F t$, $t >_F u \rightarrow s >_F u$

weak transitivity: $s \geqslant_F t$, $t \geqslant_F u \succ s \geqslant_F u$

strong irreflexivity: $s >_F s >$

weak reflexivity: \gt s \geqslant_F s

contraries: $s >_F t, t \geqslant_F s >$

subcontraries: $>_F t$, $t \geqslant_F s$

strength: $s >_F t >_F t$

preservation: Fs, $t \geqslant_F s > Ft$

DEFINITIONS

How do you define a concept?

By showing people how to use it.

Define a concept by showing how you can compose that concept out of more primitive concepts.

x is a square $=_{df}$ x is a rectangle \wedge all sides of x are equal in length.

```
x is a square =_{df} x is a rectangle \wedge all sides of x are equal in length.
```

```
x is a square =_{df} x is a rectangle \wedge all sides of x are equal in length.
```

```
x is a square =_{df} x is a rectangle \wedge all sides of x are equal in length.
```

Define a concept by showing how you can compose that concept out of more primitive concepts.

```
x is a square =_{df} x is a rectangle \wedge all sides of x are equal in length.
```

Concepts defined explicitly are *sharply delimited* (contingent on people accepting the definition).

Define a concept by showing how you can compose that concept out of more primitive concepts.

```
x is a square =_{df} x is a rectangle \wedge all sides of x are equal in length.
```

Concepts defined explicitly are *sharply delimited* (contingent on people accepting the definition).

Logical concepts are similarly sharply delimited, but they cannot *all* be given explicit definitions.

Definition through a rule for use

 $[X,A\otimes B:Y] \text{ is out of bounds}$ if and only if

[X, A, B : Y] is out of bounds

Definition through a rule for use

 $[X, A \otimes B : Y]$ is out of bounds if and only if

[X, A, B : Y] is out of bounds

$$\frac{X,A,B\succ Y}{\overline{X,A\otimes B}\succ Y}\otimes \mathit{Df}$$

What about when to *deny* a conjunction?

When do we have $X > A \otimes B$, Y?

What about when to deny a conjunction?

When do we have $X > A \otimes B$, Y?

$$\frac{X' \succ B, Y'}{X \succ A, Y} \frac{\overline{A \otimes B \succ A \otimes B}}{A, B \succ A \otimes B} \stackrel{Id}{\otimes Df} \\ \frac{X \succ A, Y}{X', A \succ A \otimes B, Y'} \stackrel{Cut}{Cut}$$

What about when to deny a conjunction?

When do we have $X > A \otimes B$, Y?

$$\frac{X' \succ B, Y'}{X \succ A, Y} \frac{\overline{A \otimes B \succ A \otimes B}}{A, B \succ A \otimes B} \stackrel{Id}{\otimes Df} \\ \frac{X \succ A, Y}{X', A \succ A \otimes B, Y'} \stackrel{Cut}{Cut}$$

So, we have

$$\frac{X \succ A, Y \quad X' \succ B, Y'}{X, X' \succ A \otimes B, Y, Y'} \otimes R$$

What we've done

We have given norms governing \otimes judgements in terms of norms governing simpler judgements.

Definitions for other logical concepts

$$\frac{X \succ A, Y}{X, \neg A \succ Y} \neg Df \qquad \frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df \qquad \frac{X \succ A, B, Y}{X \succ A \oplus B, Y} \oplus Df$$

$$\frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df$$

$$X \succ A, B, Y$$

 $X \succ A \oplus B, Y$

Definitions for other logical concepts

$$\frac{X \succ A, Y}{X, \neg A \succ Y} \neg Df \qquad \frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df \qquad \frac{X \succ A, B, Y}{X \succ A \oplus B, Y} \oplus Df$$

$$\frac{X \succ A, Y \quad X \succ B, Y}{X \succ A \land B, Y} \land Df \qquad \frac{X, A \succ Y \quad X, B \succ Y}{X, A \lor B \succ Y} \lor Df$$

Definitions for other logical concepts

$$\frac{X \succ A, Y}{X, \neg A \succ Y} \neg Df \qquad \frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df \qquad \frac{X \succ A, B, Y}{X \succ A \oplus B, Y} \oplus Df$$

$$\frac{X \succ A, Y \quad X \succ B, Y}{X \succ A \land B, Y} \land Df \qquad \frac{X, A \succ Y \quad X, B \succ Y}{X, A \lor B \succ Y} \lor Df$$

$$\frac{X \succ A|_{n}^{x}, Y}{X \succ (\forall x)A, Y} \forall Df \qquad \frac{X, A|_{n}^{x} \succ Y}{X, (\exists x)A \succ Y} \forall Df \qquad \frac{X, Fs \succ Ft, Y}{X \succ s = t, Y} = Df$$

(Where n and F are not present in X and Y.)

How does this work?

How do concepts defined in this way work?

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$

 $*Df \leftrightarrow *L/R$

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$

 $*Df \leftrightarrow *L/R$

$$*L/R + Cut + Id \Leftrightarrow *L/R + Cut$$

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$

 $*Df \leftrightarrow *L/R$

$$*L/R + Cut + Id \Leftrightarrow *L/R + Cut$$
Id Elimination

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$

 $*Df \leftrightarrow *L/R$

$$*L/R + Cut + Id \Leftrightarrow *L/R + Cut$$

Id Elimination

$$*L/R + Cut \Leftrightarrow *L/R$$

Transforming Systems of Rules

$$*Df + Cut + Id \Leftrightarrow *L/R + Cut + Id$$
 $*Df \leftrightarrow *L/R$
 $*L/R + Cut + Id \Leftrightarrow *L/R + Cut$
 $Id \ Elimination$
 $*L/R + Cut \Leftrightarrow *L/R$
 $Cut \ Elimination$

► Are uniquely defined. (If you and I use the same rule, we define the same concept.)

- ► Are uniquely defined. (If you and I use the same rule, we define the same concept.)
- ► Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)

- ► Are uniquely defined. (If you and I use the same rule, we define the same concept.)
- ► Are conservatively extending. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ► Play useful dialogical roles. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)

- ► Are uniquely defined. (If you and I use the same rule, we define the same concept.)
- ► Are conservatively extending. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ Play useful dialogical roles. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ► Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)

- ► Are uniquely defined. (If you and I use the same rule, we define the same concept.)
- ► Are conservatively extending. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ Play useful dialogical roles. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ► Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)
- ▶ In Brandom's terms, they *make explicit* some of what was implicit in the practice of assertion and denial.

A Tiny Proof

If it's Thursday, I'm in Melbourne.

It's Thursday.

Therefore, I'm in Melbourne.

A Tiny Proof

If it's Thursday, I'm in Melbourne.

It's Thursday.

Therefore, I'm in Melbourne.

$$\frac{\overline{A \to B \succ A \to B}}{A \to B, A \succ B}^{\textit{Id}}$$

A Tiny Proof

If it's Thursday, I'm in Melbourne.

It's Thursday.

Therefore, I'm in Melbourne.

$$\frac{\overline{A \to B \succ A \to B}}{A \to B, A \succ B} \stackrel{\textit{Id}}{\to \textit{Df}}$$

[It's Thursday \rightarrow I'm in Melbourne, It's Thursday : I'm in Melbourne] (This is out of bounds.)

The Undeniable

Take a context in which I've asserted it's Thursday → I'm in Melbourne and I've asserted it's Thursday, then I'm in Melbourne is undeniable.

The Undeniable

Take a context in which I've asserted it's Thursday → I'm in Melbourne and I've asserted it's Thursday, then I'm in Melbourne is undeniable.

Adding the *assertion* makes explicit what was *implicit* before that assertion.

The Undeniable

Take a context in which I've asserted it's Thursday → I'm in Melbourne and I've asserted it's Thursday, then I'm in Melbourne is undeniable.

Adding the *assertion* makes explicit what was *implicit* before that assertion.

The stance (pro or con) on I'm in Melbourne was already made.

Proofs

A proof for X > Y shows that the position [X : Y] is out of bounds, by way of the defining rules for the concepts involved in the proof.

Proofs

A proof for X > Y shows that the position [X : Y] is out of bounds, by way of the defining rules for the concepts involved in the proof.

In this sense, proofs are analytic.

They apply, given the definitions, independently of the positions taken by those giving the proof.

What Proofs Prove

A proof of A, B \succ C, D can be seen as a proof of C from [A, B : D],

What Proofs Prove

A proof of A, B \succ C, D can be seen as a proof of C from [A, B : D], and a refutation of A from [B : C, D], and more.

COUNTEREXAMPLES & KREISEL'S SQUEEZE

Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \; \textit{Cut}$$

Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \; \textit{Cut}$$

If $X \succ Y$ is not derivable then one of X, $A \succ Y$ and $X \succ A \succ Y$ is also not derivable.

Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \; \textit{Cut}$$

If $X \succ Y$ is not derivable then one of X, $A \succ Y$ and $X \succ A \succ Y$ is also not derivable.

If [X : Y] is available, then so is either [X, A : Y] or [X : A, Y]

Keep Going ...

If [X : Y] is available, we can extend it into a partition [X' : Y'] of the entire language.

Keep Going ...

If [X : Y] is available, we can extend it into a partition [X' : Y'] of the entire language.

 $U \succ V$ is not derivable for any finite $U \subseteq X'$ and $V \subseteq Y'$.

Adding Witnesses

If $(\exists x)$ A is added on the left, we also add a *witness* $A|_n^x$, where n is fresh

Adding Witnesses

If $(\exists x)A$ is added on the left, we also add a *witness* $A|_n^x$, where n is fresh

$$\frac{X,A|_{n}^{x},(\exists x)A\succ Y}{X,(\exists x)A\succ Y}\,\exists \textit{Df,W}$$

Adding Witnesses

If $(\exists x)A$ is added on the left, we also add a witness $A|_n^x$, where n is fresh and similarly when $(\forall x)A$ is added on the right.

$$\frac{X,A|_{n}^{x},(\exists x)A\succ Y}{X,(\exists x)A\succ Y}\,\exists \textit{Df,W}\qquad \frac{X\succ (\forall x)A,A|_{n}^{x},Y}{X\succ (\forall x)A,Y}\,\forall \textit{Df,W}$$

$$A \in X' \text{ iff } \neg A \not\in X' \text{ iff } \neg A \in Y'$$
,

$$A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$$

$$A \lor B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$$

$$A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$$

$$(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$$

 $(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$

$$A \in X' \text{ iff } \neg A \notin X' \text{ iff } \neg A \in Y',$$

 $A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$

 $A \lor B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$

 $A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$

 $(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$

 $(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$

This is a model, where the true formulas are in X' and the false formulas are in Y', and whose domain is the set of names.

$$A \in X' \text{ iff } \neg A \notin X' \text{ iff } \neg A \in Y',$$

 $A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$

 $A \lor B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$

 $A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$

 $(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$

 $(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$

This is a model, where the true formulas are in X' and the false formulas are in Y', and whose domain is the set of names.

(Things are *little* more delicate when the language contains the identity predicate.)

Soundness and Completeness

X ≻ Y is derivable
 iff there is no model
 in which each member of X is true
 and each member of Y is false.

Kreisel's Squeeze

X > Y is informally valid

Kreisel's Squeeze

 $X \succ Y$ has a derivation $\downarrow \downarrow$ $X \succ Y$ is informally valid

Kreisel's Squeeze

Kreisel's Squeeze

$$X \succ Y$$
 has a derivation $\downarrow \downarrow$ $X \succ Y$ is informally valid $\downarrow \downarrow$ $X \succ Y$ has no countermodel $\downarrow \downarrow$ $X \succ Y$ has a derivation.

► To say that X > Y is *informally valid* means that is a clash involved in asserting each member of X and denying each member of Y.

- ► To say that X > Y is *informally valid* means that is a clash involved in asserting each member of X and denying each member of Y.
- ▶ Axiomatic sequents (A > A) are informally valid in this sense.

- ► To say that X > Y is *informally valid* means that is a clash involved in asserting each member of X and denying each member of Y.
- ▶ Axiomatic sequents (A > A) are informally valid in this sense.
- ► Structural rules preserve informal validity.

- ► To say that X > Y is *informally valid* means that is a clash involved in asserting each member of X and denying each member of Y.
- ▶ Axiomatic sequents $(A \succ A)$ are informally valid in this sense.
- ► Structural rules preserve informal validity.
- ▶ Defining rules *define* the connectives/quantifiers.



► Refine our notion of informal validity: *Literals* (Fa, Gbc, etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.

- ► Refine our notion of informal validity: *Literals* (Fa, Gbc, etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ► Given a witnessed partition position [X : Y] (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in X and denying any in Y.

- ► Refine our notion of informal validity: *Literals* (Fa, Gbc, etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ► Given a witnessed partition position [X : Y] (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in X and denying any in Y.
- ► So, there is no clash involved in asserting *any* formulas in X and denying any formulas in Y, by appeal to the defining rules.

- ► Refine our notion of informal validity: *Literals* (Fa, Gbc, etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ► Given a witnessed partition position [X : Y] (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in X and denying any in Y.
- ▶ So, there is no clash involved in asserting *any* formulas in X and denying any formulas in Y, by appeal to the defining rules. (This is an induction on the depth of the structure of the formulas. The defining rules reduce clashes involving formulas into clashes involving subformulas.)

- ► Refine our notion of informal validity: *Literals* (Fa, Gbc, etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ► Given a witnessed partition position [X : Y] (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in X and denying any in Y.
- ▶ So, there is no clash involved in asserting *any* formulas in X and denying any formulas in Y, by appeal to the defining rules. (This is an induction on the depth of the structure of the formulas. The defining rules reduce clashes involving formulas into clashes involving subformulas.)
- ► So, a countermodel for a sequent shows *how* there is no clash involved in asserting each member of X and denying each member of Y.

(3) From Absence of Countermodel to Derivability

That's the Completeness Theorem.

Kreisel's Squeeze

The Result

Informal validity (in virtue of first order logical form), for the language given by the defining rules, is *first order classical logic*, as given by the sequent calculus and Tarski's models.

THANK YOU!

https://consequently.org/presentation/2018/defining-rules-proofs-and-counterexamples-ba-logic-vii/

@consequently on Twitter