

Isomorphisms in a Category of Propositions and Proofs

Greg Restall



THE UNIVERSITY OF
MELBOURNE

MELBOURNE LOGIC GROUP · 2 MARCH 2018

To show how a category of propositions
and *classical* proofs can give rise to
finely grained hyperintensional notions
of sameness of content.

One notion is *very* finely grained
(distinguishing p and $p \wedge p$)
others are is less finely grained.

One of these notions amounts to equivalence in
R. B. Angell's logic of analytic containment.

To apply distinctively
prooftheoretical methods
to issues in philosophical logic.

Acknowledgements

Thanks to
Rohan French,
Dave Ripley, and
Shawn Standefer for
helpful conversations
on this material.

The Category of Classical Proofs

Isomorphisms

More Proofs from A to A

Matching & Logics of Analytic Containment

Matching as Isomorphism

THE CATEGORY OF CLASSICAL PROOFS

There can be different ways to prove the same thing

$$p \wedge q \succ p \vee q$$

Four different derivations,

$$\frac{\frac{p \succ p}{p \wedge q \succ p} \wedge L}{p \wedge q \succ p \vee q} \vee R$$

$$\frac{\frac{p \succ p}{p \succ p \vee q} \vee R}{p \wedge q \succ p \vee q} \wedge L$$

$$\frac{\frac{q \succ q}{p \wedge q \succ q} \wedge L}{p \wedge q \succ p \vee q} \vee R$$

$$\frac{\frac{q \succ q}{q \succ p \vee q} \vee R}{p \wedge q \succ p \vee q} \wedge L$$

Four different derivations, two *proofs*

$$\frac{\frac{p \succ p}{p \wedge q \succ p} \wedge L}{p \wedge q \succ p \vee q} \vee R \approx \frac{p \wedge q}{p} \approx \frac{\frac{p \succ p}{p \succ p \vee q} \vee R}{p \wedge q \succ p \vee q} \wedge L$$

$$\frac{\frac{q \succ q}{p \wedge q \succ q} \wedge L}{p \wedge q \succ p \vee q} \vee R \approx \frac{p \wedge q}{q} \approx \frac{\frac{q \succ q}{q \succ p \vee q} \vee R}{p \wedge q \succ p \vee q} \wedge L$$

Motivating Idea

Proof terms are an *invariant*
for derivations under rule permutation.

δ_1 and δ_2 have the same *term* iff
some permutation sends δ_1 to δ_2 .

Four different derivations, two *proof terms*

$$\begin{array}{c}
 \frac{x \curvearrowright y}{x : p \supset y : p} \wedge L \\
 \frac{x : p \wedge q \supset y : p}{\wedge x \curvearrowright \vee y} \vee R \\
 x : p \wedge q \supset y : p \vee q
 \end{array}$$

$$\wedge x \curvearrowright \vee y$$

$$\begin{array}{c}
 \frac{x \curvearrowright x}{x : p \supset y : p} \vee R \\
 \frac{x \curvearrowright \vee y}{x : p \supset y : p \vee q} \wedge L \\
 x : p \wedge q \supset y : p \vee q
 \end{array}$$

$$\begin{array}{c}
 \frac{x \curvearrowright y}{x : q \supset y : q} \wedge L \\
 \frac{x : p \wedge q \supset y : q}{\lambda x \curvearrowright \vee y} \vee R \\
 x : p \wedge q \supset y : p \vee q
 \end{array}$$

$$\lambda x \curvearrowright \vee y$$

$$\begin{array}{c}
 \frac{x \curvearrowright y}{x : q \supset y : q} \vee R \\
 \frac{x \curvearrowright \vee y}{x : q \supset y : p \vee q} \wedge L \\
 x : p \wedge q \supset y : p \vee q
 \end{array}$$

Ingredients

λ terms • flow graphs • proof nets


A proof term for $\Sigma \succ \Delta$
encodes the flow of information
in a proof of $\Sigma \succ \Delta$.

Proof Terms


$$\begin{array}{l} \lambda x \rightarrow \lambda y \quad \lambda x \rightarrow \lambda y \quad \lambda x \rightarrow \lambda y \quad \lambda x \rightarrow \lambda y \\ x : p \wedge (q \vee r) \succ y : (p \wedge q) \vee (p \wedge r) \end{array}$$

Proof Terms as Graphs on Sequents

$\lambda x \curvearrowright \lambda \vee y \quad \lambda x \curvearrowright \lambda \vee y \quad \vee \lambda x \curvearrowright \lambda \vee y \quad \vee \lambda x \curvearrowright \lambda \vee y$
 $x : p \wedge (q \vee r) \succ y : (p \wedge q) \vee (p \wedge r)$



Finding a Proof Term from a Derivation



Finding a Proof Term from a Derivation

$$\begin{array}{c}
 \begin{array}{c} p \succ p \quad q \succ q \\ \hline p, q \succ p \wedge q \end{array} \wedge R \quad \begin{array}{c} p \succ p \quad r \succ r \\ \hline p, r \succ p \wedge r \end{array} \wedge R \\
 \hline p, q \vee r \succ p \wedge q, p \wedge r \vee L \\
 \hline p, q \vee r \succ (p \wedge q) \vee (p \wedge r) \vee R \\
 \hline p \wedge (q \vee r) \succ (p \wedge q) \vee (p \wedge r) \wedge L
 \end{array}$$

$$\begin{array}{c}
 p \wedge (q \vee r) \\
 \swarrow \quad \searrow \quad \downarrow \\
 (p \wedge q) \vee (p \wedge r)
 \end{array}$$

Finding a Proof Term from a Derivation

$$\begin{array}{c}
 \frac{p \succ p \quad q \succ q}{p, q \succ p \wedge q} \wedge R \qquad \frac{p \succ p \quad r \succ r}{p, r \succ p \wedge r} \wedge R \\
 \frac{\quad}{p, q \vee r \succ p \wedge q, p \wedge r} \vee L \\
 \frac{\quad}{p, q \vee r \succ (p \wedge q) \vee (p \wedge r)} \vee R \\
 \frac{\quad}{p \wedge (q \vee r) \succ (p \wedge q) \vee (p \wedge r)} \wedge L
 \end{array}$$

$$\begin{array}{c}
 p \wedge (q \vee r) \\
 \swarrow \quad \searrow \\
 (p \wedge q) \vee (p \wedge r)
 \end{array}$$

Finding a Proof Term from a Derivation

$$\begin{array}{c}
 \frac{p \succ p \quad q \succ q}{p, q \succ p \wedge q} \wedge R \qquad \frac{p \succ p \quad r \succ r}{p, r \succ p \wedge r} \wedge R \\
 \frac{\frac{p, q \succ p \wedge q \quad p, r \succ p \wedge r}{p, q \vee r \succ p \wedge q, p \wedge r} \vee L}{p, q \vee r \succ (p \wedge q) \vee (p \wedge r)} \vee R \\
 \frac{p, q \vee r \succ (p \wedge q) \vee (p \wedge r)}{p \wedge (q \vee r) \succ (p \wedge q) \vee (p \wedge r)} \wedge L
 \end{array}$$

$$\begin{array}{c}
 p \wedge (q \vee r) \\
 \downarrow \\
 (p \wedge q) \vee (p \wedge r)
 \end{array}$$

Finding a Proof Term from a Derivation

$$\begin{array}{c}
 \frac{p \succ p \quad q \succ q}{p, q \succ p \wedge q} \wedge R \qquad \frac{p \succ p \quad r \succ r}{p, r \succ p \wedge r} \wedge R \\
 \hline
 \frac{\qquad p, q \vee r \succ p \wedge q, p \wedge r}{p, q \vee r \succ (p \wedge q) \vee (p \wedge r)} \vee L \\
 \hline
 \frac{\qquad p, q \vee r \succ (p \wedge q) \vee (p \wedge r)}{p \wedge (q \vee r) \succ (p \wedge q) \vee (p \wedge r)} \wedge L
 \end{array}$$

$$p \wedge (q \vee r)$$




$$(p \wedge q) \vee (p \wedge r)$$

Finding a Proof Term from a Derivation




$$\begin{array}{c}
 \begin{array}{c} p \succ p \quad q \succ q \\ \hline p, q \succ p \wedge q \end{array} \wedge R \quad \begin{array}{c} p \succ p \quad r \succ r \\ \hline p, r \succ p \wedge r \end{array} \wedge R \\
 \hline p, q \vee r \succ p \wedge q, p \wedge r \vee L \\
 \hline p, q \vee r \succ (p \wedge q) \vee (p \wedge r) \vee R \\
 \hline p \wedge (q \vee r) \succ (p \wedge q) \vee (p \wedge r) \wedge L
 \end{array}$$

$$\begin{array}{c}
 p \wedge (q \vee r) \\
 \swarrow \quad \searrow \quad \downarrow \\
 (p \wedge q) \vee (p \wedge r)
 \end{array}$$

More Flow Graphs



More Flow Graphs



Links wholly internal to a *premise* or a *conclusion* are called *cups* (↪) and *caps* (↩).

Proof Term Facts

Not every directed graph on occurrences of atoms in a sequent is a proof term.

Proof Term Facts

Not every directed graph on occurrences of atoms in a sequent is a proof term.

- ▶ They *typecheck*. [An occurrence of p is linked only with an occurrence of p .]

Proof Term Facts

Not every directed graph on occurrences of atoms in a sequent is a proof term.


- ▶ They *typecheck*. [An occurrence of p is linked only with an occurrence of p .]
- ▶ They *respect polarities*. [Positive occurrences of atoms in premises and negative occurrences of atoms in conclusions are *inputs*. Negative occurrences of atoms in premises and positive occurrences of atoms in conclusions are *outputs*.]

Proof Term Facts


Not every directed graph on occurrences of atoms in a sequent is a proof term.

- ▶ They *typecheck*. [An occurrence of p is linked only with an occurrence of p .]
- ▶ They *respect polarities*. [Positive occurrences of atoms in premises and negative occurrences of atoms in conclusions are *inputs*. Negative occurrences of atoms in premises and positive occurrences of atoms in conclusions are *outputs*.]
- ▶ They must satisfy an “enough connections” condition, amounting to a non-emptiness under every *switching*. [e.g. the obvious linking between premise $p \vee q$ and conclusion $p \wedge q$ is not connected enough to be a proof term.]


Cut is chaining of proof terms



Cut is chaining of proof terms




Cut is chaining of proof terms




The *cut formula* is no longer a premise or a conclusion in the proof term.


Eliminating Cuts is Local





Eliminating Cuts is Local



Eliminating Cuts is Local



Eliminating Cuts is Local




- ▶ Cut elimination is *confluent* and *terminating*.

- ▶ Cut elimination is *confluent* and *terminating*.
[So it can be understood as a kind of *evaluation*.]



- ▶ Cut elimination is *confluent* and *terminating*.
[So it can be understood as a kind of *evaluation*.]
- ▶ Cut elimination for proof terms is *local*.

- ▶ Cut elimination is *confluent* and *terminating*.
[So it can be understood as a kind of *evaluation*.]
- ▶ Cut elimination for proof terms is *local*.
[So it is easily made parallel.]

Cuts with Caps and Cups



Cuts with Caps and Cups



\mathcal{C} is the Category of Classical Proofs

OBJECTS Formulas — A, B , etc.

ARROWS Cut-Free Proof Terms — $\pi : A \multimap B$.

COMPOSITION Composition of derivations with the elimination of *Cut* — If $\pi : A \multimap B$ and $\tau : B \multimap C$ then $\tau \circ \pi : A \multimap C$.

IDENTITY Canonical identity proofs — $Id(A) : A \multimap A$.

Identity Proofs

$$\begin{array}{c}
\frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
\frac{}{\neg p \succ \neg p} \neg R \qquad \frac{}{p \supset p \succ p \supset p} \supset R \\
\hline
\neg p \vee (p \supset p) \succ \neg p, p \supset p \qquad \qquad \qquad \vee L \\
\hline
\neg p \vee (p \supset p) \succ \neg p \vee (p \supset p) \qquad \qquad \qquad \vee R
\end{array}$$

$$\neg p \vee (p \supset p)$$

$$\neg p \vee (p \supset p)$$

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{\neg p \succ \neg p}{p \supset p \succ p \supset p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \frac{\neg p \succ \neg p}{\neg p \vee (p \supset p) \succ \neg p, p \supset p} \vee L \\
 \frac{\neg p \vee (p \supset p) \succ \neg p, p \supset p}{\neg p \vee (p \supset p) \succ \neg p \vee (p \supset p)} \vee R
 \end{array}$$

$$\neg p \vee (p \supset p)$$

$$\neg p \vee (p \supset p)$$

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{p, \neg p \succ}{\neg p \succ \neg p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \frac{\neg p \succ \neg p \quad p \supset p \succ p \supset p}{\neg p \vee (p \supset p) \succ \neg p, p \supset p} \vee L \\
 \frac{\neg p \vee (p \supset p) \succ \neg p, p \supset p}{\neg p \vee (p \supset p) \succ \neg p \vee (p \supset p)} \vee R
 \end{array}$$

$$\begin{array}{c}
 \neg p \vee (p \supset p) \\
 \uparrow \\
 \neg p \vee (p \supset p)
 \end{array}$$

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{\neg p \succ \neg p}{p \supset p \succ p \supset p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \frac{\neg p \supset (p \supset p) \succ \neg p, p \supset p}{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)} \vee L \\
 \frac{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)}{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)} \vee R
 \end{array}$$

$$\begin{array}{c}
 \neg p \vee (p \supset p) \\
 \uparrow \\
 \neg p \vee (p \supset p)
 \end{array}$$

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{\neg p \succ \neg p}{p \supset p \succ p \supset p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \frac{\neg p \supset (p \supset p) \succ \neg p, p \supset p}{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)} \vee L \\
 \frac{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)}{\neg p \supset (p \supset p) \succ \neg p \supset (p \supset p)} \vee R
 \end{array}$$

Red arrows indicate the flow of information: from the top-left $p \succ p$ to the $\neg L$ rule, then to the $\neg R$ rule, then to the $\vee L$ rule, and finally to the $\vee R$ rule. A curved red arrow also connects the two $p \succ p$ assumptions at the top to the $\supset L$ rule.

$$\begin{array}{c}
 \neg p \vee (p \supset p) \\
 \uparrow \qquad \uparrow \\
 \neg p \vee (p \supset p)
 \end{array}$$

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{\neg p \succ \neg p}{\neg p \succ \neg p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \hline
 \frac{\neg p \vee (p \supset p) \succ \neg p, p \supset p}{\neg p \vee (p \supset p) \succ \neg p \vee (p \supset p)} \vee L \\
 \hline
 \neg p \vee (p \supset p) \succ \neg p \vee (p \supset p) \quad \vee R
 \end{array}$$

Red arrows indicate the flow of information: from the top-left $p \succ p$ to the $\neg L$ rule, from the top-right $p \succ p$ to the $\supset L$ rule, from the bottom-left $\neg p \vee (p \supset p)$ to the $\vee L$ rule, and from the bottom-right $\neg p \vee (p \supset p)$ to the $\vee R$ rule.

$$\begin{array}{c}
 \neg p \vee (p \supset p) \\
 \uparrow \qquad \uparrow \\
 \neg p \vee (p \supset p)
 \end{array}$$

Red arrows indicate the flow of information: from the bottom $\neg p$ to the top $\neg p$, and from the bottom $p \supset p$ to the top $p \supset p$.

Identity Proofs

$$\begin{array}{c}
 \frac{p \succ p}{p, \neg p \succ} \neg L \qquad \frac{p \succ p \quad p \succ p}{p \supset p, p \succ p} \supset L \\
 \frac{\neg p \succ \neg p}{\neg p \succ \neg p} \neg R \qquad \frac{p \supset p, p \succ p}{p \supset p \succ p \supset p} \supset R \\
 \hline
 \frac{\neg p \vee (p \supset p) \succ \neg p, p \supset p}{\neg p \vee (p \supset p) \succ \neg p \vee (p \supset p)} \vee L \\
 \hline
 \neg p \vee (p \supset p) \succ \neg p \vee (p \supset p) \quad \vee R
 \end{array}$$

Red arrows indicate the flow of information: from the top-left $p \succ p$ to the $\neg L$ rule, from the top-right $p \succ p$ to the $\supset L$ rule, from the bottom-left $\neg p \vee (p \supset p)$ to the $\vee L$ rule, and from the bottom-right $\neg p \vee (p \supset p)$ to the $\vee R$ rule.

$$\begin{array}{c}
 \neg p \vee (p \supset p) \\
 \uparrow \qquad \uparrow \qquad \downarrow \\
 \neg p \vee (p \supset p)
 \end{array}$$



Red arrows indicate the flow of information: from the bottom-left $\neg p$ to the top-left $\neg p$, from the bottom-middle p to the top-middle p , and from the bottom-right p to the top-right p .

Identity Proofs



In the identity proof from A to A ,

- ▶ A *positive* occurrence of an atom in the premise linked *to* its mate in the conclusion.
- ▶ A *negative* occurrence of an atom in the premise is linked *from* its mate in the conclusion.
- ▶ There are no other links.



Identity and Composition in \mathcal{C}



Identity and Composition in \mathcal{C}



Identity and Composition in \mathcal{C}



The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

Being enriched in *SLat* means that proofs terms come ordered by \sqsubseteq , and compose under \cup , and these interact sensibly with composition.

The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

Being enriched in *SLat* means that proofs/terms come ordered by \sqsubseteq , and compose under \cup , and these interact sensibly with composition.

$$\pi \sqsubseteq \pi' \Rightarrow \pi \circ \tau \sqsubseteq \pi' \circ \tau$$

The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

Being enriched in *SLat* means that proofs/terms come ordered by \sqsubseteq , and compose under \cup , and these interact sensibly with composition.

$$\begin{aligned}\pi \sqsubseteq \pi' &\Rightarrow \pi \circ \tau \sqsubseteq \pi' \circ \tau \\ \tau \sqsubseteq \tau' &\Rightarrow \pi \circ \tau \sqsubseteq \pi \circ \tau'\end{aligned}$$

The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

Being enriched in *SLat* means that proofs/terms come ordered by \sqsubseteq , and compose under \cup , and these interact sensibly with composition.

$$\pi \sqsubseteq \pi' \Rightarrow \pi \circ \tau \sqsubseteq \pi' \circ \tau$$

$$\tau \sqsubseteq \tau' \Rightarrow \pi \circ \tau \sqsubseteq \pi \circ \tau'$$

$$\pi \circ (\tau \cup \tau') = (\pi \circ \tau) \cup (\pi \circ \tau')$$

The Category \mathcal{C} ...

- ▶ ... is *symmetric monoidal* and *star autonomous*
- ▶ but not *Cartesian*,
- ▶ with structural *monoids* and *comonoids*,
- ▶ and is enriched in *SLat* (the category of semilattices).

Being enriched in *SLat* means that proofs/terms come ordered by \sqsubseteq , and compose under \cup , and these interact sensibly with composition.

$$\pi \sqsubseteq \pi' \Rightarrow \pi \circ \tau \sqsubseteq \pi' \circ \tau$$

$$\tau \sqsubseteq \tau' \Rightarrow \pi \circ \tau \sqsubseteq \pi \circ \tau'$$

$$\pi \circ (\tau \cup \tau') = (\pi \circ \tau) \cup (\pi \circ \tau')$$

$$(\pi \cup \pi') \circ \tau = (\pi \circ \tau) \cup (\pi' \circ \tau)$$

\mathcal{C} is just *classical* propositional logic,
in a categorical setting.

\mathcal{C} is just *classical* propositional logic,
in a categorical setting.

(The sequent calculus plays no essential role here.
You can define proof terms on other proof systems,
e.g. *natural deduction*, *Hilbert proofs*, *tableaux*, *resolution*.)

ISOMORPHISMS

Isomorphisms in Categories

$f : A \rightarrow B$ is an *isomorphism* in a category iff
it has an *inverse* $g : B \rightarrow A$, where
 $g \circ f = id_A : A \rightarrow A$ and $f \circ g = id_B : B \rightarrow B$.

Isomorphisms in Categories

$f : A \rightarrow B$ is an *isomorphism* in a category iff
it has an *inverse* $g : B \rightarrow A$, where
 $g \circ f = id_A : A \rightarrow A$ and $f \circ g = id_B : B \rightarrow B$.

If g and g' are both inverses, we have
 $g = id_A \circ g = (g' \circ f) \circ g = g' \circ (f \circ g) = g' \circ id_B = g'$,
so any inverse is unique. We can call it f^{-1} .

Why Isomorphisms?

If A and B are isomorphic in a category \mathcal{C} ,
then what we can do with A (in \mathcal{C})
we can do with B , too.

Why Isomorphisms?

If A and B are isomorphic in a category \mathcal{C} ,
then what we can do with A (in \mathcal{C})
we can do with B , too.

If A and B are isomorphic in \mathfrak{C} , then
they agree not only on *provability*,
but also, on *proofs*.

Why Isomorphisms?

If A and B are isomorphic in a category \mathcal{C} ,
then what we can do with A (in \mathcal{C})
we can do with B , too.

If A and B are isomorphic in \mathfrak{C} , then
they agree not only on *provability*,
but also, on *proofs*.

The distinctions drawn when you analyse how something is *proved*
(from premises), are not far from what you want to understand
when you ask how something is *made true*.

Isomorphisms in \mathcal{C}

$$p \wedge q \cong q \wedge p$$

Isomorphisms in \mathcal{C}

$$p \wedge q \cong q \wedge p$$



Isomorphisms in \mathcal{C}

$$p \wedge q \cong q \wedge p$$



Isomorphisms in \mathcal{C}

$$p \vee q \cong q \vee p$$

Isomorphisms in \mathcal{C}

$$p \vee q \cong q \vee p$$



Isomorphisms in \mathcal{C}

$$p \vee q \cong q \vee p$$




Isomorphisms in \mathcal{C}

$$p \wedge (q \wedge r) \cong (p \wedge q) \wedge r$$



Isomorphisms in \mathcal{C}

$$p \wedge (q \wedge r) \cong (p \wedge q) \wedge r$$



Isomorphisms in \mathcal{C}

$$p \wedge (q \wedge r) \cong (p \wedge q) \wedge r$$




Isomorphisms in \mathcal{C}

$$\neg(p \wedge q) \cong \neg p \vee \neg q$$



Isomorphisms in \mathcal{C}

$$\neg(p \wedge q) \cong \neg p \vee \neg q$$



Isomorphisms in \mathcal{C}

$$\neg(p \wedge q) \cong \neg p \vee \neg q$$




Isomorphisms in \mathcal{C}

$$\neg\neg p \cong p$$


Isomorphisms in \mathcal{C}

$$\neg\neg p \cong p$$



Isomorphisms in \mathcal{C}

$$\neg\neg p \cong p$$




Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$

Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$


$$p \wedge (q \vee \neg q)$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$

$$p \wedge (q \vee \neg q)$$



$$p \wedge (q \vee \neg q)$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$

$$p \wedge (q \vee \neg q)$$



$$p \wedge (q \vee \neg q)$$





$$p \wedge (q \vee \neg q)$$

$$p \wedge (q \vee \neg q)$$





Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$





Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee \neg q) \not\cong p$$



Non-isomorphisms in \mathcal{C}

$$p \wedge p \not\cong p$$

Non-isomorphisms in \mathcal{C}


$$p \wedge p \not\cong p$$

$$p \wedge p$$




Non-isomorphisms in \mathcal{C}

$$p \wedge p \not\cong p$$




Non-isomorphisms in \mathcal{C}

$$p \wedge p \not\cong p$$



Non-isomorphisms in \mathcal{C}

$$p \wedge p \not\cong p$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee r) \not\cong (p \wedge q) \vee (p \wedge r)$$

Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee r) \not\cong (p \wedge q) \vee (p \wedge r)$$


$$(p \wedge q) \vee (p \wedge r)$$

$$p \wedge (q \vee r)$$

$$(p \wedge q) \vee (p \wedge r)$$

Non-isomorphisms in \mathcal{C}


$$p \wedge (q \vee r) \not\cong (p \wedge q) \vee (p \wedge r)$$



$$(p \wedge q) \vee (p \wedge r)$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee r) \not\cong (p \wedge q) \vee (p \wedge r)$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (q \vee r) \not\cong (p \wedge q) \vee (p \wedge r)$$



Non-isomorphisms in \mathcal{C}

$$p \wedge (p \vee q) \not\cong p \vee (p \wedge q)$$

Occurrence Polarity Condition

If A is isomorphic to B in \mathfrak{C}
then each variable occurs positively [negatively] in A
exactly as many times as it occurs positively [negatively] in B .

Occurrence Polarity Condition

If A is isomorphic to B in \mathfrak{C}
then each variable occurs positively [negatively] in A
exactly as many times as it occurs positively [negatively] in B .

(This condition is *necessary*, not *sufficient*: $p \wedge (p \vee q) \not\cong p \vee (p \wedge q)$.)

Characterising Isomorphisms

A is isomorphic to B
iff *A* and *B* are equivalent
in the following calculus:

$$A \wedge B \leftrightarrow B \wedge A, \quad A \wedge (B \wedge C) \leftrightarrow (A \wedge B) \wedge C.$$

$$A \vee B \leftrightarrow B \vee A, \quad A \vee (B \vee C) \leftrightarrow (A \vee B) \vee C.$$

$$\neg(A \vee B) \leftrightarrow \neg A \vee \neg B, \quad \neg(A \wedge B) \leftrightarrow \neg A \vee \neg B.$$

$$\neg\neg A \leftrightarrow A. \quad A \leftrightarrow B \Rightarrow C(A) \leftrightarrow C(B).$$

Characterising Isomorphisms

A is *isomorphic* to B
iff A and B are equivalent
in the following calculus:

$$\begin{aligned}A \wedge B &\leftrightarrow B \wedge A, & A \wedge (B \wedge C) &\leftrightarrow (A \wedge B) \wedge C. \\A \vee B &\leftrightarrow B \vee A, & A \vee (B \vee C) &\leftrightarrow (A \vee B) \vee C. \\ \neg(A \vee B) &\leftrightarrow \neg A \vee \neg B, & \neg(A \wedge B) &\leftrightarrow \neg A \wedge \neg B. \\ \neg\neg A &\leftrightarrow A. & A \leftrightarrow B &\Rightarrow C(A) \leftrightarrow C(B).\end{aligned}$$

This allows for a *negation normal form*, but not DNF or CNF.

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.
- ▶ A is isomorphic to B iff there are *diversified* A' and B' where A' and B' are isomorphic, and $A = \sigma A'$ and $B = \sigma B'$ for some substitution σ .

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.
- ▶ A is isomorphic to B iff there are *diversified* A' and B' where A' and B' are isomorphic, and $A = \sigma A'$ and $B = \sigma B'$ for some substitution σ .
- ▶ A is isomorphic to B iff their negation normal forms are isomorphic. (If A is diversified, so is its negation normal form.)

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.
- ▶ A is isomorphic to B iff there are *diversified* A' and B' where A' and B' are isomorphic, and $A = \sigma A'$ and $B = \sigma B'$ for some substitution σ .
- ▶ A is isomorphic to B iff their negation normal forms are isomorphic. (If A is diversified, so is its negation normal form.)
- ▶ If A and B are diversified, isomorphic, and in negation normal form, if $l \wedge m$ is a conjunction in A (l and m , literals), a substitution argument (substituting \top and \perp for the *other* atoms) shows that l and m must be conjunctively joined in B , too. The same goes for $l \vee m$.

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.
- ▶ A is isomorphic to B iff there are *diversified* A' and B' where A' and B' are isomorphic, and $A = \sigma A'$ and $B = \sigma B'$ for some substitution σ .
- ▶ A is isomorphic to B iff their negation normal forms are isomorphic. (If A is diversified, so is its negation normal form.)
- ▶ If A and B are diversified, isomorphic, and in negation normal form, if $l \wedge m$ is a conjunction in A (l and m , literals), a substitution argument (substituting \top and \perp for the *other* atoms) shows that l and m must be conjunctively joined in B , too. The same goes for $l \vee m$.
- ▶ Replace $l \wedge m$ by a new atom in both A and B , and repeat.

Proof Sketch (Došen and Petrić, 2012)

- ▶ If $A \leftrightarrow B$ holds in the calculus, A and B are isomorphic.
- ▶ A is isomorphic to B iff there are *diversified* A' and B' where A' and B' are isomorphic, and $A = \sigma A'$ and $B = \sigma B'$ for some substitution σ .
- ▶ A is isomorphic to B iff their negation normal forms are isomorphic. (If A is diversified, so is its negation normal form.)
- ▶ If A and B are diversified, isomorphic, and in negation normal form, if $l \wedge m$ is a conjunction in A (l and m , literals), a substitution argument (substituting \top and \perp for the *other* atoms) shows that l and m must be conjunctively joined in B , too. The same goes for $l \vee m$.
- ▶ Replace $l \wedge m$ by a new atom in both A and B , and repeat.
- ▶ This shows how to reconstruct a proof of equivalence for A and B in the syntactic calculus for isomorphic formulas.

This is a *very* tight constraint

- ▶ If A and B are isomorphic, they can play *essentially* the same role in proof.

This is a *very* tight constraint

- ▶ If A and B are isomorphic, they can play *essentially* the same role in proof.
- ▶ Replacing A by B in a proof (as a premise or conclusion) not only gives you something that is also a proof (mere logical equivalence would do *that*), but it gives you a proof which is *essentially the same*.

This is a *very* tight constraint

- ▶ If A and B are isomorphic, they can play *essentially* the same role in proof.
- ▶ Replacing A by B in a proof (as a premise or conclusion) not only gives you something that is also a proof (mere logical equivalence would do *that*), but it gives you a proof which is *essentially the same*.
- ▶ Not even A and $A \wedge A$ are equivalent in *this* sense.

This is a *very* tight constraint


- ▶ If A and B are isomorphic, they can play *essentially* the same role in proof.
- ▶ Replacing A by B in a proof (as a premise or conclusion) not only gives you something that is also a proof (mere logical equivalence would do *that*), but it gives you a proof which is *essentially the same*.
- ▶ Not even A and $A \wedge A$ are equivalent in *this* sense.
- ▶ Yet, A and $A \wedge A$ seem to have identical *subject matter* (insofar as we understand that notion).

This is a *very* tight constraint

- ▶ If A and B are isomorphic, they can play *essentially* the same role in proof.
- ▶ Replacing A by B in a proof (as a premise or conclusion) not only gives you something that is also a proof (mere logical equivalence would do *that*), but it gives you a proof which is *essentially the same*.
- ▶ Not even A and $A \wedge A$ are equivalent in *this* sense.
- ▶ Yet, A and $A \wedge A$ seem to have identical *subject matter* (insofar as we understand that notion).
- ▶ Can we use the fine-grained tools of proof theoretical analysis to generalise this notion of subject matter to arbitrary statements?

MORE PROOFS
FROM A TO A


$$Id(p \vee (p \wedge \neg p))$$



In $Id(A)$, each occurrence of an atom in the premise is linked with every its corresponding occurrence of in the conclusion.
Different occurrences of atoms in A are treated differently.

$Id(A), Hz(A), Mx(A)$

$Hz(p \vee (p \wedge \neg p))$




In $Id(A)$, each occurrence of an atom in the premise is linked with every its corresponding occurrence of in the conclusion.
Different occurrences of atoms in A are treated differently.

In $Hz(A)$, each positive [negative] occurrence of an atom in the premise is linked with every positive [negative] occurrence in the conclusion.
We treat occurrences of an atom in A —with the same polarity—equally.

$Id(A)$, $Hz(A)$, $Mx(A)$

$Mx(p \vee (p \wedge \neg p))$



In $Id(A)$, each occurrence of an atom in the premise is linked with every its corresponding occurrence of in the conclusion.
Different occurrences of atoms in A are treated differently.


In $Hz(A)$, each positive [negative] occurrence of an atom in the premise is linked with every positive [negative] occurrence in the conclusion.
We treat occurrences of an atom in A —with the same polarity—equally.

In $Mx(A)$, each syntactically possible linking is included.
We treat all occurrences of an atom in A equally.


$H\mathbf{z}(A)$, $M\mathbf{x}(A)$, Caps and Cups

Note: $H\mathbf{z}(A)$ is $M\mathbf{x}(A)$ with the caps and cups removed.

$$H\mathbf{z}(p \vee (p \wedge \neg p))$$



$$M\mathbf{x}(p \vee (p \wedge \neg p))$$



Let's look at relations like isomorphism,
but which erase distinctions, up to Hx or Mx .

Hz-Matching

Let's say that A and B *Hz-MATCH*, when there are proofs $\pi : A \succ B$ and $\pi' : B \succ A$ where $\pi' \circ \pi = \text{Hz}(A)$ and $\pi \circ \pi' = \text{Hz}(B)$.

Hz-Matching

Let's say that A and B *Hz-MATCH*, when there are proofs $\pi : A \succ B$ and $\pi' : B \succ A$ where $\pi' \circ \pi = \text{Hz}(A)$ and $\pi \circ \pi' = \text{Hz}(B)$.

We write “ \approx_{Hz} ” for the *Hz*-matching relation, and we write “ $\pi, \pi' : A \approx_{\text{Hz}} B$ ” to say that $\pi : A \succ B$ and $\pi' : B \succ A$ define a *Hz*-match between A and B .

Mx-Matching

Let's say that A and B Mx -MATCH, when there are proofs $\pi : A \succ B$ and $\pi' : B \succ A$ where $\tau \circ \pi = Mx(A)$ and $\pi \circ \pi' = Mx(B)$.

Mx-Matching

Let's say that A and B Mx -MATCH, when there are proofs $\pi : A \succ B$ and $\pi' : B \succ A$ where $\tau \circ \pi = Mx(A)$ and $\pi \circ \pi' = Mx(B)$.

We write “ \approx_{Mx} ” for the Mx -matching relation, and we write “ $\pi, \pi' : A \approx_{Mx} B$ ” to say that $\pi : A \succ B$ and $\pi' : B \succ A$ define a Mx -match between A and B .

Isomorphism \subseteq Hz-Matching

If $\pi : A \succ B$ and $\pi^{-1} : B \succ A$, then
consider $\pi' = \text{Hz}(B) \circ \pi \circ \text{Hz}(A)$
and $\tau' = \text{Hz}(A) \circ \pi^{-1} \circ \text{Hz}(B)$.

These satisfy the Hz-matching criteria,
 $\tau' \circ \pi' = \text{Hz}(A)$ and $\pi' \circ \tau' = \text{Hz}(B)$.

Proof

$$\begin{aligned} Hz(A) &= Id(A) \circ Id(A) \circ Hz(A) \\ &\subseteq Hz(A) \circ Id(A) \circ Hz(A) \\ &= Hz(A) \circ (\pi^{-1} \circ \pi) \circ Hz(A) \\ &= Hz(A) \circ (\pi^{-1} \circ Id(B) \circ Id(B) \circ \pi) \circ Hz(A) \\ &\subseteq Hz(A) \circ (\pi^{-1} \circ Hz(B) \circ Hz(B) \circ \pi) \circ Hz(A) \\ &= (Hz(A) \circ \pi^{-1} \circ Hz(B)) \circ (Hz(B) \circ \pi \circ Hz(A)) \\ &= \tau' \circ \pi' \\ &\subseteq Hz(A) \end{aligned}$$

Proof

$$\begin{aligned} Hz(A) &= Id(A) \circ Id(A) \circ Hz(A) \\ &\subseteq Hz(A) \circ Id(A) \circ Hz(A) \\ &= Hz(A) \circ (\pi^{-1} \circ \pi) \circ Hz(A) \\ &= Hz(A) \circ (\pi^{-1} \circ Id(B) \circ Id(B) \circ \pi) \circ Hz(A) \\ &\subseteq Hz(A) \circ (\pi^{-1} \circ Hz(B) \circ Hz(B) \circ \pi) \circ Hz(A) \\ &= (Hz(A) \circ \pi^{-1} \circ Hz(B)) \circ (Hz(B) \circ \pi \circ Hz(A)) \\ &= \tau' \circ \pi' \\ &\subseteq Hz(A) \end{aligned}$$

...and similarly, $Hz(B) \subseteq \pi' \circ \tau' \subseteq Hz(B)$

$\text{Hz-Matching} \subseteq \text{Mx-Matching}$

If $\pi, \pi' : A \approx_{\text{Hz}} B$, then
consider $\tau = \text{Mx}(B) \circ \pi \circ \text{Mx}(A)$
and $\tau' = \text{Mx}(A) \circ \pi' \circ \text{Mx}(B)$.

These satisfy the Mx-matching criteria,
 $\tau' \circ \pi' = \text{Mx}(A)$ and $\pi' \circ \tau' = \text{Mx}(B)$.

Proof

$$\begin{aligned} Mx(A) &= Id(A) \circ Id(A) \circ Mx(A) \\ &\subseteq Mx(A) \circ Hz(A) \circ Mx(A) \\ &= Mx(A) \circ (\pi' \circ \pi) \circ Mx(A) \\ &= Mx(A) \circ (\pi' \circ Id(B) \circ Id(B) \circ \pi) \circ Mx(A) \\ &\subseteq Mx(A) \circ (\pi' \circ Mx(B) \circ Mx(B) \circ \pi) \circ Mx(A) \\ &= (Mx(A) \circ \pi' \circ Mx(B)) \circ (Mx(B) \circ \pi \circ Mx(A)) \\ &= \tau' \circ \tau \\ &\subseteq Mx(A) \end{aligned}$$

Proof

$$\begin{aligned} Mx(A) &= Id(A) \circ Id(A) \circ Mx(A) \\ &\subseteq Mx(A) \circ Hz(A) \circ Mx(A) \\ &= Mx(A) \circ (\pi' \circ \pi) \circ Mx(A) \\ &= Mx(A) \circ (\pi' \circ Id(B) \circ Id(B) \circ \pi) \circ Mx(A) \\ &\subseteq Mx(A) \circ (\pi' \circ Mx(B) \circ Mx(B) \circ \pi) \circ Mx(A) \\ &= (Mx(A) \circ \pi' \circ Mx(B)) \circ (Mx(B) \circ \pi \circ Mx(A)) \\ &= \tau' \circ \tau \\ &\subseteq Mx(A) \end{aligned}$$

...and similarly, $Mx(B) \subseteq \pi' \circ \tau' \subseteq Mx(B)$

If $A \approx_{Mx} B$ then there are proofs
 $\pi : A \multimap B$ and $\tau : B \multimap A$.

Matching Relations are Equivalence Relations

REFLEXIVE $Hx(A), Hx(A) : A \approx_{Hx} A.$

$Mx(A), Mx(A) : A \approx_{Mx} A.$

Matching Relations are Equivalence Relations

REFLEXIVE $H_z(A), H_z(A) : A \approx_{H_z} A.$

$M_x(A), M_x(A) : A \approx_{M_x} A.$

SYMMETRIC If $\pi, \pi' : A \approx_{H_z} B$, then $\pi', \pi : B \approx_{H_z} A.$

If $\pi, \pi' : A \approx_{M_x} B$, then $\pi', \pi : B \approx_{M_x} A.$

Matching Relations are Equivalence Relations

REFLEXIVE $H_z(A), H_z(A) : A \approx_{H_z} A.$

$M_x(A), M_x(A) : A \approx_{M_x} A.$

SYMMETRIC If $\pi, \pi' : A \approx_{H_z} B$, then $\pi', \pi : B \approx_{H_z} A.$


If $\pi, \pi' : A \approx_{M_x} B$, then $\pi', \pi : B \approx_{M_x} A.$

TRANSITIVE If $\pi, \pi' : A \approx_{H_z} B$ and $\tau, \tau' : B \approx_{H_z} C$, then
 $(\tau \circ \pi), (\pi' \circ \tau') : A \approx_{H_z} C.$

If $\pi, \pi' : A \approx_{M_x} B$ and $\tau, \tau' : B \approx_{M_x} C$, then
 $(\tau \circ \pi), (\pi' \circ \tau') : A \approx_{M_x} C.$


More Matchings

$$p \vee p \approx_{Hz} p \approx_{Hz} p \wedge p$$





More Matchings

$$p \vee p \approx_{Hz} p \approx_{Hz} p \wedge p$$





More Matchings

$$p \wedge (q \vee r) \approx_{\text{Hz}} (p \wedge q) \vee (p \wedge r)$$



More Matchings

$$p \wedge (q \vee r) \approx_{\text{Hz}} (p \wedge q) \vee (p \wedge r)$$



Mx -Matching \subset Logical Equivalence

FACT: If an atom p occurs positively in A but not in B , then A and B do not Mx -match.

FACT: If an atom p occurs positively in A but not in B , then A and B do not Mx -match.

PROOF: $Mx(A) : A \succ A$ contains the link from that occurrence of p in the premise A to its corresponding occurrence in the conclusion A .

FACT: If an atom p occurs positively in A but not in B , then A and B do not Mx -match.

PROOF: $Mx(A) : A \succ A$ contains the link from that occurrence of p in the premise A to its corresponding occurrence in the conclusion A .

No proof from A to B contains a link from that occurrence to anything in B (since there is no positive occurrence in B at all).

FACT: If an atom p occurs positively in A but not in B , then A and B do not Mx -match.

PROOF: $Mx(A) : A \succ A$ contains the link from that occurrence of p in the premise A to its corresponding occurrence in the conclusion A .

No proof from A to B contains a link from that occurrence to anything in B (since there is no positive occurrence in B at all).

So, in the composition proof from A to A , there is no link from the premise occurrence to the conclusion occurrence. No proof from A to B and back can recreate $Mx(A)$.

Mx -Matching \subset Logical Equivalence

FACT: If an atom p occurs positively [negatively] in A but not in B , then A and B do not Mx -match.

PROOF: $Mx(A) : A \succ A$ contains the link from [to] that occurrence of p in the premise A to [from] its corresponding occurrence in the conclusion A .

No proof from A to B contains a link from [to] that occurrence to anything in B (since there is no positive [negative] occurrence in B at all).

So, in the composition proof from A to A , there is no link from [to] the premise occurrence to [from] the conclusion occurrence. No proof from A to B and back can recreate $Mx(A)$.

Mx -Matching \subseteq Logical Equivalence: Examples

FACT: If an atom occurs positively
[negatively] in A but not in B
then A and B do not Mx -match.

Mx -Matching \subseteq Logical Equivalence: Examples

FACT: If an atom occurs positively
[negatively] in A but not in B
then A and B do not Mx -match.

COROLLARY: $p \not\approx_{Mx} p \wedge (q \vee \neg q)$.

Mx -Matching \subseteq Logical Equivalence: Examples


FACT: If an atom occurs positively
[negatively] in A but not in B
then A and B do not Mx -match.

COROLLARY: $p \not\approx_{Mx} p \wedge (q \vee \neg q)$.

$$p \wedge \neg p \not\approx_{Mx} q \wedge \neg q.$$



$H\mathbf{z}$ -matching $\subset M\mathbf{x}$ -matching

$$(p \wedge \neg p) \wedge (q \vee \neg q) \approx_{M\mathbf{x}} (p \vee \neg p) \wedge (q \wedge \neg q)$$



$H\mathbf{z}$ -matching $\subset M\mathbf{x}$ -matching

$$(p \wedge \neg p) \wedge (q \vee \neg q) \approx_{M\mathbf{x}} (p \vee \neg p) \wedge (q \wedge \neg q)$$



Isomorphism \subset Hz-Matching \subset Mx-Matching \subset Logical Equivalence

So what *are* the *matching* relations?

MATCHING & LOGICS OF ANALYTIC CONTAINMENT

Angell's Logic of Analytic Containment

$$\text{AC1 } A \leftrightarrow \neg\neg A$$

$$\text{AC2 } A \leftrightarrow (A \wedge A)$$

$$\text{AC3 } (A \wedge B) \leftrightarrow (B \wedge A)$$

$$\text{AC4 } A \wedge (B \wedge C) \leftrightarrow (A \wedge B) \wedge C$$

$$\text{AC5 } A \vee (B \wedge C) \leftrightarrow (A \vee B) \wedge (A \vee C)$$

$$\text{R1 } A \leftrightarrow B, C(A) \Rightarrow C(B)$$

Angell's Logic of Analytic Containment

$$\text{AC1 } A \leftrightarrow \neg\neg A$$

$$\text{AC2 } A \leftrightarrow (A \wedge A)$$

$$\text{AC3 } (A \wedge B) \leftrightarrow (B \wedge A)$$

$$\text{AC4 } A \wedge (B \wedge C) \leftrightarrow (A \wedge B) \wedge C$$

$$\text{AC5 } A \vee (B \wedge C) \leftrightarrow (A \vee B) \wedge (A \vee C)$$

$$\text{R1 } A \leftrightarrow B, C(A) \Rightarrow C(B)$$

Here, $A \vee B$ is shorthand for $\neg(\neg A \wedge \neg B)$.

Angell's Logic of Analytic Containment

$$\text{AC1 } A \leftrightarrow \neg\neg A$$

$$\text{AC2 } A \leftrightarrow (A \wedge A)$$

$$\text{AC3 } (A \wedge B) \leftrightarrow (B \wedge A)$$

$$\text{AC4 } A \wedge (B \wedge C) \leftrightarrow (A \wedge B) \wedge C$$

$$\text{AC5 } A \vee (B \wedge C) \leftrightarrow (A \vee B) \wedge (A \vee C)$$

$$\text{R1 } A \leftrightarrow B, C(A) \Rightarrow C(B)$$

Here, $A \vee B$ is shorthand for $\neg(\neg A \wedge \neg B)$.

You can define $A \rightarrow B$ as $A \leftrightarrow (A \wedge B)$.

Angell's Logic of Analytic Containment

$$\text{AC1 } A \leftrightarrow \neg\neg A$$

$$\text{AC2 } A \leftrightarrow (A \wedge A)$$

$$\text{AC3 } (A \wedge B) \leftrightarrow (B \wedge A)$$

$$\text{AC4 } A \wedge (B \wedge C) \leftrightarrow (A \wedge B) \wedge C$$

$$\text{AC5 } A \vee (B \wedge C) \leftrightarrow (A \vee B) \wedge (A \vee C)$$

$$\text{R1 } A \leftrightarrow B, C(A) \Rightarrow C(B)$$

Here, $A \vee B$ is shorthand for $\neg(\neg A \wedge \neg B)$.

You can define $A \rightarrow B$ as $A \leftrightarrow (A \wedge B)$.

Famously, $A \rightarrow (A \vee B)$ is not derivable in Angell's logic.
We cannot prove $A \leftrightarrow (A \wedge (A \vee B))$.

Extensions of Angell's Logic

- ▶ The first degree fragment of *Parry's Logic of Analytic Containment* is found by adding $(A \vee (B \wedge \neg B)) \rightarrow A$ to Angell's Logic.
 - ▶ Parry's logic still satisfies this relevance constraint: $A \rightarrow B$ is provable only when the atoms in B are present in A .
- ▶ *First Degree Entailment* (FDE) is found by adding $A \rightarrow (A \vee B)$ to Angell's Logic.
 - ▶ FDE allows for a disjunctive (or conjunctive) normal form. The only difference from classical logic is that $p \vee \neg p$, and $q \wedge \neg q$ are both non-trivial, and ineliminable.
 - ▶ A simple translation encodes FDE inside classical logic. Choose, for each atom p , a fresh atom p' , its *shadow*. For each FDE formula A , its translation is the formula A' found by replacing the negative occurrences of atoms p in A by their shadows. An argument is FDE valid iff its translation is classically valid.

$$Mx(A, B)$$

DEFINITION: $Mx(A, B)$ is the set of all possible linkings which could occur in any proof from A to B .

DEFINITION: $Mx(A, B)$ is the set of all possible linkings which could occur in any proof from A to B .

That is, it contains a link from
 {positive atoms in A , negative atoms in B }
to matching {positive atoms in B , negative atoms in A }.

$Mx(A, B)$

DEFINITION: $Mx(A, B)$ is the set of all possible linkings which could occur in any proof from A to B .

That is, it contains a link from
 {positive atoms in A , negative atoms in B }
to matching {positive atoms in B , negative atoms in A }.

FACT: $Mx(A, B)$ is a proof iff
there is some proof from A to B .

$Mx(A, B)$ examples

$Mx(p, q)$

p

q

$Mx(A, B)$ examples

$Mx(p, q)$

p

q

No links.

$Mx(A, B)$ examples

$Mx(p, q)$

p


q

No links.

$Mx(p \vee \neg p, p \wedge \neg q)$

$p \vee \neg p$

$p \wedge \neg q$



$Mx(A, B)$ examples


$Mx(p, q)$

p

q

No links.

$Mx(p \vee \neg p, p \wedge \neg q)$



Not a proof.

$Mx(A, B)$ and matching

LEMMA: If $A \approx_{Mx} B$, then $Mx(A, B)$ and $Mx(B, A)$ are proofs, and $Mx(A, B), Mx(B, A) : A \approx_{Mx} B$

$Mx(A, B)$ and matching

LEMMA: If $A \approx_{Mx} B$, then $Mx(A, B)$ and $Mx(B, A)$ are proofs, and $Mx(A, B), Mx(B, A) : A \approx_{Mx} B$

PROOF: If $\pi, \pi' : A \approx_{Mx} B$, then $\pi \subseteq Mx(A, B)$ and $\pi' \subseteq Mx(B, A)$, so $Mx(A, B)$ and $Mx(B, A)$ are both proofs.

$Mx(A, B)$ and matching

LEMMA: If $A \approx_{Mx} B$, then $Mx(A, B)$ and $Mx(B, A)$ are proofs, and $Mx(A, B), Mx(B, A) : A \approx_{Mx} B$

PROOF: If $\pi, \pi' : A \approx_{Mx} B$, then $\pi \subseteq Mx(A, B)$ and $\pi' \subseteq Mx(B, A)$, so $Mx(A, B)$ and $Mx(B, A)$ are both proofs.

Since $\pi' \circ \pi = Mx(A)$, we have
 $Mx(A) = \pi' \circ \pi \subseteq Mx(B, A) \circ Mx(A, B) \subseteq Mx(A)$,
and similarly, $Mx(B) = Mx(A, B) \circ Mx(B, A)$,
so $Mx(A, B), Mx(B, A) : A \approx_{Mx} B$.

FACT: If A is classically logically equivalent to B ,
and all atoms occurring positively [negatively] in A
also occur positively [negatively] in B ,
and *vice versa*, then A and B Mx -match
— and conversely.

Proof

If A is logically equivalent to B , then $Mx(A, B)$ and $Mx(B, A)$ are both proofs.

It suffices to show that $Mx(B, A) \circ Mx(A, B) = Mx(A)$ (and similarly for B). To show this, we need to show that each positive [negative] occurrence of an atom in A is linked to any positive [negative] occurrence of that atom in A by way of some link in $Mx(A, B)$ composed with a link in $Mx(B, A)$. But since that atom occurs positively [negatively] also in B at least once, the links to accomplish this occur in $Mx(A, B)$ and $Mx(B, A)$.

Conversely, if $A \approx_{Mx} B$, we have already seen that A and B must be equivalent, and no atom occurs positively [negatively] in A but not B .

This is *not* Equivalence in Parry's Logic

FACT: A is equivalent to B in Parry's logic of analytic containment iff A is classically equivalent to B and the atoms present in A are the atoms present in B .

This is *not* Equivalence in Parry's Logic

FACT: A is equivalent to B in Parry's logic of analytic containment iff A is classically equivalent to B and the atoms present in A are the atoms present in B.

$$(p \wedge \neg p) \wedge q \not\approx_{Mx} (p \wedge \neg p) \wedge \neg q$$

But this pair satisfies Parry's variable sharing criteria.

Open Question

Does the equivalence relation of Mx -matching
occur elsewhere in the literature?

$$Hz(A, B)$$

DEFINITION: $Hz(A, B)$ is the set of all possible linkings which could occur in any proof from A to B , excluding caps and cups.

$Hz(A, B)$

DEFINITION: $Hz(A, B)$ is the set of all possible linkings which could occur in any proof from A to B , excluding caps and cups.

That is, it contains a link from positive atoms in A to corresponding positive atoms in B and negative atoms in A to corresponding negative atoms in B .

$H_z(A, B)$ examples

$$H_z(p \wedge \neg p, q \vee \neg q)$$

$H_z(A, B)$ examples

$$p \wedge \neg p$$

$$H_z(p \wedge \neg p, q \vee \neg q)$$

$$q \vee \neg q$$

$H_z(A, B)$ examples

$$p \wedge \neg p$$

$$H_z(p \wedge \neg p, q \vee \neg q)$$

No links!

$$q \vee \neg q$$

$H_z(A, B)$ examples

$$p \wedge \neg p$$

$$H_z(p \wedge \neg p, q \vee \neg q)$$

No links!

$$q \vee \neg q$$

$$H_z(p \wedge \neg p, p \vee \neg p)$$

$H_z(A, B)$ examples

$$H_z(p \wedge \neg p, q \vee \neg q)$$

$$p \wedge \neg p$$

No links!

$$q \vee \neg q$$

$$H_z(p \wedge \neg p, p \vee \neg p)$$

$$\begin{array}{c} p \wedge \neg p \\ \downarrow \uparrow \\ p \vee \neg p \end{array}$$

$Hz(A, B)$ examples

$$Hz(p \wedge \neg p, q \vee \neg q)$$

$$p \wedge \neg p$$

No links!

$$q \vee \neg q$$

$$Hz(p \wedge \neg p, p \vee \neg p)$$

$$\begin{array}{ccc} p \wedge \neg p & & p \wedge \neg p \\ \downarrow & & \uparrow \\ p \vee \neg p & & p \vee \neg p \end{array}$$

A proof, but not
the maximal one.

FACT: $H_z(A, B)$ is a proof iff
the argument from A to B is FDE valid.

FACT: $H_z(A, B)$ is a proof iff
the argument from A to B is FDE valid.

- ▶ From FDE-validity to H_z -proof: straightforward induction on an FDE-axiomatisation.

FACT: $Hz(A, B)$ is a proof iff
the argument from A to B is FDE valid.

- ▶ From FDE-validity to Hz -proof: straightforward induction on an FDE-axiomatisation.
- ▶ From the Hz -proof $Hz(A, B)$ to FDE-validity. Notice that no negative occurrences of atoms in A or B are linked to any positive occurrences of atoms in A or B . So, there is another Hz -proof $Hz(A', B')$ for the FDE translations for A and B .

$Hz(A, B)$ and Hz -matching

LEMMA: If $A \approx_{Hz} B$, then $Hz(A, B)$ and $Hz(B, A)$ are proofs, and $Hz(A, B), Hz(B, A) : A \approx_{Hz} B$

$Hz(A, B)$ and Hz -matching

LEMMA: If $A \approx_{Hz} B$, then $Hz(A, B)$ and $Hz(B, A)$ are proofs, and $Hz(A, B), Hz(B, A) : A \approx_{Hz} B$

PROOF: If $\pi, \pi' : A \approx_{Hz} B$, then then since $\pi' \circ \pi = Hz(A)$ and $\pi \circ \pi' = Hz(B)$, π and π' are cap- and cup-free, so $\pi \subseteq Hz(A, B)$ and $\pi' \subseteq Hz(B, A)$, so $Hz(A, B)$ and $Hz(B, A)$ are both proofs.

$Hz(A, B)$ and Hz -matching

LEMMA: If $A \approx_{Hz} B$, then $Hz(A, B)$ and $Hz(B, A)$ are proofs, and $Hz(A, B), Hz(B, A) : A \approx_{Hz} B$

PROOF: If $\pi, \pi' : A \approx_{Hz} B$, then then since $\pi' \circ \pi = Hz(A)$ and $\pi \circ \pi' = Hz(B)$, π and π' are cap- and cup-free, so $\pi \subseteq Hz(A, B)$ and $\pi' \subseteq Hz(B, A)$, so $Hz(A, B)$ and $Hz(B, A)$ are both proofs.

Since $\pi' \circ \pi = Hz(A)$, we have
 $Mx(A) = \pi' \circ \pi \subseteq Hz(B, A) \circ Hz(A, B) \subseteq Hz(A)$,
and similarly, $Hz(B) = Hz(A, B) \circ Hz(B, A)$,
so $Hz(A, B), Hz(B, A) : A \approx_{Mx} B$.

FACT: If A is FDE-equivalent to A ,
and all atoms occurring positively [negatively] in A
also occur positively [negatively] in B ,
and *vice versa*, then A and B *Hz*-match
— and conversely.

Proof

If A is FDE-equivalent to B , then $\text{Hz}(A, B)$ and $\text{Hz}(B, A)$ are both proofs.

It suffices to show that $\text{Hz}(B, A) \circ \text{Hz}(A, B) = \text{Hz}(A)$ (and similarly for B). To show this, we need to show that each positive [negative] occurrence of an atom in A is linked to any positive [negative] occurrence of that atom in A by way of some link in $\text{Hz}(A, B)$ composed with a link in $\text{Hz}(B, A)$. But since that atom occurs positively [negatively] also in B at least once, the links to accomplish this occur in $\text{Hz}(A, B)$ and $\text{Hz}(B, A)$.

Conversely, if $A \approx_{\text{Hz}} B$, we have already seen that A and B must be FDE-equivalent, and no atom occurs positively [negatively] in A but not B .

Hz-matching \equiv Angellic Equivalence

FACT: (Fine, Ferguson) A is equivalent to B in Angell's logic of analytic containment iff A is FDE equivalent to B , and any atom occurs positively [negatively] in A iff it occurs positively [negatively] in B .

$Hz\text{-matching} \equiv \text{Angellic Equivalence}$

FACT: (Fine, Ferguson) A is equivalent to B in Angell's logic of analytic containment iff A is FDE equivalent to B , and any atom occurs positively [negatively] in A iff it occurs positively [negatively] in B .

So, $Hz\text{-matching}$ is equivalence
in Angell's Logic of Analytic Containment.

MATCHING AS ISOMORPHISM

$H_z(A)$ and $Mx(A)$ are Idempotents

- ▶ $H_z(A) \circ H_z(A) = H_z(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.

$Hx(A)$ and $Mx(A)$ are Idempotents

- ▶ $Hx(A) \circ Hx(A) = Hx(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.
- ▶ For any category \mathcal{C} , if i_A is an idempotent for each object A , we can form a new category \mathcal{C}_i with the same objects as \mathcal{C} , and with arrows $i_B \circ f \circ i_A : A \rightarrow B$.

$Hx(A)$ and $Mx(A)$ are Idempotents

- ▶ $Hx(A) \circ Hx(A) = Hx(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.
- ▶ For any category \mathcal{C} , if i_A is an idempotent for each object A , we can form a new category \mathcal{C}_i with the same objects as \mathcal{C} , and with arrows $i_B \circ f \circ i_A : A \rightarrow B$.
- ▶ In this new category, the idempotents i_A are the new identity arrows.

$Hz(A)$ and $Mx(A)$ are Idempotents

- ▶ $Hz(A) \circ Hz(A) = Hz(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.
- ▶ For any category \mathcal{C} , if i_A is an idempotent for each object A , we can form a new category \mathcal{C}_i with the same objects as \mathcal{C} , and with arrows $i_B \circ f \circ i_A : A \rightarrow B$.
- ▶ In this new category, the idempotents i_A are the new identity arrows.
- ▶ So, \mathcal{C}_{Hz} and \mathcal{C}_{Mx} are both categories — like \mathcal{C} , but less discriminating, with fewer arrows.


$Hz(A)$ and $Mx(A)$ are Idempotents

- ▶ $Hz(A) \circ Hz(A) = Hz(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.
- ▶ For any category \mathcal{C} , if i_A is an idempotent for each object A , we can form a new category \mathcal{C}_i with the same objects as \mathcal{C} , and with arrows $i_B \circ f \circ i_A : A \rightarrow B$.
- ▶ In this new category, the idempotents i_A are the new identity arrows.
- ▶ So, \mathfrak{C}_{Hz} and \mathfrak{C}_{Mx} are both categories — like \mathfrak{C} , but less discriminating, with fewer arrows.
- ▶ Hz -matching is *isomorphism* in \mathfrak{C}_{Hz} .

$Hx(A)$ and $Mx(A)$ are Idempotents

- ▶ $Hx(A) \circ Hx(A) = Hx(A)$, $Mx(A) \circ Mx(A) = Mx(A)$.
- ▶ For any category \mathcal{C} , if i_A is an idempotent for each object A , we can form a new category \mathcal{C}_i with the same objects as \mathcal{C} , and with arrows $i_B \circ f \circ i_A : A \rightarrow B$.
- ▶ In this new category, the idempotents i_A are the new identity arrows.
- ▶ So, \mathfrak{C}_{Hx} and \mathfrak{C}_{Mx} are both categories — like \mathfrak{C} , but less discriminating, with fewer arrows.
- ▶ Hx -matching is *isomorphism* in \mathfrak{C}_{Hx} .
- ▶ Mx -matching is *isomorphism* in \mathfrak{C}_{Mx} .

\mathfrak{C}_{Mx} and \mathfrak{C}_{Hz} are nontrivial, nonetheless



These are each different proofs in \mathfrak{C}_{Mx} and \mathfrak{C}_{Hz} .

\mathfrak{C}_{Mx} and \mathfrak{C}_{Hz} are nontrivial, nonetheless

$$\begin{array}{c} p \wedge q \\ \downarrow \\ p \vee q \end{array}$$

$$\begin{array}{c} p \wedge q \\ \downarrow \quad \downarrow \\ p \vee q \end{array}$$

$$\begin{array}{c} p \wedge q \\ \downarrow \\ p \vee q \end{array}$$

These are each different proofs in \mathfrak{C}_{Mx} and \mathfrak{C}_{Hz} .

To Conclude

- ▶ Proof theoretical resources *for classical logic* provide tools for fine-grained hyperintensional distinctions.

To Conclude

- ▶ Proof theoretical resources *for classical logic* provide tools for fine-grained hyperintensional distinctions.
- ▶ Extending these results to include the units \top and \perp are not difficult. (They were left out only to ease the presentation).

To Conclude

- ▶ Proof theoretical resources *for classical logic* provide tools for fine-grained hyperintensional distinctions.
- ▶ Extending these results to include the units \top and \perp are not difficult. (They were left out only to ease the presentation).
- ▶ Relate these results to *models* of logics of content.

To Conclude

- ▶ Proof theoretical resources *for classical logic* provide tools for fine-grained hyperintensional distinctions.
- ▶ Extending these results to include the units \top and \perp are not difficult. (They were left out only to ease the presentation).
- ▶ Relate these results to *models* of logics of content.
- ▶ Extend these results to first order logic, and beyond!

THANK YOU!

Thank you!

SLIDES: <http://consequently.org/presentation/>

FEEDBACK: @consequently on *Twitter*,
or *email* at `restall@unimelb.edu.au`