

# What Proofs are For

*Greg Restall*



THE UNIVERSITY OF  
MELBOURNE

MELBOURNE–GLASGOW WORKSHOP · JUNE 12, 2018

To present an account of the nature of proof,  
with the aim of explaining how proof could  
actually play the role in reasoning that it does,  
and answering some long-standing puzzles  
about the nature of proof.

Along the way, I'll explain how Kreisel's *squeezing argument* helps us understand the connection between an informal notion of validity and the notions formalised in our accounts of proofs and models, and the relationship between proof-theoretic and model-theoretic analyses of logical consequence.

Motivation

Background

What Proofs Are & What They Do

Counterexamples & Kreisel's Squeeze

Consequences for How Proofs Work

# MOTIVATION

## Example Proof

---

Every *drink* (in our fridge) is either a *beer* or a *lemonade*

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx)).$

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*



## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ .

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why?

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an  
arbitrary *drink*.

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an  
arbitrary *drink*. It's either a *beer* or a *lemonade*.

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink* is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we have the conclusion that some *drink* is a *lemonade*.

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an  
arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we  
have the conclusion that some *drink* is a *lemonade*. If we don't have  
that conclusion, then that arbitrary *drink* is a *beer*

## Example Proof


Every *drink* (in our fridge) is either a *beer* or a *lemonade*  
 $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink*  
is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an  
arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we  
have the conclusion that some *drink* is a *lemonade*. If we don't have  
that conclusion, then that arbitrary *drink* is a *beer*, and so, all the  
*drinks* are *beers*

## Example Proof

Every *drink* (in our fridge) is either a *beer* or a *lemonade*  $(\forall x)(Dx \supset (Bx \vee Lx))$ . So either every *drink* is a *beer* or some *drink* is a *lemonade*  $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$ . Why? Take an arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we have the conclusion that some *drink* is a *lemonade*. If we don't have that conclusion, then that arbitrary *drink* is a *beer*, and so, all the *drinks* are *beers*, and so, we also have our conclusion.



# Example Proof (the formal structure)



## Example Proof (the formal structure)

$$\begin{array}{c}
 \frac{Ba \succ Ba \quad La \succ La}{Ba \vee La \succ Ba, La} \vee L \\
 \frac{Da \succ Da \quad Ba \vee La \succ Ba, La}{Da \rightarrow (Ba \vee La), Da \succ Ba, La} \rightarrow L \\
 \frac{Da \succ Da \quad Da \rightarrow (Ba \vee La), Da \succ Ba, La}{Da \rightarrow (Ba \vee La), Da \succ Ba, Da \wedge La} \wedge R \\
 \frac{Da \rightarrow (Ba \vee La), Da \succ Ba, Da \wedge La}{Da \rightarrow (Ba \vee La) \succ Da \rightarrow Ba, Da \wedge La} \rightarrow R \\
 \frac{Da \rightarrow (Ba \vee La) \succ Da \rightarrow Ba, Da \wedge La}{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ Da \rightarrow Ba, Da \wedge La} \forall L \\
 \frac{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ Da \rightarrow Ba, Da \wedge La}{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ Da \rightarrow Ba, (\exists x)(Dx \wedge Lx)} \exists R \\
 \frac{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ Da \rightarrow Ba, (\exists x)(Dx \wedge Lx)}{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ (\forall x)(Dx \rightarrow Bx), (\exists x)(Dx \wedge Lx)} \forall R \\
 \frac{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ (\forall x)(Dx \rightarrow Bx), (\exists x)(Dx \wedge Lx)}{(\forall x)(Dx \rightarrow (Bx \vee Lx)) \succ (\forall x)(Dx \rightarrow Bx) \vee (\exists x)(Dx \wedge Lx)} \exists R
 \end{array}$$

# Our Focus

---

Our focus is on *categorical* proofs from premises to a conclusion.

# Our Focus

---

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic*

# Our Focus

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic* ( $\wedge, \vee, \neg, \rightarrow, \forall, \exists, =$ ).

# Our Focus

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic* ( $\wedge, \vee, \neg, \rightarrow, \forall, \exists, =$ ).

But what I say here can be extended to proof relying on other concepts.

## Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?

## Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?



## Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?
- ▶ What *grounds* the necessity in the connection between the premises and the conclusion?

## Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?
- ▶ What *grounds* the necessity in the connection between the premises and the conclusion?
- ▶ (Notice that these are important questions for proofs in first order predicate logic, as much as for proof more generally.)

# BACKGROUND

# *Assertions and Denials*

$[X : Y]$

... in a communicative practice

---

Assertions and denials are moves in a practice.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

I can ‘try on’ assertion or denial hypothetically.



Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

I can ‘try on’ assertion or denial hypothetically.

They are connected to other speech acts, too, like imperatives, interrogatives, recognitives, observatives, *etc.*

Assertions and denials take a *stand*  
(*pro* or *con*) on something.

DENIAL clashes with assertion.  
ASSERTION clashes with denial.

# Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.

## Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY:  $[A : A]$  is out of bounds.

## Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY:  $[A : A]$  is out of bounds.
- ▶ WEAKENING: If  $[X : Y]$  is out of bounds, then  $[X, A : Y]$  and  $[X : A, Y]$  are also out of bounds.

## Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY:  $[A : A]$  is out of bounds.
- ▶ WEAKENING: If  $[X : Y]$  is out of bounds, then  $[X, A : Y]$  and  $[X : A, Y]$  are also out of bounds.
- ▶ CUT: If  $[X, A : Y]$  and  $[X : A, Y]$  are out of bounds, then so is  $[X : Y]$ .

## Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY:  $[A : A]$  is out of bounds.
- ▶ WEAKENING: If  $[X : Y]$  is out of bounds, then  $[X, A : Y]$  and  $[X : A, Y]$  are also out of bounds.
- ▶ CUT: If  $[X, A : Y]$  and  $[X : A, Y]$  are out of bounds, then so is  $[X : Y]$ .
- ▶ A position that is OUT OF BOUNDS doesn't succeed in taking a stand.

## Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.



## Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

$x$  is a square  $=_{df}$   $x$  is a rectangle  $\wedge$   
all sides of  $x$  are equal in length.

## Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

$x$  is a square  $=_{df}$   $x$  is a rectangle  $\wedge$   
all sides of  $x$  are equal in length.

Concepts defined explicitly are *sharply delimited*  
(contingent on the definition).

## Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

$x$  is a square  $=_{df}$   $x$  is a rectangle  $\wedge$   
all sides of  $x$  are equal in length.

Concepts defined explicitly are *sharply delimited*  
(contingent on the definition).

Logical concepts are similarly sharply delimited,  
but they cannot all be given explicit definitions.

## Definition through a rule for use

$[X, A \wedge B : Y]$  is out of bounds

if and only if

$[X, A, B : Y]$  is out of bounds

## Definition through a rule for use

$[X, A \wedge B : Y]$  is out of bounds

if and only if

$[X, A, B : Y]$  is out of bounds

$$\frac{X, A, B \succ Y}{X, A \wedge B \succ Y} \wedge Df$$

## What about when to *deny* a conjunction?

$$\frac{\frac{X \succ A, Y \quad \frac{X \succ B, Y \quad \frac{\frac{A \wedge B \succ A \wedge B}{A, B \succ A \wedge B} \wedge Df}{X, A \succ A \wedge B, Y} \text{Cut}}{X \succ A \wedge B, Y} \text{Cut}}{X \succ A \wedge B, Y} \text{Cut}$$

## What about when to *deny* a conjunction?

$$\frac{\frac{X \succ A, Y \quad \frac{\frac{X \succ B, Y \quad \frac{\overline{A \wedge B \succ A \wedge B} Id}{A, B \succ A \wedge B} \wedge Df}{X, A \succ A \wedge B, Y} Cut}{X \succ A \wedge B, Y} Cut}{X \succ A \wedge B, Y} Cut$$

So, we have

$$\frac{X \succ A, Y \quad X \succ B, Y}{X \succ A \wedge B, Y} \wedge R$$

## Definitions for other logical concepts

$$\frac{X \succ A, Y}{X, \neg A \succ Y} \neg Df$$

$$\frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df$$

$$\frac{X \succ A, B, Y}{X \succ A \vee B, Y} \vee Df$$



## Definitions for other logical concepts

$$\frac{X \succ A, Y}{X, \neg A \succ Y} \neg Df$$

$$\frac{X, A \succ B, Y}{X \succ A \rightarrow B, Y} \rightarrow Df$$

$$\frac{X \succ A, B, Y}{X \succ A \vee B, Y} \vee Df$$

$$\frac{X \succ A|_n^x, Y}{X \succ (\forall x)A, Y} \forall Df$$

$$\frac{X, A|_n^x \succ Y}{X, (\exists x)A \succ Y} \forall Df$$

$$\frac{X, Fs \succ Ft, Y}{X \succ s = t, Y} =Df$$

(Where  $n$  and  $F$  are not present in  $X$  and  $Y$ .)

## Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)

## Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)

## Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)

## Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ▶ Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)

## Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ▶ Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)
- ▶ In Brandom's terms, they *make explicit* some of what was implicit in the practice of assertion and denial.

# WHAT PROOFS ARE & WHAT THEY DO

## A Tiny Proof

---

*If it's Thursday, I'm in Melbourne.*

*It's Thursday.*

*Therefore, I'm in Melbourne.*



## A Tiny Proof

*If it's Thursday, I'm in Melbourne.*

*It's Thursday.*

*Therefore, I'm in Melbourne.*

$$\frac{\frac{}{A \rightarrow B \succ A \rightarrow B} Id}{A \rightarrow B, A \succ B} \rightarrow Df$$

## A Tiny Proof

*If it's Thursday, I'm in Melbourne.*

*It's Thursday.*

*Therefore, I'm in Melbourne.*

$$\frac{\frac{}{A \rightarrow B \succ A \rightarrow B} Id}{A \rightarrow B, A \succ B} \rightarrow Df$$

*[It's Thursday  $\rightarrow$  I'm in Melbourne, It's Thursday : I'm in Melbourne]*

(This is out of bounds.)

## The Undeniable

Take a context in which I've asserted  
*it's Thursday  $\rightarrow$  I'm in Melbourne*  
and I've asserted *it's Thursday*,  
then *I'm in Melbourne* is undeniable.

## The Undeniable

Take a context in which I've asserted  
*it's Thursday  $\rightarrow$  I'm in Melbourne*  
and I've asserted *it's Thursday*,  
then *I'm in Melbourne* is undeniable.

Adding the *assertion* makes explicit  
what was *implicit* before that assertion.

## The Undeniable

Take a context in which I've asserted  
*it's Thursday  $\rightarrow$  I'm in Melbourne*  
and I've asserted *it's Thursday*,  
then *I'm in Melbourne* is undeniable.

Adding the *assertion* makes explicit  
what was *implicit* before that assertion.

The *stance* (*pro* or *con*)  
on *I'm in Melbourne* was already made.

**A *proof* for  $X \succ Y$  shows that the position  $[X : Y]$  is out of bounds, by way of the defining rules for the concepts involved in the proof.**

**A proof for  $X \succ Y$  shows that the position  $[X : Y]$  is out of bounds, by way of the defining rules for the concepts involved in the proof.**

In this sense, proofs are *analytic*.

They apply, given the definitions, independently of the positions taken by those giving the proof.

A proof of  $A, B \succ C, D$  can be seen  
as a *proof* of  $C$  from  $[A, B : D]$ ,



A proof of  $A, B \succ C, D$  can be seen  
as a *proof* of  $C$  from  $[A, B : D]$ ,  
and a *refutation* of  $A$  from  $[B : C, D]$ ,  
and *more*.

COUNTEREXAMPLES  
& KREISEL'S  
SQUEEZE

## Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \textit{Cut}$$

## Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \textit{Cut}$$

If  $X \succ Y$  is not derivable  
then one of  $X, A \succ Y$  and  $X \succ A \succ Y$   
is also not derivable.

## Enlarging Positions

$$\frac{X \succ A, Y \quad X, A \succ Y}{X \succ Y} \textit{Cut}$$

If  $X \succ Y$  is not derivable  
then one of  $X, A \succ Y$  and  $X \succ A \succ Y$   
is also not derivable.

If  $[X : Y]$  is available, then  
so is either  $[X, A : Y]$  or  $[X : A, Y]$

If  $[X : Y]$  is available,  
we can extend it into  
a *partition*  $[X' : Y']$   
of the entire language.

If  $[X : Y]$  is available,  
we can extend it into  
a *partition*  $[X' : Y']$   
of the entire language.

$U \succ V$  is not derivable  
for any finite  $U \subseteq X'$  and  $V \subseteq Y'$ .

## Adding Witnesses

If  $(\exists x)A$  is added on the left, we also  
add a *witness*  $A|_n^x$ , where  $n$  is fresh



## Adding Witnesses

If  $(\exists x)A$  is added on the left, we also add a *witness*  $A|_n^x$ , where  $n$  is fresh

$$\frac{X, A|_n^x, (\exists x)A \succ Y}{X, (\exists x)A \succ Y} \exists Df, W$$

## Adding Witnesses

If  $(\exists x)A$  is added on the left, we also add a *witness*  $A|_n^x$ , where  $n$  is fresh and similarly when  $(\forall x)A$  is added on the right.

$$\frac{X, A|_n^x, (\exists x)A \succ Y}{X, (\exists x)A \succ Y} \exists Df, W$$

$$\frac{X \succ (\forall x)A, A|_n^x, Y}{X \succ (\forall x)A, Y} \forall Df, W$$

# Witnessed Limit Positions give rise to Models

---

## Witnessed Limit Positions give rise to Models

$$A \in X' \text{ iff } \neg A \notin X' \text{ iff } \neg A \in Y',$$

$$A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$$

$$A \vee B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$$

$$A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$$

$$(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$$

$$(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$$

## Witnessed Limit Positions give rise to Models

$$A \in X' \text{ iff } \neg A \notin X' \text{ iff } \neg A \in Y',$$

$$A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$$

$$A \vee B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$$

$$A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$$

$$(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$$

$$(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$$

This is a *model*, where the *true formulas* are in  $X'$  and the *false formulas* are in  $Y'$ , and whose *domain* is the set of names.

## Witnessed Limit Positions give rise to Models

$$A \in X' \text{ iff } \neg A \notin X' \text{ iff } \neg A \in Y',$$

$$A \wedge B \in X' \text{ iff } A \in X' \text{ and } B \in X'.$$

$$A \vee B \in X' \text{ iff } A \in X' \text{ or } B \in X'.$$

$$A \rightarrow B \in X' \text{ iff } A \in Y' \text{ or } B \in X'.$$

$$(\forall x)A \in X' \text{ iff } A|_n^x \in X' \text{ for each name } n.$$

$$(\exists x)A \in X' \text{ iff } A|_n^x \in X' \text{ for some name } n.$$

This is a *model*, where the *true formulas* are in  $X'$  and the *false formulas* are in  $Y'$ , and whose *domain* is the set of names.

(Things are *little* more delicate when the language contains the identity predicate.)

$X \succ Y$  is derivable  
iff there is no *model*  
in which each member of  $X$  is true  
and each member of  $Y$  is false.

$X \succ Y$  is *informally* valid



$X \succ Y$  has a derivation



$X \succ Y$  is *informally* valid

$X \succ Y$  has a derivation



$X \succ Y$  is *informally* valid



$X \succ Y$  has no countermodel

$X \succ Y$  has a derivation



$X \succ Y$  is *informally* valid



$X \succ Y$  has no countermodel



$X \succ Y$  has a derivation.

## (1) From Derivability to Informal Validity

- ▶ To say that  $X \succ Y$  is *informally valid* means that is a clash involved in asserting each member of  $X$  and denying each member of  $Y$ .

## (1) From Derivability to Informal Validity

- ▶ To say that  $X \succ Y$  is *informally valid* means that is a clash involved in asserting each member of  $X$  and denying each member of  $Y$ .
- ▶ Axiomatic sequents ( $A \succ A$ ) are informally valid in this sense.

## (1) From Derivability to Informal Validity

- ▶ To say that  $X \succ Y$  is *informally valid* means that is a clash involved in asserting each member of  $X$  and denying each member of  $Y$ .
- ▶ Axiomatic sequents ( $A \succ A$ ) are informally valid in this sense.
- ▶ Structural rules preserve informal validity.

## (1) From Derivability to Informal Validity

- ▶ To say that  $X \succ Y$  is *informally valid* means that is a clash involved in asserting each member of  $X$  and denying each member of  $Y$ .
- ▶ Axiomatic sequents ( $A \succ A$ ) are informally valid in this sense.
- ▶ Structural rules preserve informal validity.
- ▶ Defining rules *define* the connectives/quantifiers.

## (2) From Informal Validity to Absence of Countermodel

---



## (2) From Countermodel to Informal *Invalidity*

---

## (2) From Countermodel to Informal *Invalidity*

- Refine our notion of informal validity: *Literals* ( $Fa$ ,  $Gbc$ , etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.

## (2) From Countermodel to Informal *Invalidity*

- ▶ Refine our notion of informal validity: *Literals* ( $Fa$ ,  $Gbc$ , etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ▶ Given a witnessed partition position  $[X : Y]$  (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in  $X$  and denying any in  $Y$ .

## (2) From Countermodel to Informal *Invalidity*

- ▶ Refine our notion of informal validity: *Literals* ( $Fa$ ,  $Gbc$ , etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ▶ Given a witnessed partition position  $[X : Y]$  (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in  $X$  and denying any in  $Y$ .
- ▶ So, there is no clash involved in asserting *any* formulas in  $X$  and denying any formulas in  $Y$ , by appeal to the defining rules.

## (2) From Countermodel to Informal *Invalidity*

- ▶ Refine our notion of informal validity: *Literals* ( $Fa$ ,  $Gbc$ , etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ▶ Given a witnessed partition position  $[X : Y]$  (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in  $X$  and denying any in  $Y$ .
- ▶ So, there is no clash involved in asserting *any* formulas in  $X$  and denying any formulas in  $Y$ , by appeal to the defining rules. (This is an induction on the depth of the structure of the formulas. The defining rules reduce clashes involving formulas into clashes involving subformulas.)

## (2) From Countermodel to Informal Invalidity

- ▶ Refine our notion of informal validity: *Literals* ( $Fa$ ,  $Gbc$ , etc.) are informally logically independent. We *ignore* logical connections between literals—we fix on informal validity *in virtue of first order logical form*.
- ▶ Given a witnessed partition position  $[X : Y]$  (i.e., given a model), there is no informal clash (in virtue of logical form) involved in asserting any of the literals in  $X$  and denying any in  $Y$ .
- ▶ So, there is no clash involved in asserting *any* formulas in  $X$  and denying any formulas in  $Y$ , by appeal to the defining rules. (This is an induction on the depth of the structure of the formulas. The defining rules reduce clashes involving formulas into clashes involving subformulas.)
- ▶ So, a countermodel for a sequent shows *how* there is no clash involved in asserting each member of  $X$  and denying each member of  $Y$ .

### (3) From Absence of Countermodel to Derivability

---

That's the *Completeness Theorem*.

## Kreisel's Squeeze

$X \succ Y$  has a derivation



$X \succ Y$  is *informally* valid



$X \succ Y$  has no countermodel



$X \succ Y$  has a derivation.



Informal validity (in virtue of first order logical form), for the language given by the defining rules, is *first order classical logic*, as given by the sequent calculus and Tarski's models.

# CONSEQUENCES FOR HOW PROOFS WORK

## Observation o: Proofs are *definitionally analytic*

Validity is grounded in the *rules*  
*defining* the concepts used in them.

## Observation 1: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.

## Observation 1: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.
- ▶ However, given plausible (minimal) assumptions concerning  $T$ , we can show that (*for example*) if  $A, B \succ C$  then  $T\langle A \rangle, T\langle B \rangle \succ T\langle C \rangle$ .

## Observation 1: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.
- ▶ However, given plausible (minimal) assumptions concerning  $T$ , we can show that (*for example*) if  $A, B \succ C$  then  $T\langle A \rangle, T\langle B \rangle \succ T\langle C \rangle$ .
- ▶ This *follows from* the concepts of consequence and truth.

## Observation 2: *Specification* outstrips *Recognition*

---

Our ability to *specify* concepts and consequence far outstrips our ability to *recognise* that consequence.

# Peano Arithmetic and Goldbach's Conjecture

## SUCCESSOR AXIOMS:

PA1:  $\forall x \forall y (x' = y' \rightarrow x = y)$ ;

PA2:  $\forall x (0 \neq x')$ .

## ADDITION AXIOMS:

PA3:  $\forall x (x + 0 = x)$ ;

PA4:  $\forall x (x + y' = (x + y)')$ .

## MULTIPLICATION AXIOMS:

PA5:  $\forall x (x \times 0 = 0)$ ;

PA6:  $\forall x \forall y (x \times y' = (x \times y) + x)$ .

## INDUCTION SCHEME:

PA7:  $(\phi(0) \wedge \forall x (\phi(x) \rightarrow \phi(x'))) \rightarrow \forall x \phi(x)$ .

## GOLDBACH'S CONJECTURE:

GC:  $\forall x \exists y \exists z (\text{Prime } y \wedge \text{Prime } z \wedge 0'' \times x = y + z)$



## Observation 2: *Specification* outstrips *Recognition*

Is  $[PA1, \dots, PA7 : GC]$  out of bounds?

## Observation 2: *Specification* outstrips *Recognition*

Is  $[PA1, \dots, PA7 : GC]$  out of bounds?

We have no idea.

## Observation 2: *Specification* outstrips *Recognition*

Is  $[PA1, \dots, PA7 : GC]$  out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

## Observation 2: *Specification* outstrips *Recognition*

Is  $[PA1, \dots, PA7 : GC]$  out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

Our concepts are rich and expressive.  
We can say things whose significance  
we continue to work out.

## Observation 2: *Specification* outstrips *Recognition*

Is  $[PA1, \dots, PA7 : GC]$  out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

Our concepts are rich and expressive.  
We can say things whose significance  
we continue to work out.

Verifying a putative proof is straightforward.  
Checking that something *has* a proof is not so easy.

## Are we logically omniscient?

Suppose that  $PA \succ GC$  is derivable  
(but we don't possess that proof)  
and that we *know*  $PA$ .

Do we know  $GC$ ?

In a weak sense of 'know', *yes*, we *do* know CC

- ▶ It's a logical consequence of what we know.

## In a weak sense of 'know', *yes*, we *do* know CC

- ▶ It's a logical consequence of what we know.
- ▶ It is implicitly present in what we already know.



## In a weak sense of 'know', *yes*, we *do* know GC

- ▶ It's a logical consequence of what we know.
- ▶ It is implicitly present in what we already know.
- ▶ There is no epistemic possibility (no circumstance consistent with our knowledge) that leaves GC out.

## In a weak sense of ‘know’, *yes*, we *do* know GC

- ▶ It's a logical consequence of what we know.
- ▶ It is implicitly present in what we already know.
- ▶ There is no epistemic possibility (no circumstance consistent with our knowledge) that leaves GC out.
- ▶ The means to come to know GC (the derivation) is “there” to be found.

In a not-so-weak sense, we don't know GC

---

- ▶ Do we *believe* GC?

## In a not-so-weak sense, we don't know GC

- ▶ Do we *believe* GC?
- ▶ If we do believe it, do we believe it *in the right way*?

## In a not-so-weak sense, we don't know GC

- ▶ Do we *believe* GC?
- ▶ If we do believe it, do we believe it *in the right way*?
- ▶ There “is” evidence for GC (its proof from PA, for example), but if that evidence plays no role in our belief...

## Observation 3: Proofs Can Transfer Warrant

- ▶ The account of consequence does not use the concept of *warrant*.

## Observation 3: Proofs Can Transfer Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (*for example*) if  $p$  is a proof for  $A, B \succ C$  then  $x : A, y : B \succ p(x, y) : C$ .

## Observation 3: Proofs Can Transfer Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (*for example*) if  $p$  is a proof for  $A, B \succ C$  then  $x : A, y : B \succ p(x, y) : C$ .
- ▶ Here,  $p$  transforms warrants for the premises into warrant for the conclusion.



## Observation 3: Proofs Can Transfer Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (for example) if  $p$  is a proof for  $A, B \succ C$  then  $x : A, y : B \succ p(x, y) : C$ .
- ▶ Here,  $p$  transforms warrants for the premises into warrant for the conclusion.
- ▶ This works only for *categorical, conclusive* warrants (*grounds*), not for *defeasible* warrants.

Consider the “Lottery Paradox.”

Consider the “Lottery Paradox.”

$$\begin{aligned} & [ (\exists x)(Tx \wedge Wx), \\ & (\forall x)(Tx \equiv (x = t_1 \vee x = t_2 \vee \dots \vee x = t_{1\,000\,000})) ) \\ & \quad : Wt_1, Wt_2, \dots, Wt_{1\,000\,000} ] \end{aligned}$$

Consider the “Lottery Paradox.”

$$\begin{aligned} [ & (\exists x)(Tx \wedge Wx), \\ & (\forall x)(Tx \equiv (x = t_1 \vee x = t_2 \vee \dots \vee x = t_{1\,000\,000})) \\ & \quad : Wt_1, Wt_2, \dots, Wt_{1\,000\,000} ] \end{aligned}$$

We have a very high degree of confidence in each part.

Each component is highly probable.

But the whole position is out of bounds.

## Observation 4: Achilles and the Tortoise

“Well, now, let’s take a little bit of the argument in that First Proposition—just *two* steps, and the conclusion drawn from them. Kindly enter them in your note-book. And in order to refer to them conveniently, let’s call them *A*, *B*, and *Z* :—

(*A*) Things that are equal to the same are equal to each other.

(*B*) The two sides of this Triangle are things that are equal to the same.

(*Z*) The two sides of this Triangle are equal to each other.

Readers of Euclid will grant, I suppose, that *Z* follows logically from *A* and *B*, so that any one who accepts *A* and *B* as true, *must* accept *Z* as true?”

“Undoubtedly! The youngest child in a High School—as soon as High Schools are invented, which will not be till some two thousand years later—will grant *that*.”

“And if some reader had *not* yet accepted *A* and *B* as true, he might still accept the *sequence* as a *valid* one, I suppose?”

## Observation 4: Achilles and the Tortoise

“No doubt such a reader might exist. He might say ‘I accept as true the Hypothetical Proposition that, *if*  $A$  and  $B$  be true,  $Z$  must be true; but, I *don't* accept  $A$  and  $B$  as true.’ Such a reader would do wisely in abandoning Euclid, and taking to football.”

“And might there not *also* be some reader who would say ‘I accept  $A$  and  $B$  as true, but I *don't* accept the Hypothetical’?”

“Certainly there might. *He*, also, had better take to football.”

“And *neither* of these readers,” the Tortoise continued, “is *as yet* under any logical necessity to accept  $Z$  as true?”

“Quite so,” Achilles assented.

“Well, now, I want you to consider *me* as a reader of the *second* kind, and to force me, logically, to accept  $Z$  as true.”

“A tortoise playing football would be—” Achilles was beginning

“—an anomaly, of course,” the Tortoise hastily interrupted. “Don’t wander from the point. Let’s have  $Z$  first, and football afterwards!”

“I’m to force you to accept  $Z$ , am I?” Achilles said musingly. “And your present position is that you accept  $A$  and  $B$ , but you *don't* accept the Hypothetical—”

“Let’s call it  $C$ ,” said the Tortoise.

“—but you *don't* accept

( $C$ ) If  $A$  and  $B$  are true,  $Z$  must be true.”

“That is my present position,” said the Tortoise.

“Then I must ask you to accept  $C$ .”

## Our Analysis

---

$$A, A \rightarrow Z \succ Z$$

## Our Analysis

$$A, A \rightarrow Z \succ Z$$

This *doesn't* mean when I accept  $A$   
and I accept  $A \rightarrow Z$ ,  
I ought to also accept  $Z$ .



## Our Analysis

$$A, A \rightarrow Z \succ Z$$

This *doesn't* mean when I accept  $A$   
and I accept  $A \rightarrow Z$ ,  
I ought to also accept  $Z$ .

However, if I assert  $A$  and  $A \rightarrow Z$  then  $Z$  is *undeniable*.

If I assert  $A$  and *if  $A$  then  $Z$*  and *deny  $Z$* ,  
then I am using ‘*if...then*’ in a way that  
deviates from the defining rule for  $\rightarrow$ ,  
or I am explicitly contradicting myself.

If I assert  $A$  and *if*  $A$  *then*  $Z$  and *deny*  $Z$ ,  
then I am using ‘*if...then*’ in a way that  
deviates from the defining rule for  $\rightarrow$ ,  
or I am explicitly contradicting myself.

$$\frac{A \rightarrow B \succ A \rightarrow B}{A \rightarrow B, A \succ B} \rightarrow Df$$

If I assert  $A$  and *if  $A$  then  $Z$*  and ask whether  $Z$  holds?

- ▶ We need to understand connections between defining rules and norms for questions and answers, as well as assertions and denials.

An account of the logical concepts given in terms of defining rules governing assertions and denials

An account of the logical concepts given in terms of defining rules governing assertions and denials helps explain how (*first order predicate logic*) proof works,

An account of the logical concepts given in terms of defining rules governing assertions and denials helps explain how (*first order predicate logic*) proof works, how possessing a proof can expand our knowledge, while proofs *make explicit* what is *implicit* in what we know.

# THANK YOU!

[https://consequently.org/presentation/2018/  
what-proofs-are-for-melbourne-glasgow/](https://consequently.org/presentation/2018/what-proofs-are-for-melbourne-glasgow/)

@consequently on Twitter