

Wavefunction

Probability of Finding the Particle between (a,b)

$$P_{a < x < b} = \int_a^b |\Psi(x, t)|^2 dx$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

21

Time-independent Shrodinger's Equation

Assume

$$-\frac{\hbar^2}{2m_0} \frac{d^2\Psi}{dx^2} + U(x)\Psi = i\hbar \frac{d\Psi}{dt}$$

$$\Psi(x, t) = \psi(x) e^{-iEt/\hbar}$$

$$-e^{-\frac{iEt}{\hbar}} \frac{\hbar^2}{2m_0} \frac{d^2\psi(x)}{dx^2} + e^{-\frac{iEt}{\hbar}} U(x)\psi(x) = i\hbar \frac{-iE}{\hbar} \psi(x) e^{-\frac{iEt}{\hbar}}$$

$$-\frac{\hbar^2}{2m_0} \frac{d^2\psi}{dx^2} + U(x)\psi = E\psi$$

$$\frac{d^2\psi}{dx^2} + \frac{2m_0}{\hbar^2} (E - U)\psi = 0$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

22

Time-independent Schrodinger's Equation

$$\frac{d^2\psi}{dx^2} + \frac{2m_0}{\hbar^2}(E - U)\psi = 0$$

If $E > U$, then

$$k \equiv \frac{\sqrt{2m_0[E - U]}}{\hbar} \quad \frac{d^2\psi}{dx^2} + k^2\psi = 0 \quad \psi(x) = A \sin(kx) + B \cos(kx) \\ \equiv A_+ e^{ikx} + A_- e^{-ikx}$$

If $U > E$, then

$$\alpha \equiv \frac{\sqrt{2m_0[U - E]}}{\hbar} \quad \frac{d^2\psi}{dx^2} - \alpha^2\psi = 0 \quad \psi(x) = D e^{-\alpha x} + E e^{+\alpha x}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

23

Basic Steps to Solve the Equation

$$-\frac{\hbar^2}{2m_0} \frac{d^2\psi}{dx^2} + U(x)\psi = E\psi$$

- Obtain $U(x)$ and the boundary conditions for a given problem.
- Solve the 2nd order equation – pretty basic
- Interpret $|\psi|^2 = \psi^* \psi$ as the probability of finding an electron at x
- Compute anything else you need, e.g.,

$$p = \int_0^\infty \Psi^* \left[\frac{\hbar}{i} \frac{d}{dx} \right] \Psi dx \quad E = \int_0^\infty \Psi^* \left[-\frac{\hbar^2}{2m_0} \frac{d^2}{dx^2} + U(x) \right] \Psi dx$$




University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

24

Solution of Schrodinger's Equation



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

25

Five Steps to Solve this Problem

- 1) $\frac{d^2\psi}{dx^2} + k^2\psi = 0$ \longrightarrow $2N$ unknowns for N regions
- 2) $\psi(x = -\infty) = 0$
 $\psi(x = +\infty) = 0$ \longrightarrow Reduces 2 unknowns
- 3) $\psi|_{x=x_B^-} = \psi|_{x=x_B^+}$
 $\frac{d\psi}{dx}|_{x=x_B^-} = \frac{d\psi}{dx}|_{x=x_B^+}$ \longrightarrow Set $2N-2$ equations for $2N-2$ unknowns (for continuous U)
- 4) Det (coefficient matrix)=0
And find E by graphical or numerical solution
- 5) $\int_{-\infty}^{\infty} |\psi(x, E)|^2 dx = 1$
for wave function




University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

26

Case 1: Bounded Levels in a Finite Well



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

27

Continuity of Wavefunction

$$3) \psi|_{x=x_B^-} = \psi|_{x=x_B^+}$$

$$\frac{d\psi}{dx}\bigg|_{x=x_B^-} = \frac{d\psi}{dx}\bigg|_{x=x_B^+}$$


$$C = B$$

$$\alpha C = -kA$$

$$A \sin(ka) + B \cos(ka) = D e^{-\alpha a}$$

$$kA \cos(ka) - kB \sin(ka) = -\alpha D e^{-\alpha a}$$

$$\psi = A \sin kx + B \cos kx$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

28

Continuity of Wavefunction

$$C = B$$

$$\alpha C = -kA$$

$$A \sin(ka) + B \cos(ka) = De^{-\alpha a}$$

$$kA \cos(ka) - kB \sin(ka) = -\alpha De^{-\alpha a}$$



$$\begin{pmatrix} 0 & 1 & -1 & 0 \\ k & 0 & \alpha & 0 \\ \sin(ka) & \cos(ka) & 0 & -e^{-\alpha a} \\ \cos(ka) - \sin(ka) & 0 & \alpha e^{-\alpha a} / k & 0 \end{pmatrix} \begin{bmatrix} A \\ B \\ C \\ D \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics


Bao Ve320 S12

29

Step 4: Bound-level in Finite Well

$$\det(\text{Matrix})=0$$

$$\tan(\alpha a \sqrt{\xi}) = \frac{2\sqrt{\xi(1-\xi)}}{2\xi-1} \quad \xi \equiv \frac{E}{U_0} \quad \alpha \equiv \sqrt{\frac{2mU_0}{\hbar^2}}$$



Only unknown is E

- (i) Use Matlab function
- (ii) Use graphical method



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics



Bao Ve320 S12

30

Graphical Method for Bound Levels

$$x^2 = x + 5$$

$$y_1 = x^2 \quad y_2 = x + 5$$




University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

31

Graphical Method for Bound Levels



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

32


Wave Function

Step 5:

$$\begin{pmatrix} 0 & 1 & -1 & 0 \\ k & 0 & \alpha & 0 \\ \sin(ka) & \cos(ka) & 0 & e^{-\alpha a} \\ \cos(ka) & -\sin(ka) & 0 & -\alpha D e^{-\alpha a} / k \end{pmatrix} \begin{pmatrix} A \\ B \\ C \\ D \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 & 0 \\ 0 & \alpha & 0 \\ \cos(ka) & 0 & e^{-\alpha a} \end{pmatrix} \begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} 0 \\ -kA \\ -A \sin(ka) \end{pmatrix}$$

$$\begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & \alpha & 0 \\ \cos(ka) & 0 & e^{-\alpha a} \end{pmatrix}^{-1} \begin{pmatrix} 0 \\ -kA \\ -A \sin(ka) \end{pmatrix}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

33

Calculation of Wave Function

$$\psi = A \sin kx + B \cos kx$$

$$\begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & \alpha & 0 \\ \cos(ka) & 0 & e^{-\alpha a} \end{pmatrix}^{-1} \begin{pmatrix} 0 \\ -kA \\ -A \sin(ka) \end{pmatrix}$$

$$\int_{-\infty}^{\infty} |\psi|^2 dx = 1 \Rightarrow \int_{-\infty}^0 C^2 e^{2\alpha x} dx + \int_0^a [A \sin(kx) + B \cos(kx)]^2 dx + \int_a^{\infty} D^2 e^{-2\alpha x} dx$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

34

Case 2: Infinite Quantum Well

$$\frac{d^2\psi}{dx^2} + k^2\psi = 0 \quad k \equiv \frac{\sqrt{2m_0[E-U]}}{\hbar}$$


1) Solutions: $\psi = A \sin kx + B \cos(kx)$

2) Boundary conditions

$$\psi(x=0) = 0 = A \sin k(0) + B \cos k(0)$$

$$\psi(x=a) = 0 = A \sin(ka) = A \sin(n\pi)$$

$$k_n = \frac{n\pi}{a} = \frac{\sqrt{2m_0 E_n}}{\hbar} \quad E_n = \frac{\hbar^2 n^2 \pi^2}{2m_0 a^2}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

35

Five Steps to Solve this Problem

1) $\frac{d^2\psi}{dx^2} + k^2\psi = 0 \longrightarrow$ 2N unknowns for N regions

2) $\psi(x = -\infty) = 0$
 $\psi(x = +\infty) = 0 \longrightarrow$ Reduces 2 unknowns

3) $\psi|_{x=x_B^-} = \psi|_{x=x_B^+}$
 $\frac{d\psi}{dx}|_{x=x_B^-} = \frac{d\psi}{dx}|_{x=x_B^+} \longrightarrow$ Set 2N-2 equations for 2N-2 unknowns (for continuous U)

4) Det (coefficient matrix)=0
And find E by graphical or numerical solution

5) $\int_{-\infty}^{\infty} |\psi(x, E)|^2 dx = 1$
for wave function



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics


Bao Ve320 S12

36

Case 3: $E \gg U$

$$\frac{d^2\psi}{dx^2} + k^2\psi = 0 \quad k \equiv \frac{\sqrt{2m_0[E-U]}}{\hbar}$$

1) Solution $\psi(x) = A \sin(kx) + B \cos(kx)$
 $\equiv A_+ e^{ikx} + A_- e^{-ikx}$



2) Boundary condition $\psi(x) = A_+ e^{ikx}$ positive going wave
 $= A_- e^{-ikx}$ negative going wave



University of Michigan – Shanghai Jiao Tong University Joint Institute
 Center of Optics and Optoelectronics

Bao Ve320 S12


37

Free Particle

$$\psi(x) = A \sin(kx) + B \cos(kx)$$

$$\equiv A_+ e^{ikx} + A_- e^{-ikx}$$

$\psi(x) = A_+ e^{ikx}$ positive going wave
 $= A_- e^{-ikx}$ negative going wave



Probability: $|\psi|^2 = \psi\psi^* = |A_+|^2 \text{ or } |A_-|^2$

Momentum: $p = \int_0^\infty \Psi^* \left[\frac{\hbar}{i} \frac{d}{dx} \right] \Psi dx = \hbar k \text{ or } -\hbar k$




University of Michigan – Shanghai Jiao Tong University Joint Institute
 Center of Optics and Optoelectronics

Bao Ve320 S12


38

Tunneling



Electron can stay in the potential barrier. Even if the electron energy is smaller than the potential barrier. It has a chance to go through the barrier!

This is different from classical particles!



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

39

Summary

- We have solved the Schrodinger's equation for finite well, infinite potential well, and free particle.

Next Week

- Solution of Schrodinger's equation in periodic potential
- E-k diagram, band structure
- Electrons and holes



University of Michigan – Shanghai Jiao Tong University Joint Institute
Center of Optics and Optoelectronics

Bao Ve320 S12

40