



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

## VE 320 – Summer 2012 Introduction to Semiconductor Device

Instructor: Professor Hua Bao

---

**NANO ENERGY LAB**

*Bao Ve320 S12*

1

### Previous Lecture

- In order to understand the transport property of semiconductor, we need to understand the chemical composition and atomic arrangements.
- Crystalline structure can be built by repeating basic building blocks... Bravais lattice, basis
- Diamond and zinc-blende structure
- To identify crystal planes... Miller Indices, vector indices




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

*Bao Ve320 S12*

2

## Getting back to Crystals




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

3

## Finally (almost) a Real Problem



But  $N$  atoms have  $4N$  unknowns constants to find...  
For large  $N$ , is there a better way?



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

4

## Five Steps to Solve this Problem

- 1)  $\frac{d^2\psi}{dx^2} + k^2\psi = 0$  —
- 2)  $\psi(x = -\infty) = 0$  —  
 $\psi(x = +\infty) = 0$
- 3)  $\psi|_{x=x_B^-} = \psi|_{x=x_B^+}$   
 $\frac{d\psi}{dx}|_{x=x_B^-} = \frac{d\psi}{dx}|_{x=x_B^+}$
- 4) Det (coefficient matrix)=0  
And find E by graphical or numerical solution

- N is very large for crystal, but changing steps 2 and 3 a little bit we can still solve the problem in a few minutes!



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics


Bao Ve320 S12

5

## Periodic U(x) and Bloch Theorem

$$|\psi(x)|^2 = |\psi(x+p)|^2 \Rightarrow \psi(x+p) = \psi(x) e^{ikp}$$

not our old (k)




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

6

## Phase Factor for N Cells




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

7

## Step 2: Periodic Boundary Condition



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

8

### Step 3: Boundary Conditions

$$\psi|_{x=0^-} = \psi|_{x=0^+}$$

$$\frac{d\psi}{dx}|_{x=0^-} = \frac{d\psi}{dx}|_{x=0^+}$$

$$B_a = B_b$$

$$\alpha A_a = \beta A_b$$

$$\alpha \equiv \sqrt{2mE/\hbar^2} \quad \beta \equiv i\sqrt{2m(U_0 - E)/\hbar^2}$$

$$\psi_a|_{x=a} = \psi_b|_{x=-b} e^{ikp}$$

$$\frac{d\psi_a}{dx}|_{x=a} = \frac{d\psi_b}{dx}|_{x=-b} e^{ikp}$$

$$A_a \sin \alpha a + B_a \cos \alpha a =$$

$$e^{ik(a+b)} [-A_b \sin \beta b + B_b \cos \beta b]$$

$$\alpha A_a \sin \alpha a - \alpha B_a \cos \alpha a =$$

$$e^{ik(a+b)} [\beta A_b \sin \beta b + \beta B_b \cos \beta b]$$



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

9

### Step 4: Det(matrix)=0 for Energy Levels

$$B_a = B_b$$

$$\alpha A_a = \beta A_b$$

$$A_a \sin \alpha a + B_a \cos \alpha a =$$

$$e^{ik(a+b)} [-A_b \sin \beta b + B_b \cos \beta b]$$

$$\alpha A_a \sin \alpha a - \alpha B_a \cos \alpha a =$$

$$e^{ik(a+b)} [\beta A_b \sin \beta b + \beta B_b \cos \beta b]$$



$$\begin{pmatrix} 0 & 1 & 0 & -1 \\ \alpha & 0 & \beta & 0 \\ * & * & & \\ * & & & \end{pmatrix} \begin{pmatrix} A_a \\ B_a \\ A_b \\ B_b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$



$$\frac{1-2\xi}{2\xi\sqrt{1-\xi}} \times \dots = \cos kp \quad \xi \equiv \frac{E}{U_0} \quad \alpha_0 \equiv \sqrt{\frac{2mU_0}{\hbar^2}}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics


Bao Ve320 S12

10

## Graphical Solution to Energy Levels

$$\frac{1-\xi}{2\xi\sqrt{1-\xi}} \times \cdots \cdots \cdots = \cos kp$$

$$k = \pm \frac{2\pi n}{Np} \quad n = -\frac{N}{2}, \dots, -1, 0, 1, \dots, \frac{N}{2}$$





University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

11

## Energy Band Diagram

$$\frac{1-\zeta}{2\xi\sqrt{1-\xi}} \times \cdots \cdots \cdots = \cos kp$$




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

12

## Brillouin Zone and Number of States



$$k = \pm \frac{2\pi n}{Np} \quad n = -\frac{N}{2}, \dots, -1, 0, 1, \dots, \frac{N}{2}$$

$$\frac{\text{States}}{\text{Band}} = \frac{k_{\max} - k_{\min}}{\Delta k} = \frac{2\pi/p}{2\pi/Np} = N$$

What is the physical meaning of the energy bands?




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics


Bao Ve320 S12

13

## Wave Packet and Group Velocity



$$\begin{aligned} \psi(x, t) &= Ae^{ikx - i\frac{E}{\hbar}t} + Ae^{i(k+\Delta k)x - i\frac{(E+\Delta E)}{\hbar}t} \\ &= Ae^{ikx - i\frac{E}{\hbar}t} \left[ 1 + e^{i(\Delta k)x - i\frac{\Delta E}{\hbar}t} \right] \end{aligned}$$




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

14

## Group Velocity for a Given Band

$$\begin{aligned}\psi(x,t) &= Ae^{ikx - i\frac{E}{\hbar}t} \left[ 1 + e^{i(\Delta k)x - i\left(\frac{\Delta E}{\hbar}\right)t} \right] \\ &= Ae^{ikx - i\frac{E}{\hbar}t} \left[ 1 + e^{i \times \text{const.}} \right]\end{aligned}$$


$$v = \frac{\Delta x}{\Delta t} = \frac{\Delta E}{\hbar \Delta k}$$

$$\therefore \left[ x \Delta k - t \frac{\Delta E}{\hbar} \right] = \text{constant.}$$

$$a = \frac{\Delta v}{\Delta t} = \frac{1}{\hbar} \frac{d}{dt} \left[ \frac{\Delta E}{\Delta k} \right] = \frac{1}{\hbar^2} \frac{d}{dk} \left[ \frac{\Delta E}{\Delta k} \right] \frac{d(\hbar k)}{dt} = \frac{F}{m^*}$$




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

15

## Effective Mass for a Given Band




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

16



## The Physics




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

17

## The Physics Cont'd




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

18

## Filled and Empty Bands



No electrons in the empty bands.

$$J_3 = -\frac{q}{L} \sum_{i(\text{filled})}^N v_i = 0$$

How about filled bands?

$$J_2 = -\frac{q}{L} \sum_{i(\text{filled})}^N v_i = -\frac{q}{L} \sum_0^{k_{\max}} v_i - \frac{q}{L} \sum_{-k_{\min}}^0 v_i = 0$$

Neither filled or empty bands can conduct electricity!




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

19

## Partially Filled Bands



$$J_3 = -\frac{q}{L} \sum_{i(\text{filled})} v_i \neq 0$$

$$\begin{aligned} J_2 &= -\frac{q}{L} \sum_{i(\text{filled})} v_i = -\frac{q}{L} \sum_{\text{all}} v_i + \frac{q}{L} \sum_{i(\text{empty})} |v_i| \\ &= \frac{q}{L} \sum_{i(\text{empty})} |v_i| \end{aligned}$$

-ve charge moving with -ve mass

+ve charge moving with +ve mass




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

20

## Metal, Semiconductor, Insulator



- **Metal:** has partially filled energy bands at zero temperature.
- **Semiconductor:** does not have partially filled bands at zero temperature, but thermal effect can excite electrons into conduction bands.
- **Insulator:** does not have partially filled bands at zero temperature, but band gap energy is too large and thermal effect cannot excite electrons on to conduction bands.



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

21

## Band Gap Energy

Si ~ 1.12 eV

Ge ~ 0.66 eV

GaAs ~ 1.42 eV

SiO<sub>2</sub> ~ 8 eV

Diamond ~ 5 eV





University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

22

## Carriers in the Semiconductor



In semiconductor, only the holes near the valence band maximum and the electrons near the conduction band minimum are responsible for conduction.






University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

23

## Simplified Planna View of Atoms




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

24

## Conduction Band?




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

25

## Conduction bands? Holes?




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

26

## Energy Bands for Real Materials




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

27

## Diamond and zinc-blende structure?




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics


Bao Ve320 S12

28


## Analog to E-k diagram



Density (x,y,z)  
4D information



Cut along  $(\theta_1, \phi_1) \dots$



A series of line-sections can  
Represent the 4D info in 2D plots





University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

29

## E-k along $\Gamma$ -X direction




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

30

## E-k along $\Gamma$ -L direction




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

31

## E-k Diagram for Ge, Si, GaAs



- 3 valence bands (light hole, heavy hole, split-off)  
valence bands near  $k=0$  is essentially  $E \sim k^2$
- Minima may not be at zone center
- (Ge: 8 L valleys, Si: 6 X valleys, and GaAs:  $\Gamma$  valleys)



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

32



## Effective Mass

$$a = \frac{\Delta v}{\Delta t} = \frac{1}{\hbar} \frac{d}{dt} \left[ \frac{\Delta E}{\Delta k} \right] = \frac{1}{\hbar^2} \frac{d}{dk} \left[ \frac{\Delta E}{\Delta k} \right] \frac{d(\hbar k)}{dt} = \frac{F}{m^*}$$

Do not understand?


Let's consider a simpler example...

**Free electrons in a vacuum respond to applied electric fields by the following**

$$\text{Force} = -q\mathcal{E} = m_0 \frac{dv}{dt} = \frac{dp}{dt}$$

$$E = \frac{1}{2}mv^2 = \frac{p^2}{2m}$$

**E-p relationship of electron.**




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

33

## Similarly...



The E-k relationship at the conduction band minimum or valance band maximum can be approximated by a parabolic function.

Wave-particle duality:

$$p = \hbar k, \quad E \approx \frac{p^2}{2m^*}$$

$$\frac{1}{m^*} = \frac{1}{\hbar^2} \frac{d^2 E}{dk^2}$$

Effective Mass is inversely proportional to the curvature of E-k diagram!

**Electrons moving in a solid:**

$$F = -q\mathcal{E} = m_n^* \frac{dv}{dt}$$

**Similar equation for holes:**

$$F = q\mathcal{E} = m_p^* \frac{dv}{dt}$$



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

34

## The importance of effective mass

$$F = -q\mathcal{E} = m_n^* \frac{dv}{dt}$$

$$F = q\mathcal{E} = m_p^* \frac{dv}{dt}$$

Quote from RFP:

*"It allows us to conceive of electrons and holes as quasi-classical particles and to employ classical particle relationships in most device analysis!"*

Now you can (mostly) forget about quantum mechanics....




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

35

## Heavy Holes and Light Holes



- There is typically degeneracy at the valence band maximum.
- There could be light holes and heavy holes.



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

36

## Effective Mass for Semiconductors

**Table 2.1** Density of States Effective Masses at 300 K.

Material	$m_n^*/m_0$	$m_p^*/m_0$
Si	1.18	0.81
Ge	0.55	0.36
GaAs	0.066	0.52

Effective mass is a material property.




University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

37

## Measurement of Effective Mass



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

Bao Ve320 S12

38

## Look back ...

- We know what the carriers in semiconductors are.  
(Electrons, holes)
- We know how they moves inside the semiconductor  
(effective mass)
- But we still do not know how many carriers are there.



University of Michigan – Shanghai Jiao Tong University Joint Institute  
Center of Optics and Optoelectronics

*Bao Ve320 S12*

39