

cover_letter_v6_ULTIMATE

2026-04-27

[Date: 2026-04-27]

Editor-in-Chief, *npj Precision Oncology*

Dear Editor,

We submit our manuscript “**An 8-gene RAI-responsiveness biomarker is the deployable molecular quantification of the WHO 2022 morphologic axis in papillary thyroid carcinoma**” for consideration as an Article. This v6 ULTIMATE version is the result of a comprehensive pre-submission self-audit and integration sprint that consolidates four lines of validation our v5 preprint did not include.

Why this manuscript

Decisions about radioiodine (RAI) therapy in papillary thyroid carcinoma rely on BRAF V600E mutation status and clinician judgement about differentiation, without a quantitative pre-treatment biomarker that outperforms the mutation alone. We provide an eight-gene panel of canonical thyroid-differentiation genes (NIS/SLC5A5, TPO, TG, TSHR, PAX8, NKX2-1, FOXE1, DIO1 — all measurable by NanoString or qPCR) that operationalises the WHO 2022 morphologic axis (cPTC + infiltrative FVPTC = BRAF-like, IEFVPTC = RAS-like) at the molecular level.

Pre-submission self-audit and corrections

We identified two issues in our v5 preprint and corrected them transparently in v6 ULTIMATE:

1. **Circular validation issue.** The original DM1/DM2 cluster was constructed using a 67-gene panel (TIERA67) that *included* the eight RAI-responsiveness genes; predicting that cluster from those same eight genes is autocorrelation. We re-derived the cluster four times with leak-free gene sets (zero overlap with the eight-gene panel): variant A (TIERA67 minus eight-gene, 47 genes, biology-preserving)

yields CV AUC **0.962** [0.940-0.979]; variant B (MAPK + immune + EMT + cell-cycle, 30 genes, mechanistically motivated zero-overlap) yields **0.925** [0.895-0.952]; variant D (BRS71 minus eight-gene, 26 genes, Chakravarty 2011 framework) yields **0.957** [0.936-0.976]; variant C (immune-only 5-feature) collapses to 0.664, demonstrating biology specificity. Δ AUC over BRAF V600E baseline is sustained at +0.111 to +0.130 across the three biology-preserving variants. The v5 figure of 0.954 is replaced with this honest range.

2. **Cluster orientation correction.** The leak-free re-derivation revealed that v5's labelling convention ("DM1 = aggressive, DM2 = differentiated") is reversed relative to the actual cluster identity. Four independent lines of evidence agree: (i) WHO 2022 mapping shows DM2 contains 85.6% BRAF-like classic PTC, while DM1 contains 77.5% RAS-like FVPTC; (ii) Pu 2021 dedifferentiation-signature direction concordance is 88.9% (8/9 marker genes); (iii) GSE97466 methylation cross-modality places 100% of aggressive histologies (ATC/PDTC/FTC/Hürthle/mFTC) in met_DM2; (iv) the largest local cohort GSE213647 (n=632) shows cPTC dm-score > PDTC > ATC, consistent with DM1 = differentiated. We adopt the corrected convention throughout v6 (DM1 = differentiated, DM2 = aggressive); the underlying biology and effect sizes are preserved — only the human-readable labels change orientation.

Four new validation layers added in v6 ULTIMATE

(1) **WHO 2022 morphologic alignment (n=476).** The DM1/DM2 axis aligns with the WHO 2022 morphologic axis with Cohen's $\kappa = 0.567$, sensitivity (BRAF-like $\rightarrow$ DM2) = 0.856, specificity (RAS-like $\rightarrow$ DM1) = 0.775, accuracy = 0.838, **odds ratio = 20.4 (Fisher exact p = 2.5×10^{-33})**. We position the eight-gene panel as the deployable molecular quantification of the morphologic axis.

(2) **Cross-modality methylation validation.** In an independent cohort (GSE97466, n=141, Illumina 450K), unsupervised K=2 clustering on the top-5,000 β -values produces two methylation clusters whose histology composition is striking: met_DM1 (n=41) = 100% PTC; met_DM2 (n=33) = 19 PTC + 4 FTC + 4 mFTC + 2 Hürthle + 1 PDTC + 3 ATC. Every aggressive histology lands in met_DM2 — independent confirmation that the differentiation axis is real and recovered at the methylation level.

(3) **Seven-cohort meta-analysis with true RAI clinical ground truth (★).** Across seven external cohorts (TCGA + GSE76039 + GSE27155 + GSE126698 + GSE213647 + GSE65144 + GSE60542 + GSE151179), the random-effects pooled AUC = **0.898** [0.835, 0.961], $I^2 = 75.8\%$. **The most clinically critical entry is**

GSE151179 (n=39 tumour specimens with directly-labelled RAI-refractory vs RAI-avid status; the open-access companion to the controlled-access Mu et al. JCEM 2024 cohort): the eight-gene panel achieves AUC = **0.671** [0.514, 0.946]. This is, to our knowledge, the first cross-validation of a deployable RAI panel against true clinical refractoriness labels rather than histology proxies.

(4) Literature integration with Korean parallel work. We position this work as the deployable minimal version of axes previously described by Han SC, Park YJ et al. (Endocrinol Metab 2023) — BL/RL-PTC transcriptomic phenotypes — and by Pu W et al. (Nat Commun 2021, BRAF-like-B subtype; Oncogene 2022, four-subtype framework). We extend these by (i) decision-curve clinical net benefit, (ii) cross-modality methylation, (iii) seven-cohort meta-analysis with RAI ground truth, and (iv) explicit WHO 2022 alignment. We additionally cite Yang H et al. (Endocrinol Metab 2022, n=2,092 Korean PTC) for population-level TERT promoter prevalence and present a 3-cohort prevalence range (Korean 2.8% PTC → TCGA 7.1% → MSK 63.4%) reflecting clinical-stage enrichment.

Limitations transparently reported

1. TCGA-THCA does not consistently distinguish encapsulated FVPTC (IEFVPTC) from infiltrative FVPTC; the FVPTC mixed category is treated as a RAS-like proxy at medium confidence — the main ceiling on the observed κ = 0.567.
2. GSE213647 (n=632) shows cross-platform attenuation (effect size 0.700 vs TCGA in-sample 0.972), reported transparently rather than excluded.
3. Mu et al. JCEM 2024 NGDC HRA004166 RAI-avidity dataset is controlled-access (DAC application required); GSE151179 was used as the best open alternative.
4. Korean prospective cohort validation is presently a stated next step; collaborative outreach to the Han / Park YJ group (SNU) and Bundang SNUH is in progress and will be reported in revision-round.
5. Drug-actionability findings (Fig. 7) remain hypothesis-generating (PRISM/CCLE n=5 vs 5).

Suggested reviewers

1. **Park YJ** (Seoul National University) — Han SC 2023 senior author, Korean parallel work; outreach in progress
2. **Pu W** (Sichuan University) — Nat Commun 2021 (scRNA, BRAF-like-B) + Oncogene 2022 (4-subtype)

3. **Landa I** (Memorial Sloan Kettering) — BRAF-like / RAS-like axis (J Clin Invest 2016)
4. **Fagin JA** (Memorial Sloan Kettering) — RAI refractoriness clinical mechanism
5. **Xing M** (Mayo Clinic) — TERT promoter prognosis framework

Reproducibility and data availability

All data are open-access: TCGA-THCA via GDC, six GEO cohorts, GSE97466 methylation, cBioPortal API for cross-cohort prevalence (six thyroid studies, n=2,194 samples). 32 executed Jupyter notebooks reproduce every figure end-to-end ([anonymous_code.zip](#) , ~2 MB; reproducibility verified via fresh re-runs). The interactive 8-gene RAI calculator ([rai_calculator.html](#)) and the unified interactive submission report ([interactive_report.html](#)) are included for reviewer convenience and remain client-side (no server roundtrip).

The authors declare no competing interests. This work has not been submitted elsewhere.

Sincerely,

Co-corresponding authors:

Seungho Cook Independent Researcher (with part-time PhD affiliation), Seoul, Republic of Korea kukshomr@gmail.com

[Yu Hyeong Won — full name + email TBD by user] [Affiliation TBD], Seoul, Republic of Korea

Internal note (NOT for editorial)

This v6 ULTIMATE cover letter incorporates the v5 STRENGTHEN content + v6 REAL FIX update + v6 ULTIMATE additions (WHO 2022 alignment + methylation cross-modality + 7-cohort meta with RAI clinical + literature integration + cluster orientation correction). All five additions are independently grounded in published-cohort data (no proprietary data, no controlled-access).

Submission scenario: **A · npj Precision Oncology**, 6/9 ULTIMATE criteria pass + 1 partial (RAI clinical CI brushes 0.5) + 1 limitation (GSE213647 attenuation) — all transparently reported.