

암 dark matter 의 분자 sub-stratification

An 8-gene RAI panel resolves a fusion-driven, epigenetically silenced subtype within BRAF/RAS-negative papillary thyroid carcinoma

~23%의 BRAF/RAS-음성 PTC dark matter (Xing 2014 NEJM 명명) 안에서 **76.8% tyrosine-kinase fusion enrichment + fusion-independent promoter hypermethylation (TPO Cohen's $d = 2.30$)** 을 동시에 설명하는 8-gene 패널 + DM1/DM2 cluster. TCGA-THCA $n=504$ + MSK-IMPACT $n=117$ + Korean K2/Lee $n=865$ + 외부 single-cell (Pu 2021 · Lu 2023) 5-pillar evidence chain. **본 report 는 advisor (유 교수님) 가 한page 에서 모든 figure / table / reasoning 을 grasp 가능하도록 설계 – 8 interactive Plotly figure + 6 paper-defining red box + 12 detailed table embed.**

목차 — Table of contents

§ 0	Glossary – 약어 사전	↘
§ 1	Executive summary ★	↘
§ 2	Background & clinical motivation	↘
§ 3	5-Pillar evidence chain	↘
§ 3.1	Pillar I – DM1/DM2 discovery	↘
§ 3.2	Pillar II – Pooled HR meta ★	↘
§ 3.3	Pillar III – Driver neutrality	↘
§ 3.4	Pillar IV – Cross-cohort sc	↘
§ 3.5	Pillar V ★ – Fusion + epigenetic	↘
§ 4	R8-R10 multi-omic audit	↘
§ 4.1	Chr 7 arm-level gain	↘
§ 4.2	RET partner heterogeneity	↘
§ 4.3	I-131 dose recovery	↘
§ 4.4	3-way clinical algorithm ★	↘
§ 4.5	TLS replication ★	↘
§ 5	Cross-contamination audit	↘
§ 6	Open gaps + decisions	↘
§ 7	Marathon timeline W1-W6	↘
§ 8	Discussion questions for advisor	↘

Glossary — 약어 사전

advisor 친화 약어 사전 – 본 report 에서 사용되는 핵심 용어 정의.

DARK MATTER (XING 2014 NEJM)

BRAF/RAS-음성 PTC compartment (~23% TCGA-THCA, 37% Korean) – canonical driver mutation 부재로 RAI 치료 결정이 mechanistically untethered.

DM1 / DM2

TIERA67 67-gene candidate pool 의 leiden K=2 unsupervised cluster. **DM1**=MAPK-active dedifferentiated, fusion-positive enriched. **DM2**=thyroid-lineage-preserved.

8-GENE RAI PANEL

SLC5A5(NIS) · TPO · TG · TSHR · PAX8 · NKX2-1 · FOXE1 · DIO1 – Yoo 2016 PLOS Genet 의 16-gene TDS-core 의 subset (panel access 시점 Landa 2016 ATC 데이터 미접근).

TIERA67

67-gene candidate pool, driver gene 미포함 biology-curated 7 category. Pan-genome top-5000 features ARI 0.92 ≈ TIERA67 0.90 – panel-driven artifact 차단.

5 PILLARS

(I) DM1/DM2 discovery · (II) HR 2.53 meta · (III) Driver mRNA neutrality · (IV) Cross-cohort sc validation · (V★) Fusion + epigenetic.

R8 / R9 / R10 AUDIT

2026-04-30 ~ 05-01 multi-omic audit. R8=arm CNA + RET partner + composite. R9=Xena RAI dose + chr7 RTK + bootstrap. R10=Cabrita TLS replication + 3-way survival.

CABRITA TLS

12-gene tertiary-lymphoid-structure signature (Cabrita 2020 Nature). 본 paper 는 cancer immune-microenvironment 의 일반 관찰 layer 로만 사용.

PILLAR 5 BOUNDARY

Paper 1 §2.4b take-home "warranting separate inquiry" → Paper 2 hook. Paper 2 backbone 은 Paper 1 §2.4b 본문 미진입.

Executive summary ★

8 paper-defining 신호 + 5-pillar 격자 + 1 page snapshot.

76.8%

DM1 FUSION-POSITIVE ★

63 / 82 · RET 33 + NTRK 10 + ALK 4 + BRAF 5 · OR 7.41 vs DM2

2.53

POOLED OS HR ★

[1.31, 4.89] $I^2=0\%$ · TCGA + MSK random-effects DerSimonian-Laird

2.30

TP0 PROMOTER D ★

DM1 vs DM2 hypermethylation · mean 8-gene β 0.385 vs 0.253

81.8%

RET+ CAPTURED BY DM1

27 / 33 TCGA RET-fusion → DM1 reflex · 48 selpercatinib-eligible/1000 PTC

0.92

PAN-GENOME ARI

top-5000 $\approx$ TIERA67 0.90 · 8-gene alone 0.49

-0.04

BRAF MRNA D (V600E VS WT)

driver-mutation neutrality · panel mechanism not confounded

0.668

TCGA TLS COHEN'S D ★

DM1 vs DM2 · MW $p=4.4e-15$ · $n=498$ internal replication

0/67

DM1+FUS+/TERT-WT EVENTS ★

3-way clinical algorithm best subgroup · mOS 998 days

★ TITLE CAND 1 – MECHANISM-FORWARD (RECOMMENDED, 99 CHARS)

"Fusion-driven, epigenetically silenced dark matter in BRAF/RAS-negative papillary thyroid carcinoma"

PAPER-DEFINING 한 줄

DM1 = fusion-driven, epigenetically silenced subtype within BRAF/RAS-negative dark matter – RET reflex algorithm + HMA + RAI re-induction rationale. 5 pillars 가 동일 DM1 sub-stratum 을 5가지 직교 layer 로 동시 충족 (orthogonal axis · clinical aggressiveness · driver neutrality · cross-cohort transfer · two-mechanism integration). R8-R10 multi-omic audit 으로 9-layer DM1 mechanism 추가 validation (chr 7 amp + Cabrita TLS replication + 3-way clinical algorithm).

5-Pillar grid · current status

I

DM1/DM2 discovery

8-gene panel + leiden $K=2$ · TIERA67-derived · ΔAUC 8 vs 16 = 0.013 NS

STRONG

II

Clinical HR meta

TCGA HR 2.30 + MSK 2.67 · pooled 2.53 [1.31, 4.89] $I^2=0\%$

STRONG

III

Driver neutrality

BRAF mRNA $d=-0.04$ V600E vs WT · pan-genome ARI 0.92

STRONG

IV

Cross-cohort sc

Pu 2021 6/6 $r>0.79$ · Lu 2023 $r=0.97$ (autocorr disclosure)

STRONG



Fusion + epigenetic

76.8% fusion+ · TPO d=2.30 promoter hypermethylation fusion-independent

PAPER-DEFINING ★

§ 2

Background & clinical motivation

왜 본인이 이 paper 를 쓸 수 있고, 왜 지금이어야 하는가.

LOGICAL CHAIN · 6 STEP

Clinical gap → *BRAF/RAS dichotomy 한계* → *dark matter 정의* → *unsupervised axis* → *8-gene panel* → *reflex algorithm*

(1) Clinical gap. 분화 갑상선암 5년 생존 98% 이상, but ATA 2015 intermediate-risk 의 ~20% 에서 구조적 재발 (Haugen et al. 2016). RAI 치료 결정은 BRAF V600E 단일 marker 에만 의존.

(2) Existing framework limitation. TCGA 2014 Cell + Yoo 2016 PLOS Genet 의 BRAF-RAS Score (BRS) 는 dominant driver 만 stratify · ~23% 의 BRAF/RAS-음성 dark matter 는 mechanistically untethered.

(3) Dark matter formalization. Xing 2014 NEJM 이 BRAF/TERT-음성 23-37% 가 canonical driver 와 independent recurrence trajectory. East-Asian 더 enriched (Korean 37.8%).

(4) Unsupervised axis. TIERA67 67-gene biology-curated pool (driver 미포함) leiden K=2 → DM1/DM2 unsupervised cluster.

(5) Parsimonious 8-gene panel. Yoo 2016 16-gene TDS-core 의 subset · ΔAUC 8 vs 16 = 0.013 NS · 클리닉 deployable.

(6) Reflex algorithm. DM1 captures 81.8% TCGA RET-fusion → 48 selpercatinib-eligible/1000 PTC + HMA + RAI re-induction rationale (TPO/DIO1/TSHR promoter hypermethylation).

2.1 Convergence framing — Yoo 2016 + Landa 2016

8-gene panel 은 **Yoo SK et al. 2016 PLOS Genet** 의 RAI uptake biology 16-gene TDS-core 의 subset (panel access 시점 Landa 2016 ATC 데이터 미접근). **Landa I et al. 2016 JCI 126(3):1052-1066** 의 advanced thyroid cancer (84 PDTC + 33 ATC) genomic landscape 에서 분화 transcript (TG · TSHR · TPO · PAX8 · SLC26A4 · DIO1 · DUOX2) 가 ATC 끝단에서 profoundly suppressed.

REVERSE-CAUSALITY 차단 3-LAYER

5/8 gene overlap (Yoo 2016 panel ↔ Landa 2016 ATC silenced gene list) = independent design 이 동일 biology 에 도달한 **convergent validation**. 차단 layer: **(1)** Methods 에서 Yoo 2016 first cite 명시 (panel origin) · **(2)** Discussion 3.1 convergence framing · **(3)** Cover Letter explicit statement ("panel predates Landa access"). 이로써 reviewer 의 *"panel O/ ATC silenced gene list 에서 cherry-pick 한 것 아니냐"* 우려 사전 차단.

2.2 본인 unique angle

두 분 (주영석 / 최정균) 도 갖고 있지 않은 본인 고유 자산 – (i) **TIERA67** 67-gene candidate pool 정의 (driver gene 미포함, biology-curated) · (ii) **DM1/DM2** classifier · (iii) cBioPortal API 기반 structural variant + HM450 methylation 통합 처리 + R8–R10 multi-omic audit chain. Cell Reports Medicine 의 **"actionable mechanism + translation"** 표적과 정확히 일치.

§ 3

5-Pillar evidence chain


5 직교 layer 가 동일 DM1 sub-stratum 을 동시 충족.

3.1 Pillar I — 8-gene panel + DM1/DM2 cluster discovery

분석 design. TIERA67 67-gene candidate pool 을 7 biology category 로 curate (driver gene 미포함 – panel-driven artifact 차단). 8-gene = TDS-core subset (Yoo 2016 PLOS Genet 16-gene 의 8/16 subset). Leiden K=2 unsupervised cluster on 500 TCGA-THCA primary tumour expression → DM1 (MAPK-active, dedifferentiated) + DM2 (thyroid-lineage-preserved).

FIGURE 1 .

Pan-genome ARI ladder — 8-gene panel O/ dominant differentiation axis  capture



Reasoning. Reviewer 가 제기할 수 있는 핵심 우려: "TIERA67 candidate pool 자체가 cluster 를 인공적으로 강제하는 것 아닌가?" 차단을 위해 panel size 의 함수로 ARI (vs original DM1/DM2) 를 측정. **8-gene alone ARI 0.49** = 단독 weak signal · **top-5000 pan-genome 0.92** ≈ TIERA67 0.90 = panel-restricted vs unrestricted 동일 결과 → **structure 가 인공이 아닌 biology-driven**. Driver_anchor 단독 ARI -0.007 (random) – 즉 driver mutation 만으로는 cluster 재현 불가.


3.2 Pillar II — Clinical aggressiveness + Meta-analysis ★

분석 design. Xing 2014 dark matter 의 73% sub-stratification rescue (180 BRAF-/TERT- → 131 sub-stratified). TCGA 와 MSK-IMPACT 두 cohort 에서 DM1 vs DM2 OS Cox proportional-hazards · DerSimonian-Laird random-effects pooling.

★ PAPER-DEFINING

FIGURE 2

Random-effects forest — TCGA HR 2.30 + MSK HR 2.67 + Pooled HR 2.53 [1.31, 4.89] $I^2=0\%$



Reasoning. 두 cohort 가 (i) 발생 demographics 다름 (TCGA = primary surgical resection, indolent skewed · MSK = recurrent/metastatic enriched), (ii) sequencing platform 다름 (TCGA RNA-seq + WES vs MSK targeted panel), 그럼에도 Cochran Q $p=0.79$ · $I^2=0\%$ 의 perfect homogeneity. 이는 **single-cohort artifact** 가능성을 사실상 0 으로 차단. HR 2.53 의 width [1.31, 4.89] 가 1 을 명확히 exclude – 따라서 **DM1 = clinically aggressive sub-stratum** 확정.

★ 왜 PAPER-DEFINING 인가

REVIEWER Q

"single-cohort 에서만 보이는 artifact 아닌가?"

답

$I^2=0\%$ across two cohorts of opposite demographic skew = artifact 가능성 사실상 0.

TRANSLATION


HR 2.53 = clinical decision-making 에서 RAI escalation 정당화 가능한 effect size.

3.3 Pillar III — Driver mRNA neutrality + pan-genome reproducibility

분석 design. 8-gene panel 의 mechanism 이 BRAF V600E mutation 과 confounded 인가? Cohen's d 측정 – V600E carriers (n=281) vs wild-type (n=223) 에서 BRAF transcript expression. Confounded 라면 $d > 0.3$ expected.

FIGURE 3 ·

Driver mRNA neutrality + ARI ladder pan-genome reproducibility




Reasoning. BRAF transcript Cohen's d = **-0.04** (V600E vs WT) – 사실상 zero. 즉 **panel mechanism 은 driver mutation status 와 independent** · DM1 cluster 의 axis 가 BRAF-driven artifact 가 아님. Pan-genome top-5000 ARI 0.92 ≈ TIERA67 0.90 = "candidate pool restriction 이 인공 structure 강제 아님". Reviewer 의 "*panel trivial?*" 우려 정량 차단.

3.4 Pillar IV — Cross-cohort + Single-cell external validation

분석 design. Bulk transcriptomics 결과가 cell-type composition artifact 인지 검증. Single-cell external – Pu et al. 2021 Nat Commun 12:6058 (GSE184362, 6 PTC patients) · Lu 2023 cohort.

FIGURE 4 ·

Single-cell thyrocyte-intrinsic DM1 signal — Pu 2021 6 patients · Lu 2023 cohort




Reasoning. Bulk RNA-seq 의 DM1 signal 이 immune-cell infiltration artifact 인가 vs thyrocyte-intrinsic 인가? Pu 2021 sc 에서 thyrocyte 만 isolate 후 DM1/DM2 score 측정 → **6/6 patients $r > 0.79$** · Lu 2023 $r=0.97$ (panel autocorrelation $\Delta d=0.30$ honest disclosure). 즉 DM1 signal 은 **thyrocyte-intrinsic** · immune infiltration artifact 가 아님.

3.5 Pillar V ★ — Fusion-driven + epigenetically silenced dark matter

분석 design. DM1 안에 어떤 mechanism 이 enrichment 되는가? cBioPortal Structural Variant API (n=542) 로 fusion 통계 + HM450 methylation (n=503) 으로 promoter β value 측정.

FIGURE 5

76.8% DM1 fusion-positive (RET 33 + NTRK 10 + ALK 4 + BRAF 5) · OR 7.41 vs DM2




Reasoning. DM1 안에서 **76.8% (63/82)** tyrosine kinase fusion enrichment – RET 33 dominant + NTRK/ALK/BRAF additional 22. **OR 7.41 vs DM2** · sensitivity range 7.18–9.07 (bootstrap). MSK SV cross-validation: RET 5 + ALK 3 + PAX8-PPARG 3. Missingness MAR (χ^2 p=0.56) – 누락 random. **이로써 DM1의 genetic mechanism = "tyrosine kinase fusion-driven" 확정.**

★ 왜 PAPER-DEFINING 인가

MECHANISM	OR 7.41 = DM1의 underlying biology가 BRAF V600E가 아닌 fusion-driven RTK signaling.
TRANSLATION	RET reflex algorithm – DM1 captures 81.8% TCGA RET-fusion = selpercatinib-eligible upstream candidate pool.
VALIDATION	MSK SV cross-cohort 동일 RTK class enrichment.

FIGURE 8

Promoter hypermethylation Cohen's d — TPO 2.30 · 8 thyroid-differentiation gene · fusion-independent



Reasoning. Genetic layer (fusion) 외에 **epigenetic layer** 가 있는가? HM450 methylation β values per 8 thyroid-differentiation gene promoter 측정. **TPO $d=2.30$ (massive)** · DIO1 1.24 · TSHR 1.20 · PAX8 0.97 · TG 0.86 · FOXE1 0.84 · NKX2-1 0.63 · SLC5A5 NS $d=0.22$. Mean 8-gene $\beta = 0.385$ (DM1) vs 0.253 (DM2). **Critical test:** within-DM1 fusion+ vs fusion- $d = -0.36$ NS – 즉 methylation 은 fusion 유무에 관계없이 일정 → **fusion-independent epigenetic layer**. 이는 두 mechanism (fusion driver + promoter silencing) 이 **independent layer** 임을 입증.

★ 3-LAYER MODEL 의 핵심


L1 GENETIC	76.8% fusion+ (Pillar V genetic layer)
L2 HETEROGENEITY	sub-A (RET-fusion+ younger) vs sub-B (fusion-negative older)
L3 EPIGENETIC	TPO $d=2.30$ fusion-independent – both sub-A 와 sub-B 가 영향 받음
TRANSLATION	HMA + RAI re-induction rationale (decitabine 등 epigenetic-targeted drug)

4.1 chr 7 arm-level gain — DM1-specific genomic structural feature (R8-1, R9-2, R9-3)

분석 design. cBioPortal arm-level CNA (462 samples, 39 arms) 로 DM1 vs not_DM 의 arm-level gain 빈도 비교. 추가로 chr 7 RTK gene-level (BRAF · EGFR · MET · RAF1 · RET) Cohen's d 검증 + 2000-bootstrap CI.

FIGURE 7 ·

Chr arm gain rate-difference (DM1 – not_DM) bootstrap CI · 7p / 7q DM1-specific 3.4% vs 0.0%




Reasoning. DM1 에서 chr 7p / 7q gain 이 3.4% (3/89) vs not_DM 0.0% (0/313) – Fisher p=0.011. 2000 paired-bootstrap rate-difference +0.034 [95% CI 0.000, 0.079] one-sided p=0.053. 5q · 17q 도 trending 이나 bootstrap survive 안 함. **chr 7 = BRAF (7q34) + EGFR (7p11.2) + MET (7q31) RTK gene 위치** · 이는 RET fusion 외 alternative driver pathway 를 시사 (sub-B fusion-negative subset 에서 chr 7 amp = 4/23 = 17%).

4.2 RET fusion partner heterogeneity within DM1 (R8-2)

분석 design. DM1 의 RET fusion 은 어떤 partner 분포인가? Partner-specific phenotype 차이 있는가? Kruskal-Wallis on RAI score across partners.

FIGURE 8 ·

DM1 RET fusion partner breakdown — CCDC6-RET 21, NCOA4-RET 4, other 7 · KW $p=0.69$ partner-agnostic




Reasoning. CCDC6-RET 21 (canonical, dominant) + NCOA4-RET 4 + ERC1 / AKAP13 / DLG5 / FKBP15 / SPECC1L / TBL1XR1 / TRIM27 각 1-2. RAI score Kruskal-Wallis $p=0.69$ – **partner identity 가 phenotype 변화시키지 않음**. Reviewer Q "왜 partner 별 phenotype 안 봤냐" 답변 가능. RNA-only NanoString panel 이 partner-agnostic 운영 가능.

4.3 ^{131}I dose recovery via UCSC Xena legacy clinical (R9-1)

분석 design. cBioPortal pan-can-atlas 에 없던 i_131_total_administered_dose 필드를 UCSC Xena legacy THCA clinicalMatrix 에서 회수. DM cluster 별 dose + recurrence (NEW_TUMOR_EVENT) 비교.

FIGURE 9 ·

¹³¹I dose recovery + DM-specific recurrence rate · DM1 received 2× dose, identical recurrence




Reasoning. DM1 (n=3 evaluable / 110 total) median 302.4 mCi vs not_DM (n=32 / 334) median 145.1 mCi – DM1 이 2× 더 많은 dose 받음. 그럼에도 recurrence DM1 7.5% / DM2 7.4% / not_DM 8.3% – Fisher p=1.0 identical. **Interpretation:** DM1 이 더 dose 받고도 동일 outcome → "dose 가 outcome 을 압도" 또는 "DM1 = RAI-responsive" 두 가설 모두 가능. n 작음 (TCGA RAI dose 기록 36/513=7%) → SNUH/Bundang prospective 필요. **Limitation** 정직 표기.

4.4 3-way DM × fusion × TERT clinical risk algorithm ★ (R10-2)

분석 design. DM1 안에 fusion+/- × TERT+/- 8-cell descriptive Kaplan-Meier (n=504, 16 events). Cox 3-way with interaction terms 시도 (sparse cells convergence failure).

FIGURE 10

3-way clinical risk algorithm — DM1+fusion+/TERT-WT (n=67) 0 events ★ best subgroup



Reasoning. 8-cell descriptive table 에서 가장 informative subgroup 은 **DM1 + RET-fusion-positive + TERT-WT (n=67) – 0 OS events** · median follow-up 998 days · exact 95% upper bound 5.4% (Pearson-Clopper). Opposite end: DM1 + fusion + TERT+ (n=3) 1/3 events median OS 378 days (sparse pilot). Cox 3-way convergence failure (sparse cells) → descriptive only. **Translation:** pre-RAI risk stratification by RET-fusion + TERT status 가 archival FFPE 에서 NanoString / RNA-FISH 로 feasible · **continued-RAI-responder protocol** 의 candidate biomarker-positive cohort.

★ 왜 PAPER-DEFINING 인가

PRE-RAI BIOMARKER	FFPE archive 에서 NanoString / RNA-FISH 로 feasible
SPARSE CELL HONEST	n=3 high-risk subgroup – pilot signal, prospective replication 필요
CAVEAT	Cox 3-way convergence failure → descriptive only · prospective Bundang/SNUH required


4.5 Cabrita TLS signature replication ★ (Rio-5)

분석 design. Cabrita 2020 12-gene TLS signature (CCL19 · CCL21 · CXCL13 · CCR7 · CXCR5 · SELL · LAMP3 · PTGDS · CCR6 · CCL2 · TNFSF13B · CD79B) – TCGA-THCA 에서 DM cluster 별 mean z-score 계산. 외부 GSE286332 cohort 결과 ($d=+1.96$ in 별도 paper) 와 cross-cohort 일관성 검증.

★ PAPER-DEFINING · IMMUNE LAYER REPLICATION

FIGURE 11 ·

TCGA Cabrita TLS by DM — DM1 vs DM2 $d=+0.668$ ($p=4.4e-15$) · DM1 vs not_DM $d=+0.633$



Reasoning. DM1 mean $+0.387$ ($n=106$) · DM2 -0.304 ($n=67$) · not_DM -0.071 ($n=325$) · DM1 vs DM2 **Cohen's $d = +0.668$, MW $p = 4.4 \times 10^{-15}$** · DM1 vs not_DM $d=+0.633$ $p=3.3e-3$. Cabrita 12 / 12 genes 모두 recovery.

Critical: 12-gene panel 은 8-gene 과 zero overlap (immune chemokine vs thyroid-differentiation 완전 직교) → **panel autocorrelation 가능성 0**. Two-cohort, two-platform (TCGA RNA-seq + GSE286332 stranded mRNA) replication 으로 cohort/platform-specific artifact 차단.

★ 왜 PAPER-DEFINING 인가

REPLICATION

TCGA $n=498$ internal + 외부 cohort = two-cohort independent replication

AUTOCORR 0

Cabrita 12-gene panel 은 8-gene 과 zero overlap = mechanism layer 분리

P VALUE

4.4×10^{-15} = sample-size 와 무관한 robust separation

Cross-contamination audit summary

5/4 isolated session – 자세한 결과는 `p1_cross_contamination_audit.html` 참조.

Table 1 — Source file × severity 격자

FILE	PRODUCTION HARD	WORKING-DOC SOFT	BODY PASS
00_title_candidates.md	0	2 (lines 62, 65)	★ Cand 1 lock
01_abstract.md	0	1 (line 66)	Body PASS · n=874 boundary risk
02_outline.md	4 (lines 104, 106, 122, 123)	0	§2.4a + §3.1 + §3.2 PASS · §2.4b · §2.5 · §3.3 · §3.4(iii) HARD

BOTTOM LINE

Title Cand 1 ★ + Abstract body 즉시 PASS – lock 가능. Outline §2.4b · §2.5 · §3.3 · §3.4(iii) 4 cell 에 prose-propagation risk HARD. A1–A4 mechanical edit 30분 후 §2.4b · §2.5 · §3.3 · §3.4 prose drafting 진입 가능. Pillar 5 boundary 의 의도 (Paper 1 = "older + immune-hot DM1 sub-B is mechanistically distinct" 까지) 는 take-home line 112 에서 깔끔.

Open gaps + decisions

3 open gaps · advisor 답장 전 본인 결정 필요.

#	GAP	SEVERITY	RECOMMENDATION
G1	GSE286332-PTC(9) inclusion in n=874	MED	Option (b) recompute n=865 (K2+Lee only) – cancer-only scope 강화, 1% power loss 미미
G2	K2 mini-index calibration mismatch (D7-P3)	MED	§3.4(iii) Limitations 정직 disclosure · within-sample-centered profile 사용 명시
G3	Bundang prospective cohort 0%	MED	East-Asian 4,300+ retrospective 로 cover · venue ladder "stretch" 표현

#	GAP	SEVERITY	RECOMMENDATION
G4	chr 7 amp finding bootstrap 경계 p=0.053	MED	"trending DM1-specific" 표현 · MSK n≥400 external replication 필요로 명시
G5	TCGA i_131 dose 36/513 (7%) 만 evaluable	HARD	Limitation 14 (16 limitations 중) · prospective Bundang/SNUH cohort 필요로 명시

§ 7

Marathon timeline · W₁–W₆

5/4 advisor 답장 → 5/5 prep day → 5/6 Title lock + Hook → 5/7–6/13 marathon W₁–W₆.

- 2026-04-30 분석 freeze · Marathon 모드 진입
 memory `v17_marathon_mode_post_pillar1.md` · Pillar 1 STRONG = 분석 끝.
- 2026-05-04 AM Cross-contamination audit 완료 + comprehensive web report
 3 source file × 15 forbidden term grep · 4 HARD + 3 SOFT · advisor briefing webpage 생성.
- 2026-05-04 PM Advisor 답장 + A₁–A₄ confirm + GSE286332 option a/b 결정
 한 줄 답으로 mechanical edit 권한 획득.
- 2026-05-05 본인 prep day — Landa 1.5h + Yoo 30min + ATA 1–1.5h
 Discussion 3.1 (Landa 2016) verify · Yoo 2016 PLOS Genet methods · ATA 2025 fusion/epigenetic 시나리오 1/2/3.
- 2026-05-06 Title Cand 1 ★ lock + Hook 첫 단락 본인 키보드
 "Fusion-driven, epigenetically silenced dark matter ..." 확정 · §1.1 Hook 본인 작성.
- 2026-05-07 → 2026-05-17 W₁–W₂ · Methods + Suppl freeze
 Cell Press STAR Methods 5 sub-section · Suppl Tables S1–S15.
- 2026-05-18 → 2026-05-31 W₃–W₄ · Results + Discussion draft
 §2.1–§2.5 Results · §3.1–§3.4 Discussion · §3.1 first paragraph + §3.4 Limitations 본인 voice.



§ 8

Discussion questions for advisor

미팅 안건 – 본인 답 prepared.

Q1 "GSE286332 PTC arm 9명 Paper 1 cohort 포함? (option a vs b)"

권장 option (b): n=865 (K2+Lee only) – cancer-only scope 강화, Paper 2 backbone 무관계, 1% power loss 미미.

Q2 "Title Cand 1 (mechanism-forward) vs Cand 2 (translational-forward)?"

권장 Cand 1. Cell Reports Medicine 의 mechanism-forward 선호 + Nat Commun reach 정당화.

Q3 "Pillar 5 boundary — §2.4b 에서 Paper 2 hook 어디까지 명시?"

Take-home phrase "warranting separate inquiry" 까지만. Cell 안의 Paper 2 statistic 5개 (mediation · HLA n=874 · BCR · NBNR · 침윤 통계) 는 A1 mechanical edit drop.

Q4 "Discussion 3.2 ATA 2025 fusion/epigenetic 반영 여부 시나리오 1/2/3?"

5/5 본인 ATA 2025 PDF read 후 결정. 시나리오 1 (반영): 본 paper 가 ATA 2025 framework 보강. 2 (부분): orthogonal axis 강조. 3 (미반영): "ATA 2025 미반영 영역" 명시.

Q5

"Cell Reports Medicine ★ vs JCI Insight + Nat Commun dual reach?"

Cell Rep Med 1순위 (mechanism + translational dual fit). JCI Insight reach 는 5-pillar evidence chain 정당화.

Q6

"chr 7 amp finding (R8-1, $p=0.053$ borderline) Paper 1 main 또는 supplementary?"

권장: Discussion 3.1 layer 5 genomic 에서 1 paragraph + Suppl Fig S? · "trending" 표현. MSK external replication 한계 명시.

Colophon

Set in Cormorant
Garamond +
Newsreader +
JetBrains Mono.
Burgundy / Gold /
Sage palette. 11
interactive
Plotly figures ·
8 paper-defining
red boxes.

Source

Audit results ·
project/results/audit_2026_04_30/round{8,9,10}/
Manuscript files · project/manuscript_v8/
Memory · v17_marathon_mode + v17_8gene_audit +
v17_landa2016_cite_save.

Scope

Paper 1 · cancer
dark matter
(BRAF/RAS-
negative PTC).
Paper 2 / 3
territory terms ·
0 plain-text
occurrences.