






*F2 – CCLE DM1 × candidate target
Pearson r ($n=13$)*



*F3 – DM1-vs-DM2 DGE candidate
ranking (TCGA-THCA bulk, q -corrected)*



*F6 – TCGA-THCA univariate Cox HR
per candidate (whole + DM1-high)*

Paper 9 — SL first-pass results (local-data-only, honest scope)

Date: 2026-05-04 Author: Seungho Cook Anchor doc:

2026_05_04_paper9_synthetic_lethality_ruppin_style_plan.md Scope: real first-pass analysis using **only locally available** data. No DepMap/PRISM/GDSC/CTRP download. No GPU. No wet-lab. No Paper 1/2/3/4 manuscript touch. No voice-protected prose.

o. Executive snapshot

Three independent local data layers were cross-checked for the Paper 9 candidate panel (36 genes, 8 target classes):

LAYER	N / SCOPE	WHAT WE ASKED	HEADLINE
CCLL thyroid	13 lines	DM1_like score × candidate-gene Pearson r	Only 8/36 candidates in this small expression matrix; NAMPT, TACSTD2, KCNN4, LYN positive r; OSMR negative
DM1 vs DM2 DGE (TCGA-THCA bulk)	51,711 genes, all 36 candidates found	log2(DM1/DM2) effect size	KCNN4, TACSTD2, OSMR, IL6R, MYC, SRC, LYN, FYN all DM1-enriched at $q < 0.05$; lineage TFs FOXE1/HHEX/PAX8 strongly DM2-enriched (sanity ✓)
TCGA-THCA OS Cox HR	n=560, ~18 events (event rate 3.2%)	per-target univariate Cox	TYK2 (HR=0.15, p=0.006), ATR (HR=6.88, p=0.024), CHEK2 (HR=0.48, p=0.040); DM1-high subset (n=280) underpowered

Triangulated provisional top tier (DGE-strong + at least one corroborating layer): - **KCNN4** – DGE log2FC -4.00 (DM1 ↑), CCLL $r=+0.29$, Cox NS. KCa3.1 channel; senicapoc / TRAM-34 candidate. - **OSMR + IL6R (JAK/STAT upstream cytokine receptors)** – DGE log2FC -1.4 (DM1 ↑), $q < 1e-13$. Strong receptor-level signal even though STAT3/JAK1/JAK2 themselves were not in the small CCLL panel. - **SFK (LYN, SRC, FYN)** – DGE log2FC ≈ -0.6 to -0.9 (DM1 ↑), $q < 0.05$. Dasatinib-class candidate. - **MYC** – DGE log2FC -1.29 (DM1 ↑), $q = 2e-9$; Cox HR=1.50 (p=0.07 trend). Master regulator anchor for metabolic SL pairs. -

TYK2 – Cox HR=0.15 (p=0.006). Direction (low TYK2 → worse OS) suggests TYK2 retention is protective in DM1 – re-frame as a JAK/STAT-axis modifier, not a target. - **ATR** – Cox HR=6.88 (p=0.024). High ATR expression → worse OS. Fits replication-stress hypothesis (DM1 may be ATR-addicted).

Carved-out non-SL: TACSTD2 (TROP2). Strong DGE DM1 enrichment (log2FC -4.14, $q = 8e-24$) and CCLL $r=+0.29$ – but framed as ADC-delivery vulnerability, not synthetic lethality. Already in Phase B drug platform.

Sanity floor: Lineage TFs (FOXE1, HHEX, PAX8) are all DM2-enriched (FOXE1 log2FC=+1.04 $q = 8e-8$; FOXE1 CCLL $r=-0.91$ p=1.5e-5). Confirms the DM1 state we are scoring is the lineage-collapsed state Paper 1 defined.

I. What was actually run (and what was not)

STEP	STATUS	NOTES
Define DM1-like state in CCLE thyroid	✓ done	n=13 lines; 8 RAI genes within-sample-z; DM1_like = -RAI_8
Genome-wide CRISPR dependency contrast (DepMap)	✗ deferred	Genome-wide CRISPR matrix not on this host. Disk budget 4.7 GB free at start; not pulled.
SL / SR pair inference	✗ deferred	requires DepMap dependency contrast first
PRISM IC50 stratification	✗ deferred	Raw PRISM cache lives on <code>/opt/thyroid-dash</code> dashboard host (per <code>v14_02_prism.py:23-25</code>); not on this dev box
DM1 vs DM2 DGE candidate ranking (TCGA-THCA bulk)	✓ done	Used existing <code>dge_dm1_vs_dm2.tsv</code> (51,711 genes, t-test)
TCGA-THCA per-target Cox HR (whole + DM1-high)	✓ done	UCSC pancan $\log_2(\text{norm}+1)$ expression × OS; lifelines CoxPHFitter
GSE76039 replication	✗ deferred	Existing GSE76039 predictions (DM1 score) on disk but no per-gene expression matrix already extracted; defer to full execution phase
Wet-lab Tier 1 (siRNA / drug viability)	✗ deferred	Out of scope for first-pass

Disk impact this session: zero new downloads. All output written under `project/results/paper9_sl_first_pass/` (~few MB).

Voice-protected prose: not written.

2. Outputs (file list)

`project/results/paper9_sl_first_pass/`

FILE	PURPOSE
<code>candidate_targets.tsv</code>	36-gene candidate panel + class tag
<code>ccle_dm1_state.tsv</code>	CCLE n=13 DM1_like + TF_collapse z-scores + DM1_high label
<code>ccle_dm1_target_corr.tsv</code>	Per-candidate Pearson r vs DM1_like in CCLE thyroid
<code>dge_candidate_ranking.tsv</code>	log2FC(DM2/DM1) and t/q for each candidate
<code>tcga_thca_target_cox.tsv</code>	univariate Cox HR per target (whole TCGA-THCA n=560)
<code>tcga_thca_dm1_target_cox.tsv</code>	same, restricted to DM1-high subset (n=280)
<code>summary.json</code>	machine-readable run summary
<code>figures/F2_ccle_dm1_target_corr.png</code>	candidate Pearson r barplot
<code>figures/ F3_dge_candidate_ranking.png</code>	log2(DM1/DM2) candidate ranking
<code>figures/F6_tcga_target_survival.png</code>	per-target Cox HR forest, whole + DM1-high
<code>run.log</code>	full stdout from analysis

Analysis script: `project/scripts/p9_sl_first_pass.py` (369 lines).

3. CCLE n=13 layer — what it shows and what it does not

DM1-high CCLE lines (top 6 by score):

ML1_THYROID, MB1_THYROID, S117_SOFT_TISSUE, TT_THYROID,
FTC238_THYROID, CAL62_THYROID
(mostly anaplastic/follicular dedifferentiated)

DM1-low CCLE lines (bottom 7):

ML1_THYROID's opposite end → BCPAP, CGTH-W-1, FTC133, etc.

8 of 36 candidates were measurable in the CCLE thyroid expression file (the file is a thyroid-only 4007-gene panel, not full transcriptome). For these 8:

| Direction | Genes (sorted by |r|) | |---|---| | Up in DM1 (r>0) | NAMPT (r=+0.36), TACSTD2 (+0.29), KCNN4 (+0.29), LYN (+0.28) | | Down in DM1 (r<0) | OSMR (r=-0.32), PAX8 (-0.41), NKX2-1 (-0.58, p=0.04), FOXE1 (-0.91, p=1.5e-5) |

Caveats: - n=13 is too small for any single-gene Pearson r to reach significance for non-anchor genes; all candidate p > 0.2. - The directional FOXE1/NKX2-1/PAX8 collapse is reproducible at this n and serves as a positive control: the DM1_like axis we are scoring is real. - Most Paper 9 targets (DDR, DNMT, full SFK, JAK/STAT core) are absent from the local 4007-gene CCLE matrix and need full CCLE expression for proper test. Deferred.

4. DGE layer — strongest signal, all 36 candidates measured

Source: `project/results/dark_matter_phase2/web/data/dge_dm1_vs_dm2.tsv` . 51,711 genes; mean expression per state + log2FC + t + qBH.

Top DM1-enriched candidates (log2FC < 0 means DM1 > DM2):

GENE	CLASS	LOG2FC(DM2/DM1)	T	Q_BH
TACSTD2	TROP2 ADC (non-SL)	-4.14	13.78	8.2e-24
KCNN4	Ion channel	-4.00	19.60	3.8e-32
OSMR	JAK/STAT upstream	-1.44	9.92	1.6e-15
IL6R	JAK/STAT upstream	-1.34	9.17	5.3e-14
MYC	Metabolism (master)	-1.29	7.11	2.1e-09
SRC	SFK	-0.89	8.54	6.7e-12
LYN	SFK	-0.88	9.72	3.2e-15
FYN	SFK	-0.63	5.12	1.4e-05

Top DM2-enriched candidates (DM1-down):

GENE	CLASS	LOG2FC(DM2/DM1)	T	Q_BH
FOXE1	Lineage TF anchor	+1.04	-6.32	8.0e-08
HHEX	Lineage TF anchor	+0.73	-4.48	1.3e-04
PAX8	Lineage TF anchor	+0.64	-4.41	1.9e-04
DNMT3B	Epigenetic	+0.31	-2.62	3.4e-02
IDH2	Metabolism	+0.29	-2.25	7.6e-02
YES1	SFK	+0.21	-3.07	1.1e-02
PARP1	DDR	+0.08	-0.95	0.50
ATR	DDR	+0.08	-1.04	0.46

Reading: - **JAK/STAT upstream receptors (OSMR, IL6R)** are the strongest druggable JAK-axis signal – stronger than STAT3 / JAK1/2 / TYK2 themselves at the bulk level. This suggests the SL window may live at receptor blockade (anti-IL6R / anti-OSMR / dual JAK targeting) rather than downstream STAT3. - **SFK (LYN, SRC, FYN)** is consistently DM1-up and consistent in CCLE direction – a coherent class signal. - **DDR (ATR, PARP1)** is essentially flat at bulk DGE level – the ATR Cox HR signal in section 5 likely reflects state-conditional dependency, not bulk over-expression. Frame ATR as a vulnerability *given* DM1 state, not a DM1 marker. - **Lineage TF DM2-enrichment (FOXE1, HHEX, PAX8)** is the floor sanity check: the axis we are calling "DM1" is the lineage-collapsed state.

5. TCGA-THCA Cox layer — strong powered cohort but very low event rate

n=560 patients, ~18 OS events, event rate 3.2%. TCGA-THCA's well-known issue: small number of OS events. Cox is therefore **underpowered** for most genes; nominal $p < 0.05$ hits are suggestive, not definitive.

5.1 Whole-cohort top hits (univariate Cox per gene)

GENE	CLASS	HR	95% CI	P
TYK2	JAK/STAT	0.15	0.04–0.59	0.006
ATR	DDR	6.88	1.28–36.83	0.024
CHEK2	DDR	0.48	0.24–0.97	0.040
YES1	SFK	3.63	0.97–13.57	0.055
HDAC1	Epigenetic	0.33	0.10–1.04	0.058
MYC	Metabolism	1.50	0.97–2.32	0.070
GLS	Metabolism	1.82	0.93–3.57	0.080

Reading: - **ATR HR=6.88, p=0.024**: high ATR expression → worse OS. Together with the flat DGE direction, this is consistent with DM1 / advanced thyroid being **state-conditionally addicted** to ATR / replication-stress response. Olaparib + ATRi (ceralasertib) combo becomes a coherent Paper 9 candidate even though ATR alone is not DM1-enriched at the bulk level. - **TYK2 HR=0.15, p=0.006**: low TYK2 → worse OS. Not a target – re-frame as a protective modifier of the JAK/STAT axis. Loss-of-TYK2 enriched aggressive disease may indicate TYK2 acting as a brake; this needs confirmation. - **CHEK2 HR=0.48, p=0.040**: protective. DDR loss (low CHEK2) → worse OS. Direction is consistent with DDR-loss-driven aggressive disease; this is consonant with the ATR finding.

5.2 DM1-high subset (n=280) — underpowered

No gene reaches $p < 0.05$ in the DM1-high subset Cox. Top hits at $p < 0.20$ are MYC (1.87, $p=0.05$), HHEX (0.67, $p=0.10$), LDHA (3.68, $p=0.10$), GLS (2.19, $p=0.15$), IDH2 (0.43, $p=0.17$), HDAC1 (0.32, $p=0.17$). Direction is consistent with metabolic-axis (MYC / GLS / LDHA / IDH2) reading, but this layer needs either (a) more events or (b) a meta-analytic combination with GSE76039, which is deferred.

5.3 Honest power statement

With ~18 events, a univariate Cox at $\alpha=0.05$ has roughly 80% power to detect $HR \geq \sim 3.5$ (per +1 SD of expression). Anything weaker is shrunk into noise, so the absence of significance for any specific candidate (e.g., NAMPT $p=0.38$) is **not evidence of irrelevance** – it is a power statement.

6. Triangulation — provisional Paper 9 priority list

Combining the three layers (DGE strong, CCLE direction, Cox direction-or-magnitude), we arrive at a *provisional* priority for the deferred full-execution phase:

RANK	TARGET / CLASS	WHY IT SURVIVES TRIANGULATION	OPEN WORK NEEDED
1	KCNN4	DGE $q=4e-32$ (DM1 $\uparrow$), CCLE $r=+0.29$, well-characterized channel	DepMap CRISPR contrast; senicapoc IC50 in DM1 lines
2	OSMR + IL6R (JAK/STAT upstream)	DGE $q<1e-13$ each (DM1 $\uparrow$); cytokine-receptor SL window	Antibody / soluble-receptor screen; ruxolitinib-class drug response
3	SFK (LYN, SRC, FYN)	All three DGE-up at $q<0.05$; DGE coherent across class	Dasatinib IC50 in DM1 lines; CRISPR LYN/SRC/ FYN dependency
4	ATR + CHEK2 axis (DDR)	Cox HR strong (ATR $p=0.024$, CHEK2 $p=0.040$) despite flat bulk DGE – state-conditional	ATRi (ceralasertib) and PARPi combo IC50 in DM1 lines
5	MYC + metabolic (GLS, LDHA, IDH2)	DGE $q=2e-9$ (MYC), DM1-high Cox trends; classical SL pair logic	GLS1i (CB-839) IC50 + MYC-knockdown in DM1 lines
6	NAMPT / NAPRT (NAD)	CCLE $r=+0.36$ (DM1 $\uparrow$) but DGE NS; partner NAPRT not yet measurable here	DepMap NAMPT dependency $\times$ NAPRT expression
7 (carve-out)	TACSTD2 (TROP2 ADC, non-SL)	DGE $q=8e-24$ (DM1 $\uparrow$), CCLE $r=+0.29$ – strongest DM1 surface marker after KCNN4	Already in Phase B – keep separate from SL framing

Drop / re-frame: - **STAT3 itself** as a single-agent target – DGE log2FC small (q NS); JAK/STAT signal lives at the receptor level (rank 2), not at STAT3. - **TYK2** – flip from "target" to "protective modifier" pending replication. - **KDM1A, HDAC6, EZH2, DNMT3B** – no signal in any of the three layers at this first pass. Keep in panel for full execution but not in the provisional priority list.

7. Caveats and what this first-pass does NOT prove

- 1. CCLE n=13 with a thyroid-only 4007-gene matrix is **not** a synthetic-lethality test – it is a state-axis sanity check. The full DepMap CRISPR contrast remains the gating analysis for Paper 9.
 - 2. TCGA-THCA Cox is **underpowered** at 18 events. Effect sizes from this cohort should be treated as directional priors for GSE76039 + (eventually) Korean K2 + Bundang replication, not standalone evidence.
 - 3. PRISM drug-response stratification is **not** in this first pass. Any drug-class claim is hypothesis-level.
 - 4. SL / SR pair inference is **not** in this first pass. ATR's signal in section 5 is a Cox correlation, not an SL-pair claim.
 - 5. Wet-lab Tier 1 (siRNA / IC50) has not run.
 - 6. None of this overrides the **scope rule**: full execution is gated on Paper 1 in print + marathon close ($\geq$ 2026-06-13) + Yu sign-off.
-

8. Reproducibility

ITEM	VALUE
Script	<code>project/scripts/p9_sl_first_pass.py</code>
Python	3.12 (project/.venv)
Key libs	pandas 2.3, scipy, lifelines 0.30, matplotlib
CCLE source	<code>project/results/v8p1_rigor/f_ccle_validation/{ccle_thyroid_expression,ccle_brs_labels,ccle_thyroid_metadata}.tsv</code>
DGE source	<code>project/results/dark_matter_phase2/web/data/dge_dm1_vs_dm2.tsv</code>
TCGA expr	<code>project/data/raw/TCGA_pancan/pancan_geneExp.gz</code> (UCSC pancan log2(norm+1))
TCGA scored	<code>project_external_st/results/extra/s_tcg_thca_scored.tsv</code>
Random seeds	none required (analytic statistics only)

Re-run: `source project/.venv/bin/activate && python project/scripts/p9_sl_first_pass.py` .

Run summary written to `project/results/paper9_sl_first_pass/summary.json` .

9. What changes for the strategic plan as a result of this first pass

- **F2 / F3 / F6** sketches now have local-data drafts; replace placeholder schematics with current PNGs as the working figure baseline.
- **Section 6 (candidate target classes)** of the strategic plan should reflect the triangulation: KCNN4, OSMR/IL6R, SFK, ATR/CHEK2 promoted; NAMPT and pure STAT3 demoted; TYK2 reframed; TROP2 confirmed as carved-out non-SL.
- **Section 8 (analysis plan)** Step A is now demonstrably executable on local data; Step B (DepMap genome-wide CRISPR) remains the first paper-blocking step in the post-marathon execution phase.

The strategic plan document itself is **not edited in this turn**. Edits to that plan happen only after Paper 1 print + Yu sign-off, per the standing scope rule.