


Supplementary Fig. S2



Supplementary Fig. S2 Enrichment of CD36⁺ pro-inflammatory macrophage in cancer correlates with poor outcome of patients with PTC. **A**, Heatmap showing top marker genes (obtained from unsupervised clustering) used for custom classification of the myeloid compartment. **B**, Comparison of the absolute infiltration proportion of CD36⁺ macrophages between paired peritumor tissues and tumor tissues in the TCGA-THCA

cohort ($n = 58$ clinical samples). Data are represented as mean $\pm$ SEM. A paired Student's t -test was used to assess the difference. $***, P = 0.0006$. **C**, Representative gating strategy of CD36⁺ macrophages in peritumor tissues and tumor tissues from patients with PTC. **D**, IHC staining of SPP1 expression in PTC tissues (scale bar: 50 μ m). **E**, A representative result of IHC staining of SPP1 expression in paired PTC tissue and peritumor tissue. **F**, Bar plot showing proportion of the relative protein levels of SPP1 in 28 paired PTC and peritumor tissues as measured by IHC staining. **G**, A representative result of IHC staining of CD68 expression in paired PTC tissue and peritumor tissue. **H**, Table showing relationship between CD68 expression and clinico-pathologic characteristics in patients with PTC. **I**, Bar plot showing proportion of the relative protein levels of CD68 in 28 paired PTC and peritumor tissues as measured by IHC staining. **J**, Correlation analysis between the expression level of SPP1 and the infiltration level of pro-inflammatory and anti-inflammatory macrophages in thyroid cancer. **K**, Heatmap displaying metabolic pathway activity scores in each macrophage subtype.