

---

# Image Processing

## Lecture Notes: Color Correction


Kai-Lung Hua

---


---

### Color Images

- m Are constructed from three overlaid intensity maps.
- m Each map represents the intensity of a different “primary” color.
- m The actual hues of the primaries do not matter as long as they are distinct.
- m The primaries are 3 vectors (or axes) that form a “basis” of the color space.



## Vector-Valued Pixels



26 March 2020

3



## Color Space for standard digital images

- primary image colors red, green, and blue
  - correspond to R,G, and B axes in color space.
- 8-bits of intensity resolution per color
  - correspond to integers 0 through 255 on axes.
- no negative values
  - color “space” is a cube in the first octant of 3-space.
- color space is discrete
  - $256^3$  possible colors = 16,777,216 elements in cube.

26 March 2020

4



## Color Cube: Faces (outer)



26 March 2020

5

## Color Cube: Faces (inner)





26 March 2020

6

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

7

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

8

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

9

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

10

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

11

---

## Color Cube: Faces (inner and outer)





---

26 March 2020

12


## Color Cube: Faces (inner and outer)




26 March 2020

13

## Different Axis Sets in Color Space



RGB axes




CMY axes

26 March 2020

14

## Color With Respect To Different Axes



The same color has different RGB and CMY coordinates.

26 March 2020

15

## Color Correction

Global changes in the coloration of an image to alter its tint, its hues or the saturation of its colors with minimal changes to its luminant features




26 March 2020

16



## Gamma Adjustment of Color Bands

original




David Peters, producer, and representatives of the IA, The International Alliance of Theatrical Stage Employees, Moving Picture Technicians, Artists and Allied Crafts, on the set of *Frozen Impact* (PorchLight Entertainment, 2003).

26 March 2020

17

## Gamma Adjustment of Color Bands

red  $\gamma=2$




26 March 2020

18

## Gamma Adjustment of Color Bands

original




26 March 2020

19

## Gamma Adjustment of Color Bands

red  $\gamma=0.5$




reduced red = increased cyan

26 March 2020

20

## Gamma Adjustment of Color Bands

original




26 March 2020

21

## Gamma Adjustment of Color Bands

green  $\gamma=2$




26 March 2020

22

## Gamma Adjustment of Color Bands

original




26 March 2020

23

## Gamma Adjustment of Color Bands

green  $\gamma=0.5$




reduced green = incr. magenta

26 March 2020

24

## Gamma Adjustment of Color Bands

original




26 March 2020

25

## Gamma Adjustment of Color Bands

blue  $\gamma=2$




26 March 2020

26



## Gamma Adjustment of Color Bands

original




26 March 2020

27

## Gamma Adjustment of Color Bands

blue  $\gamma=0.5$




reduced blue = incr. yellow

26 March 2020

28

## Color Images


are represented by three bands (not uniquely) *e.g.*, R, G, & B or L,  $a^*$ , &  $b^*$ .



26 March 2020

29


## RGB to LHS: A Perceptual Transformation



26 March 2020

30


## Color Cube: Equi-value Triangle



26 March 2020

31

## Color Cube: Equi-value Triangle




26 March 2020

32




## HSV Color Representation



26 March 2020

33

## Color Point on Equivalence Triangle




26 March 2020

34

---

## Color Vector Associated with Point




---

26 March 2020

35

---

## Color Coordinates and Component Vectors




---

26 March 2020

36

---

## Color Cube, Equivalence Triangle, & Gray Line




---

26 March 2020

37

---

## Color Point and Gray Line




---

26 March 2020

38

---

## Saturation Component of Color Vector




---

26 March 2020

39

---

## Saturation and Value Components of Color Vector




---

26 March 2020

40

---

## Hue, Saturation, and Value




---

26 March 2020

41

---

## Hue and Saturation on Equivalence Plane




---

26 March 2020

42


## Hue, Saturation, and Value with Gray Line



26 March 2020

43

## HSV Color Representation




26 March 2020

44

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

45

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

46

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

47

---

## Equivalence Plane Intersecting Color Cube



---


26 March 2020

48



---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

49

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

50

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

51

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

52

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

53

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

54

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

55

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

56

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

57

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

58

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

59

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

60

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

61

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

62

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

63

---

## Equivalence Plane Intersecting Color Cube



---


26 March 2020

64



---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

65

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

66

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

67

---

## Equivalence Plane Intersecting Color Cube




---

26 March 2020

68

## Equivalence Plane Intersecting Color Cube



26 March 2020

69


## RGB to HSV Conversion

$$\mathbf{v}_0 = \frac{1}{3} \begin{bmatrix} c \\ c \\ c \end{bmatrix}, \text{ where } c = r_0 + g_0 + b_0.$$

$$v_0 = \frac{1}{3} c, \text{ or } v_0 = \|\mathbf{v}_0\| = \frac{\sqrt{3}}{3} c.$$

$$\mathbf{s}_0 = \mathbf{p}_0 - \mathbf{v}_0 = \begin{bmatrix} r_0 - \frac{1}{3}c \\ g_0 - \frac{1}{3}c \\ b_0 - \frac{1}{3}c \end{bmatrix}. \quad \mathbf{p}_0 = \begin{bmatrix} r_0 \\ g_0 \\ b_0 \end{bmatrix}.$$

$$s_0 = \|\mathbf{s}_0\| = \sqrt{\left(r_0 - \frac{1}{3}c\right)^2 + \left(g_0 - \frac{1}{3}c\right)^2 + \left(b_0 - \frac{1}{3}c\right)^2}.$$



26 March 2020

70

## RGB to HSV Conversion


$\begin{bmatrix} c \end{bmatrix}$   
 $c/3$  is the usual value-image intensity (the average of  $r, g, \& b$ ) ...  
 here  $c = r_0 + g_0 + b_0$  ... either def. of  $v_0$  can be used, but  $c/3$  has the advantage of being in the range  $[0, 255]$ .

$$v_0 = \frac{1}{3}c, \text{ or } v_0 = \|\mathbf{v}_0\| = \frac{\sqrt{3}}{3}c.$$

...  $c\sqrt{3}/3$  is the length of the value vector...

$$\mathbf{s}_0 = \mathbf{p}_0 - \mathbf{v}_0 = \begin{bmatrix} r_0 - \frac{1}{3}c \\ g_0 - \frac{1}{3}c \\ b_0 - \frac{1}{3}c \end{bmatrix}, \quad \mathbf{p}_0 = \begin{bmatrix} r_0 \\ g_0 \\ b_0 \end{bmatrix}.$$

$$s_0 = \|\mathbf{s}_0\| = \sqrt{\left(r_0 - \frac{1}{3}c\right)^2 + \left(g_0 - \frac{1}{3}c\right)^2 + \left(b_0 - \frac{1}{3}c\right)^2}.$$



26 March 2020


71

## RGB to HSV Conversion

$$\mathbf{x} = \mathbf{R} - \mathbf{v}_0 = \begin{bmatrix} c \\ 0 \\ 0 \end{bmatrix} - \frac{1}{3} \begin{bmatrix} c \\ c \\ c \end{bmatrix} = \frac{c}{3} \begin{bmatrix} 2 \\ -1 \\ -1 \end{bmatrix}$$

$$\mathbf{s}_0 = \mathbf{p}_0 - \mathbf{v}_0 = \begin{bmatrix} r_0 - \frac{1}{3}c \\ g_0 - \frac{1}{3}c \\ b_0 - \frac{1}{3}c \end{bmatrix}.$$

$$h_0 = \angle(\mathbf{s}_0, \mathbf{x}) = \cos^{-1} \left( \frac{\mathbf{s}_0 \cdot \mathbf{x}}{\|\mathbf{s}_0\| \|\mathbf{x}\|} \right).$$



26 March 2020

47 72

## RGB to HSV Conversion

In summary,

$$v_0 = \frac{1}{3}c, \text{ or } v_0 = \|\mathbf{v}_0\| = \frac{\sqrt{3}}{3}c,$$


where  $c = r_0 + g_0 + b_0$ ,

$$s_0 = \sqrt{\left(r_0 - \frac{1}{3}c\right)^2 + \left(g_0 - \frac{1}{3}c\right)^2 + \left(b_0 - \frac{1}{3}c\right)^2},$$

and

$$h_0 = \cos^{-1}\left(\frac{\mathbf{s}_0 \cdot \mathbf{x}}{\|\mathbf{s}_0\| \|\mathbf{x}\|}\right).$$

Usually,  $s_0$  is normalized to lie in the interval  $(0,1)$  and  $h_0$  is shifted to lie in  $(0,2\pi)$ .



26 March 2020

73

## HSV to RGB Conversion

The equivalue plane is perpendicular to the value vector,  $\mathbf{v}$ .


The plane contains vector  $\mathbf{x}$  defined on slide 45.

Therefore,  $\mathbf{v}$  is perpendicular to  $\mathbf{x}$  and  $\mathbf{y} = \mathbf{v} \times \mathbf{x}$  is also in the plane.

If we keep the directions but ignore the magnitudes, the unit vectors

$$\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \hat{\mathbf{y}} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \hat{\mathbf{v}} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix},$$

form an orthonormal basis with respect to the equivalue plane.



26 March 2020

74

## HSV to RGB Conversion


Given values  $h$ ,  $s$ , and  $v$ , where

$$h \in [0, 2\pi), \quad s \in [0, 1], \quad \text{and} \quad v \in [0, 255],$$

the saturation vector is

$$[\mathbf{s}]_{\text{xyv}} = \begin{bmatrix} s \cos(h) \\ s \sin(h) \\ 0 \end{bmatrix}_{\text{xyv}},$$

with respect to unit vectors  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , and  $\hat{\mathbf{v}}$ , in the equivalence plane.



$$\mathbf{s} = s \cos(h) \hat{\mathbf{x}} + s \sin(h) \hat{\mathbf{y}} + 0 \hat{\mathbf{v}}.$$

26 March 2020

75

## HSV to RGB Conversion


Given values  $h$ ,  $s$ , and  $v$ , where

$$h \in [0, 2\pi), \quad s \in [0, 1], \quad \text{and} \quad v \in [0, 255],$$

the These are the coordinates of  $\mathbf{s}$  with respect to  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , &  $\hat{\mathbf{v}}$ .

$$[\mathbf{s}]_{\text{xyv}} = \begin{bmatrix} s \cos(h) \\ s \sin(h) \\ 0 \end{bmatrix}_{\text{xyv}},$$

with res This is  $\mathbf{s}$  written as a linear combination of vectors  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , &  $\hat{\mathbf{v}}$ . and  $\hat{\mathbf{v}}$ , in the equivalence plane.



$$\mathbf{s} = s \cos(h) \hat{\mathbf{x}} + s \sin(h) \hat{\mathbf{y}} + 0 \hat{\mathbf{v}}.$$

26 March 2020

76

## HSV to RGB Conversion


$\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , and  $\hat{\mathbf{v}}$ , are not in the same directions as the red, green, and blue unit vectors,  $\hat{\mathbf{r}}$ ,  $\hat{\mathbf{g}}$ , and  $\hat{\mathbf{b}}$ .

Therefore,  $[\mathbf{s}]_{\text{xyv}}$  — which we know — is not equal to  $[\mathbf{s}]_{\text{rgb}}$  — which we need in order to find the color,  $\mathbf{p}_0$ , with respect to  $\hat{\mathbf{r}}$ ,  $\hat{\mathbf{g}}$ , and  $\hat{\mathbf{b}}$ .

$$[\mathbf{s}]_{\text{rgb}} = [r_0 \ g_0 \ b_0]^T$$

$$\mathbf{s} \leftrightarrow r_0 \hat{\mathbf{r}} + g_0 \hat{\mathbf{g}} + b_0 \hat{\mathbf{b}},$$

$$\mathbf{s} \leftrightarrow s \cos(h) \hat{\mathbf{x}} + s \sin(h) \hat{\mathbf{y}} + 0 \hat{\mathbf{v}}. \quad \text{We need to find } r_0, g_0, \& b_0.$$



26 March 2020

77

## HSV to RGB Conversion

Vector  $\mathbf{s}$  written as a linear combination of vectors,  $\hat{\mathbf{r}}$ ,  $\hat{\mathbf{g}}$ , and  $\hat{\mathbf{b}}$ , and  $\mathbf{s}$  written as a linear combination of vectors,  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , and  $\hat{\mathbf{v}}$  both refer to the same point on the equi-value plane.


$$\mathbf{s} \leftrightarrow r_0 \hat{\mathbf{r}} + g_0 \hat{\mathbf{g}} + b_0 \hat{\mathbf{b}},$$

$$\mathbf{s} \leftrightarrow s \cos(h) \hat{\mathbf{x}} + s \sin(h) \hat{\mathbf{y}} + 0 \hat{\mathbf{v}}.$$

The specific numbers in  $[\mathbf{s}]_{\text{rgb}}$  and in  $[\mathbf{s}]_{\text{xyv}}$  (that represent the point w.r.t. the two coordinate systems) are, however, different.

$$[\mathbf{s}]_{\text{rgb}} = [r_0 \ g_0 \ b_0]^T \quad \text{and}$$

$$[\mathbf{s}]_{\text{xyz}} = [s \cos(h) \ s \sin(h) \ 0]^T \quad \text{but} \quad [\mathbf{s}]_{\text{rgb}} \neq [\mathbf{s}]_{\text{xyz}}$$



26 March 2020

78


## HSV to RGB Conversion

We can find  $r_0$ ,  $g_0$ , and  $b_0$ , from  $h_0$ ,  $s_0$ , and  $v_0$ , if we know how the unit vectors  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , and  $\hat{\mathbf{v}}$ , are expressed with respect to  $\hat{\mathbf{r}}$ ,  $\hat{\mathbf{g}}$ , and  $\hat{\mathbf{b}}$ . That relationship is in the form of a rotation matrix,  $A$ , such that,

$$[\hat{\mathbf{x}}]_{\text{rgb}} = A[\hat{\mathbf{x}}]_{\text{xyv}}, \quad [\hat{\mathbf{y}}]_{\text{rgb}} = A[\hat{\mathbf{y}}]_{\text{xyv}}, \quad [\hat{\mathbf{v}}]_{\text{rgb}} = A[\hat{\mathbf{v}}]_{\text{xyv}}.$$

Then,

$$\begin{aligned} [\mathbf{s}]_{\text{rgb}} &= A[\mathbf{s}]_{\text{xyv}} \\ &= A[s \cos(h)[\hat{\mathbf{x}}]_{\text{xyv}} + s \sin(h)[\hat{\mathbf{y}}]_{\text{xyv}} + 0[\hat{\mathbf{v}}]_{\text{xyv}}] \\ &= s \cos(h)A[\hat{\mathbf{x}}]_{\text{xyv}} + s \sin(h)A[\hat{\mathbf{y}}]_{\text{xyv}} + 0A[\hat{\mathbf{v}}]_{\text{xyv}} \\ &= s \cos(h)[\hat{\mathbf{x}}]_{\text{rgb}} + s \sin(h)[\hat{\mathbf{y}}]_{\text{rgb}} + 0[\hat{\mathbf{v}}]_{\text{rgb}}. \end{aligned}$$



26 March 2020

79

## HSV to RGB Conversion

When written w.r.t the  $\mathbf{xyz}$  coordinate system we have


$$\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \hat{\mathbf{y}} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \hat{\mathbf{v}} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix},$$

So that,

$$[\hat{\mathbf{x}}]_{\text{rgb}} = A \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad [\hat{\mathbf{y}}]_{\text{rgb}} = A \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad [\hat{\mathbf{v}}]_{\text{rgb}} = A \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

But that implies,

$$A = \begin{bmatrix} [\hat{\mathbf{x}}]_{\text{rgb}} & [\hat{\mathbf{y}}]_{\text{rgb}} & [\hat{\mathbf{v}}]_{\text{rgb}} \end{bmatrix}.$$



26 March 2020

80



## HSV to RGB Conversion

$\hat{\mathbf{v}}$  is the unit vector in the direction  $[1 \ 1 \ 1]^T$  when written w.r.t **rgb** coordinates.


$$[\hat{\mathbf{v}}]_{\text{rgb}} = \frac{\sqrt{3}}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$\hat{\mathbf{x}}$  is perpendicular to  $\hat{\mathbf{v}}$  and has equal  $\hat{\mathbf{g}}$  and  $\hat{\mathbf{b}}$  components.

$$[\hat{\mathbf{x}}]_{\text{rgb}} = \frac{\sqrt{6}}{6} \begin{bmatrix} 2 \\ -1 \\ -1 \end{bmatrix}$$

$\hat{\mathbf{y}}$  is the cross product of  $\hat{\mathbf{v}}$  with  $\hat{\mathbf{x}}$ .

$$\begin{aligned} [\hat{\mathbf{y}}]_{\text{rgb}} &= [\hat{\mathbf{v}}]_{\text{rgb}} \times [\hat{\mathbf{x}}]_{\text{rgb}} \\ &= \frac{\sqrt{2}}{2} \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \end{aligned}$$



26 March 2020

81

## HSV to RGB Conversion

Therefore, the rotation matrix is


$$A = \frac{\sqrt{6}}{6} \begin{bmatrix} 2 & 0 & \sqrt{2} \\ -1 & \sqrt{3} & \sqrt{2} \\ -1 & -\sqrt{3} & \sqrt{2} \end{bmatrix}$$

and

$$\begin{aligned} [\mathbf{s}]_{\text{rgb}} &= s \frac{\sqrt{6}}{6} \cos(h) \begin{bmatrix} 2 \\ -1 \\ -1 \end{bmatrix} + s \frac{\sqrt{2}}{2} \sin(h) \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} + 0 \frac{\sqrt{3}}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \\ &= s \frac{\sqrt{6}}{6} \cos(h) \begin{bmatrix} 2 \\ -1 \\ -1 \end{bmatrix} + s \frac{\sqrt{2}}{2} \sin(h) \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}. \end{aligned}$$

Finally,  $[\mathbf{s}]_{\text{rgb}}$  must be shifted to the value vector to obtain the **rgb** color of  $\mathbf{p}_0$ :

$$\mathbf{p}_0 = [\mathbf{p}]_{\text{rgb}} = [\mathbf{s}]_{\text{rgb}} + [\mathbf{v}]_{\text{rgb}}, \text{ where } \mathbf{s}_0 = [\mathbf{s}]_{\text{rgb}} \text{ and } [\mathbf{v}]_{\text{rgb}} = \mathbf{v}_0 \text{ as def'd. on slide 15.}$$




26 March 2020

82

## Saturation Adjustment

original




26 March 2020

83

## Saturation Adjustment

saturation + 50%




26 March 2020

84

## Saturation Adjustment

original




26 March 2020

85

## Saturation Adjustment


saturation - 50%



26 March 2020

86


## Hue Shifting



26 March 2020

87


## Hue Shifting



26 March 2020

88


## Hue Shifting



26 March 2020

89


## Hue Shifting



26 March 2020

90


## Hue Shifting



26 March 2020

91

## Hue Shifting




26 March 2020

92




## Hue Shifting



26 March 2020

93


## Hue Shifting



26 March 2020

94


## Hue Shifting



26 March 2020

95

## Hue Shifting




26 March 2020

96




## Hue Shifting



26 March 2020

97


## Hue Shifting



26 March 2020

98


## Hue Shifting



26 March 2020

99

## Hue Shifting



26 March 2020

100

## Color Correction Via Transformation

- is a point process; the transformation is applied to each pixel as a function of its color alone.

$$J(r,c) = \Phi[I(r,c)] \quad \forall (r,c) \in \text{supp}(I)$$

- Each pixel is vector valued, therefore the transformation is a vector space operator.

$$I(r,c) = \begin{bmatrix} R_I(r,c) \\ G_I(r,c) \\ B_I(r,c) \end{bmatrix} \quad J(r,c) = \begin{bmatrix} R_J(r,c) \\ G_J(r,c) \\ B_J(r,c) \end{bmatrix} = \Phi\{I(r,c)\} = \Phi\left\{ \begin{bmatrix} R_I(r,c) \\ G_I(r,c) \\ B_I(r,c) \end{bmatrix} \right\}$$

26 March 2020

101

## Linear Transformation of Color



26 March 2020

102

## Color Vector Space Operators

Linear operators  
are matrix  
multiplications

$$\begin{bmatrix} r_1 \\ g_1 \\ b_1 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} r_0 \\ g_0 \\ b_0 \end{bmatrix}$$


$$\begin{bmatrix} r_1 \\ g_1 \\ b_1 \end{bmatrix} = 255 \cdot \begin{bmatrix} (r_0/255)^{1/\gamma_r} \\ (g_0/255)^{1/\gamma_g} \\ (b_0/255)^{1/\gamma_b} \end{bmatrix}$$

Example of a  
nonlinear operator:  
gamma correction

26 March 2020

103


## Linear Transformation of Color



26 March 2020

104


## Linear Transformation of Color



26 March 2020

105


## Linear Transformation of Color



26 March 2020

106

## Linear Transformation of Color



26 March 2020

107

## Color Transformation

Assume  $J$  is a discolored version of image  $I$  such that  $J = \Phi[I]$ . If  $\Phi$  is linear then it is represented by a  $3 \times 3$  matrix,  $A$ :

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}.$$

Then  $J = AI$  or, more accurately,  
 $J(r,c) = AI(r,c)$  for all pixel locations  
 $(r,c)$  in image  $I$ .

26 March 2020

108

## Color Transformation

If at pixel location  $(r, c)$ ,

then  $J(r, c) = AI(r, c)$ , or

$$\text{image } I(r, c) = \begin{bmatrix} \rho_I \\ \gamma_I \\ \beta_I \end{bmatrix} \text{ and}$$

$$\begin{bmatrix} \rho_J \\ \gamma_J \\ \beta_J \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} \rho_I \\ \gamma_I \\ \beta_I \end{bmatrix}$$

$$\text{image } J(r, c) = \begin{bmatrix} \rho_J \\ \gamma_J \\ \beta_J \end{bmatrix},$$

$$= \begin{bmatrix} a_{11}\rho_I + a_{12}\gamma_I + a_{13}\beta_I \\ a_{21}\rho_I + a_{22}\gamma_I + a_{23}\beta_I \\ a_{31}\rho_I + a_{32}\gamma_I + a_{33}\beta_I \end{bmatrix}.$$

26 March 2020

109

## Color Transformation

The inverse transform  $\Phi^{-1}$  (if it exists) maps the discolored image,  $J$ , back into the correctly colored version,  $I$ , i.e.,  $I = \Phi^{-1}[J]$ . If  $\Phi$  is linear then it is represented by the inverse of matrix  $A$ :

$$A^{-1} = (a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31})^{-1} \cdot \begin{bmatrix} a_{22}a_{33} - a_{23}a_{32} & a_{13}a_{32} - a_{12}a_{33} & a_{12}a_{23} - a_{13}a_{22} \\ a_{23}a_{31} - a_{21}a_{33} & a_{11}a_{33} - a_{13}a_{31} & a_{13}a_{21} - a_{11}a_{23} \\ a_{21}a_{32} - a_{22}a_{31} & a_{12}a_{31} - a_{11}a_{32} & a_{11}a_{22} - a_{12}a_{21} \end{bmatrix}.$$

26 March 2020

110

## Color Correction

Assume we know  $n$  colors in the discolored image,  $J$ , that correspond to another set of  $n$  colors (that we also know) in the original image,  $I$ .

$$\left\{ \begin{bmatrix} \rho_{J,k} \\ \gamma_{J,k} \\ \beta_{J,k} \end{bmatrix} \right\}_{k=1}^n \quad \begin{bmatrix} \rho_{J,k} \\ \gamma_{J,k} \\ \beta_{J,k} \end{bmatrix} \leftrightarrow \begin{bmatrix} \rho_{I,k} \\ \gamma_{I,k} \\ \beta_{I,k} \end{bmatrix} \quad \left\{ \begin{bmatrix} \rho_{I,k} \\ \gamma_{I,k} \\ \beta_{I,k} \end{bmatrix} \right\}_{k=1}^n$$

for  $k = 1, \dots, n$ .

known  
wrong  
colors

known  
correspondence

known  
correct  
colors

26 March 2020

111

## Color Correction

To remap the discolored image so that the result matches the original image in a linearly optimal way, we need to find the matrix,  $A$ , that minimizes

$$\mathcal{E}^2 = \sum_{k=1}^n \left\| \begin{bmatrix} \rho_{I,k} \\ \gamma_{I,k} \\ \beta_{I,k} \end{bmatrix} - A^{-1} \begin{bmatrix} \rho_{J,k} \\ \gamma_{J,k} \\ \beta_{J,k} \end{bmatrix} \right\|^2$$

26 March 2020

112



---

## Color Correction

To find the solution of this problem, let

$$Y = \begin{bmatrix} \begin{bmatrix} \rho_{I,1} \\ \gamma_{I,1} \\ \beta_{I,1} \end{bmatrix} & \cdots & \begin{bmatrix} \rho_{I,n} \\ \gamma_{I,n} \\ \beta_{I,n} \end{bmatrix} \end{bmatrix} \quad \text{and} \quad X = \begin{bmatrix} \begin{bmatrix} \rho_{J,1} \\ \gamma_{J,1} \\ \beta_{J,1} \end{bmatrix} & \cdots & \begin{bmatrix} \rho_{J,n} \\ \gamma_{J,n} \\ \beta_{J,n} \end{bmatrix} \end{bmatrix}.$$

Then  $X$  and  $Y$  are known  $3 \times n$  matrices such that

$$Y \approx A^{-1}X,$$

where  $A$  is the  $3 \times 3$  matrix that we want to find.

---

26 March 2020

113

---

## Color Correction

The linearly optimal solution is the least mean squared solution that is given by

$$B = A^{-1} = YX^T (XX^T)^{-1}$$

where  $X^T$  represents the transpose of matrix  $X$ .

- Notes:
1.  $n$ , the number of color pairs, must be  $\geq 3$ ,
  2.  $XX^T$  must be invertible, *i.e.*,  $\text{rank}(XX^T) = 3$ ,
  3. If  $n=3$ , then  $X^T(XX^T)^{-1} = X^{-1}$ .

---

26 March 2020

114

## Color Correction

The linearly optimal solution that is given by

input colors (to be changed):

$$\begin{bmatrix} \rho_{j,1} \\ \gamma_{j,1} \\ \beta_{j,1} \end{bmatrix} \cdots \begin{bmatrix} \rho_{j,n} \\ \gamma_{j,n} \\ \beta_{j,n} \end{bmatrix} \cdot \begin{bmatrix} \rho_{j,1} & \gamma_{j,1} & \beta_{j,1} \\ \vdots & \vdots & \vdots \\ \rho_{j,n} & \gamma_{j,n} & \beta_{j,n} \end{bmatrix}$$

$$B = A^{-1} = YX^T (XX^T)^{-1}$$

where  $X^T$  represents

output colors (wanted):

$$\begin{bmatrix} \rho_{i,1} \\ \gamma_{i,1} \\ \beta_{i,1} \end{bmatrix} \cdots \begin{bmatrix} \rho_{i,n} \\ \gamma_{i,n} \\ \beta_{i,n} \end{bmatrix}$$

of matrix  $X$ .

- Notes:
1.  $n$ , or pairs, must be  $\geq 3$ ,
  2.  $XX^T$  must be invertible, i.e.,  $\text{rank}(XX^T) = 3$ ,
  3. If  $n=3$ , then  $X^T(XX^T)^{-1} = X^{-1}$ .

26 March 2020

115

## Color Correction

The linearly optimal solution that is given by

input colors (to be changed):

$$\begin{bmatrix} \rho_{j,1} \\ \gamma_{j,1} \\ \beta_{j,1} \end{bmatrix} \cdots \begin{bmatrix} \rho_{j,n} \\ \gamma_{j,n} \\ \beta_{j,n} \end{bmatrix} \cdot \begin{bmatrix} \rho_{j,1} & \gamma_{j,1} & \beta_{j,1} \\ \vdots & \vdots & \vdots \\ \rho_{j,n} & \gamma_{j,n} & \beta_{j,n} \end{bmatrix}$$

$$B = A^{-1} = YX^T (XX^T)^{-1}$$

where  $X^T$  represents

output colors (wanted):

$$\begin{bmatrix} \rho_{i,1} \\ \gamma_{i,1} \\ \beta_{i,1} \end{bmatrix} \cdots \begin{bmatrix} \rho_{i,n} \\ \gamma_{i,n} \\ \beta_{i,n} \end{bmatrix}$$

of matrix  $X$ .

- Notes:
1.  $n$ , or pairs, must be  $\geq 3$ ,
  2.  $XX^T$  must be invertible, i.e.,  $\text{rank}(XX^T) = 3$ ,
  3. If  $n=3$ , then  $X^T(XX^T)^{-1} = X^{-1}$ .

26 March 2020

116

---

## Color Correction

Then the image is color corrected by performing

$$I(r,c) = B J(r,c), \text{ for all } (r,c) \in \text{supp}(J).$$

In Matlab this is easily performed by

```
I = reshape(((B*(reshape(J,R*C,3))')'),R,C,3);
```

where  $B=A^{-1}$  is computed directly through the LMS formula on the previous page, and  $R$  &  $C$  are the number of rows and columns in the image.

---

26 March 2020

117

---

## Linear Color Correction

NASA Summer Faculty Fellows at Ellington Air Force Base, Houston, TX, July 2002. Airplane is a T-38.



Original Image



"Aged" Image

---

26 March 2020

118

## Color Mapping 1



Original Image

"Aged" Image

26 March 2020

119

## Color Mapping 2



Original Image

"Aged" Image

26 March 2020

120

## Color Mapping 3



Original Image

"Aged" Image

26 March 2020

121

## Color Mapping 4




Original Image

"Aged" Image

26 March 2020

122

## Color Transformations




The aging process was a transformation,  $\Phi$ , that mapped:

$$\begin{bmatrix} 17 \\ 122 \\ 114 \end{bmatrix} = \Phi \left\{ \begin{bmatrix} 17 \\ 121 \\ 171 \end{bmatrix} \right\} \quad \begin{bmatrix} 222 \\ 222 \\ 185 \end{bmatrix} = \Phi \left\{ \begin{bmatrix} 222 \\ 222 \\ 218 \end{bmatrix} \right\} \quad \begin{bmatrix} 240 \\ 171 \\ 103 \end{bmatrix} = \Phi \left\{ \begin{bmatrix} 240 \\ 171 \\ 160 \end{bmatrix} \right\} \quad \begin{bmatrix} 236 \\ 227 \\ 106 \end{bmatrix} = \Phi \left\{ \begin{bmatrix} 240 \\ 230 \\ 166 \end{bmatrix} \right\}$$

26 March 2020

123

## Color Transformations



To undo the process we need to find,  $\Phi^{-1}$ , that maps:

$$\begin{bmatrix} 17 \\ 121 \\ 171 \end{bmatrix} = \Phi^{-1} \left\{ \begin{bmatrix} 17 \\ 122 \\ 114 \end{bmatrix} \right\} \quad \begin{bmatrix} 222 \\ 222 \\ 218 \end{bmatrix} = \Phi^{-1} \left\{ \begin{bmatrix} 222 \\ 222 \\ 185 \end{bmatrix} \right\} \quad \begin{bmatrix} 240 \\ 171 \\ 160 \end{bmatrix} = \Phi^{-1} \left\{ \begin{bmatrix} 240 \\ 171 \\ 103 \end{bmatrix} \right\} \quad \begin{bmatrix} 240 \\ 230 \\ 166 \end{bmatrix} = \Phi^{-1} \left\{ \begin{bmatrix} 236 \\ 227 \\ 106 \end{bmatrix} \right\}$$

26 March 2020

124

## Correction Using 3 Mappings

$$B = A^{-1} = YX^{-1}$$



$$X = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 122 & 171 \\ 185 & 114 & 103 \end{bmatrix}$$



$$Y = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 121 & 171 \\ 218 & 171 & 160 \end{bmatrix}$$

26 March 2020

125

## Correction Using 3 Mappings

$$B = A^{-1} = YX^{-1}$$

original



$$X = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 122 & 171 \\ 185 & 114 & 103 \end{bmatrix}$$

corrected



$$Y = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 121 & 171 \\ 218 & 171 & 160 \end{bmatrix}$$

26 March 2020

126



## Another Correction Using 3 Mappings

$$B = A^{-1} = YX^{-1}$$



$$X = \begin{bmatrix} 222 & 17 & 236 \\ 222 & 122 & 227 \\ 185 & 114 & 106 \end{bmatrix}$$



$$Y = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 121 & 230 \\ 218 & 171 & 166 \end{bmatrix}$$

26 March 2020

127

## Another Correction Using 3 Mappings

$$B = A^{-1} = YX^{-1}$$

original



$$X = \begin{bmatrix} 222 & 17 & 236 \\ 222 & 122 & 227 \\ 185 & 114 & 106 \end{bmatrix}$$

corrected



$$Y = \begin{bmatrix} 222 & 17 & 240 \\ 222 & 121 & 230 \\ 218 & 171 & 166 \end{bmatrix}$$

26 March 2020

128



## Correction Using All 4 Mappings

$$B = A^{-1} = YX^T(XX^T)^{-1}$$



$$X = \begin{bmatrix} 222 & 17 & 236 & 240 \\ 222 & 122 & 227 & 171 \\ 185 & 114 & 106 & 103 \end{bmatrix}$$



$$Y = \begin{bmatrix} 222 & 17 & 240 & 240 \\ 222 & 121 & 230 & 171 \\ 218 & 171 & 166 & 160 \end{bmatrix}$$

26 March 2020

129

## Correction Using All 4 Mappings

$$B = A^{-1} = YX^T(XX^T)^{-1}$$

original

corrected



$$X = \begin{bmatrix} 222 & 17 & 236 & 240 \\ 222 & 122 & 227 & 171 \\ 185 & 114 & 106 & 103 \end{bmatrix}$$



$$Y = \begin{bmatrix} 222 & 17 & 240 & 240 \\ 222 & 121 & 230 & 171 \\ 218 & 171 & 166 & 160 \end{bmatrix}$$

26 March 2020

130

## Random Sampling of Color Values

```
>> rr = round(R*rand([1 n]));
>> rc = round(C*rand([1 n]));
>> idx = [rr;rc];
>> Y(:,1) = diag(I(rr,rc,1));
>> Y(:,2) = diag(I(rr,rc,2));
>> Y(:,3) = diag(I(rr,rc,3));
>> X(:,1) = diag(J(rr,rc,1));
>> X(:,2) = diag(J(rr,rc,2));
>> X(:,3) = diag(J(rr,rc,3));
```

R = number of rows in image  
C = number of columns in image  
n = number of pixels to select

rand([1 n]): 1 × n matrix  
of random numbers  
between 0 and 1.

diag(I(rr,rc,1)): vector  
from main diagonal of  
matrix I(rr,rc,1).

26 March 2020

131

## Correction Using 128 Mappings

$$B = A^{-1} = YX^T (XX^T)^{-1}$$



$$X = \begin{bmatrix} 111 & 235 \\ 103 & \dots & 233 \\ 22 & & 210 \end{bmatrix}$$


$$Y = \begin{bmatrix} 111 & 234 \\ 102 & \dots & 233 \\ 71 & & 229 \end{bmatrix}$$

26 March 2020

132

## Correction Using 128 Mappings

$$B = A^{-1} = YX^T (XX^T)^{-1}$$



26 March 2020


133

## Correction Using 128 Mappings

$$B = A^{-1} = YX^T (XX^T)^{-1}$$

original

corrected



26 March 2020

134

## Correction Using 4 Mappings

$$B = A^{-1} = YX^T (XX^T)^{-1}$$

original



corrected



$$X = \begin{bmatrix} 222 & 17 & 236 & 240 \\ 222 & 122 & 227 & 171 \\ 185 & 114 & 106 & 103 \end{bmatrix}$$

$$Y = \begin{bmatrix} 222 & 17 & 240 & 240 \\ 222 & 121 & 230 & 171 \\ 218 & 171 & 166 & 160 \end{bmatrix}$$

26 March 2020

135

## Linear Color Transformation Program

```
function J = LinTrans(I,A)

[R C B] = size(I);

I = double(I);

J = reshape(((A*(reshape(I,R*C,3))')'),R,C,3);

J = uint8(J);

return;
```

26 March 2020

136