


PAPER

Hofmann-Mislove type definitions of non-Hausdorff spaces[†]

Chong Shen^{1*} , Xiaoyong Xi², Xiaoquan Xu³ and Dongsheng Zhao⁴

¹School of Science, Beijing University of Posts and Telecommunications, Beijing, China, ²School of Mathematics and Statistics, Yancheng Teachers University, Jiangsu, Yancheng, China, ³School of Mathematics and Statistics, Minnan Normal University, Fujian, Zhangzhou, China and ⁴Mathematics and Mathematics Education, National Institute of Education, Nanyang Technological University, 1 Nanyang Walk, Singapore

*Corresponding author. Email: shenchong0520@163.com

(Received 26 August 2021; revised 18 May 2022; accepted 2 June 2022; first published online 27 June 2022)

Abstract

One of the most important results in domain theory is the Hofmann-Mislove Theorem, which reveals a very distinct characterization for the sober spaces via open filters. In this paper, we extend this result to the d -spaces and well-filtered spaces. We do this by introducing the notions of Hofmann-Mislove-system (HM-system for short) and Ψ -well-filtered space, which provide a new unified approach to sober spaces, well-filtered spaces, and d -spaces. In addition, a characterization for Ψ -well-filtered spaces is provided via Ψ -sets. We also discuss the relationship between Ψ -well-filtered spaces and H -sober spaces considered by Xu. We show that the category of complete Ψ -well-filtered spaces is a full reflective subcategory of the category of T_0 spaces with continuous mappings. For each HM-system Ψ that has a designated property, we show that a T_0 space X is Ψ -well-filtered if and only if its Smyth power space $P_s(X)$ is Ψ -well-filtered.

Keywords: d -space; H -sober; Hofmann-Mislove Theorem; Ψ -well-filtered space; well-filtered space; sober space

1. Introduction

Traditionally, topologists were interested in Hausdorff spaces much more than non-Hausdorff spaces. The development of domain theory has inspired the heavy interests in non-Hausdorff spaces. Sober spaces, well-filtered spaces, and d -spaces are three of the mostly well-studied non-Hausdorff spaces in domain theory. Recent researches revealed that these three classes of spaces share quite a number of common properties: (i) their categories are reflective in the category of T_0 spaces (Eršov 1999; Hoffmann 1981; Liu et al. 2020; Wu et al. 2020; Wyler 1981; Xu et al. 2020a); (ii) they can be defined in terms of special subsets (i.e., irreducible sets, KF-sets, directed sets) (Shen et al. 2019); (iii) they are preserved under Cartesian product (Hoffmann 1979; Keimel and Lawson 2009); (iv) their open sets are Scott open in the specialization order (Gierz et al. 2003; Goubault-Larrecq 2003). Recently, Li, Yuan, and Zhao (2021) introduced the notion of Θ -fine space, which provided another unified approach to such properties. Also, using a generalized notion of Rudin set (KF-set), directed set, and irreducible set in T_0 spaces, Xu (2021)

[†] This research was supported by the National Natural Science Foundation of China (Nos. 1210010153, 12071199, 12071188, 11661057, 11871097), Jiangsu Provincial Department of Education (21KJB110008) and the Natural Science Foundation of Jiangxi Province, China (No. 20192ACBL20045).

introduced the notion of H -sober space and particularly proved that the category of H -sober spaces is a full reflective subcategory of the category of T_0 spaces.

Sober spaces have very rich properties. The single most important result about such spaces is the Hofmann-Mislove Theorem, which states that sober spaces are exactly the spaces such that there is a natural correspondence between the open filters of the lattice of open subsets and the compact saturated subsets.

In this paper, we shall extend the Hofmann-Mislove Theorem to a general class of spaces, including sober spaces, well-filtered spaces, and d -spaces. One byproduct of this approach is the finding of some new classes of T_0 spaces.

Here is the outline of the paper. In Section 3, we introduce the Ψ -well-filtered spaces, where Ψ is a property on open filters. We then show that sober spaces, well-filtered spaces, and d -spaces are all special types of Ψ -well-filtered spaces. In Section 4, we introduce the notion of Ψ -set and use it to characterize Ψ -well-filtered spaces. In Section 5, the interlink between H -sober spaces and Ψ -well-filtered spaces is discussed. Especially, it is shown that the complete Ψ -well-filtered spaces are exactly the H_Ψ -sober spaces, where H_Ψ is an R -subset system induced by Ψ . As an immediate result, the category of complete Ψ -well-filtered spaces is reflective in the category of T_0 spaces. In the last section, for an HM-system Ψ and a T_0 space X , we show that the Smyth power space $P_s(X)$ is Ψ -well-filtered if and only if X is Ψ -well-filtered and Ψ has property Q for X . As a corollary, it is deduced that a T_0 space X is sober (resp., well-filtered) if and only if $P_s(X)$ is sober (resp., well-filtered).

2. Preliminary

Next, we introduce some basic concepts and notations that will be used in the paper. For more details, see Engelking (1989), Gierz *et al.* (2003), Goubault-Larrecq (2003).

Let P be a poset. A nonempty subset D of P is *directed* (resp., *filtered*) if every two elements in D have an upper (resp., lower) bound in D . P is called a *directed complete poset* or a *dcpo* for short, if for any directed subset $D \subseteq P$, the supremum of D , denoted by $\bigvee D$, exists.

For any subset A of a poset P , we use the following standard notations:

$$\uparrow A = \{y \in P : \exists x \in A, x \leq y\}; \downarrow A = \{y \in P : \exists x \in A, y \leq x\}.$$

In particular, for each $x \in X$, we write $\uparrow x = \uparrow \{x\}$ and $\downarrow x = \downarrow \{x\}$. A subset A of P is called a *lower* (resp., *upper*) *set* if $A = \downarrow A$ (resp., $A = \uparrow A$).

A subset U of P is *Scott open* if (i) $U = \uparrow U$ and (ii) for any directed subset D of P for which $\bigvee D$ exists, $\bigvee D \in U$ implies $D \cap U \neq \emptyset$. All Scott open subsets of P form a topology, called the *Scott topology* on P , denoted by $\sigma(P)$. The space $\Sigma P = (P, \sigma(P))$ is called the *Scott space* of P .

Let X be a T_0 space. A subset A of X is called *saturated* if A equals the intersection of all open sets containing it. The specialization order \leq on X is defined by $x \leq y$ if and only if $x \in \text{cl}(\{y\})$, where cl is the closure operator. It is important to note that a subset A of X is saturated if and only if $A = \uparrow A$ in the specialization order.

Remark 2.1. Let X be a T_0 space.

- (1) Every open (resp., closed) set is an upper (resp., lower) set. In particular, $\text{cl}(\{x\}) = \downarrow x$ (Gierz *et al.* 2003; Goubault-Larrecq 2003).
- (2) For each subset K of X , K is compact if and only if $\uparrow K$ is compact (Gierz *et al.* 2003; Goubault-Larrecq 2003).
- (3) It is well-known that a subset K of X is compact saturated if and only if $\min K$ is compact and $K = \uparrow \min K$, where $\min K$ is the set of all minimal elements of K in the specialization order (see, e.g., Ern  2009, pp. 2068).

Definition 2.2 (Gierz et al. 2003; Goubault-Larrecq 2003). (1) A nonempty subset A of a topological space X is called *irreducible* if for any closed sets F_1, F_2 of X , $A \subseteq F_1 \cup F_2$ implies $A \subseteq F_1$ or $A \subseteq F_2$.
 (2) A T_0 space X is called *sober* if every irreducible closed subset of X is the closure of a (unique) point.

For a T_0 space X , $A \subseteq X$, and $\mathcal{F} \subseteq 2^X$, we use the following notations:

- $\mathcal{O}(X)$, the family of all open subsets of X ;
- $\mathcal{C}(X)$, the family of all closed subsets of X ;
- $\mathcal{Q}(X)$, the family of all compact saturated subsets of X ;
- $\mathcal{N}(A)$, the family $\{U \in \mathcal{O}(X) : A \subseteq U\}$;
- $\mathcal{M}(A)$, the family $\{U \in \mathcal{O}(X) : A \cap U \neq \emptyset\}$;
- $\mathfrak{M}(\mathcal{F})$, the family $\{C \in \mathcal{C}(X) : \forall F \in \mathcal{F}, C \cap F \neq \emptyset\}$;
- $m(\mathcal{F})$, the family of all minimal members in $(\mathfrak{M}(\mathcal{F}), \subseteq)$.

Definition 2.3 (Gierz et al. 2003; Goubault-Larrecq 2003). A T_0 space X is called *well-filtered* if for any filtered family \mathcal{F} of $\mathcal{Q}(X)$ and $U \in \mathcal{O}(X)$, $\bigcap \mathcal{F} \subseteq U$ implies $K \subseteq U$ for some $K \in \mathcal{F}$.

Definition 2.4 (Shen et al. 2019; Xu et al. 2020a). Let X be a T_0 space. A nonempty subset A of X is called a *KF-set* (or a *Rudin set*), if there exists a filtered family \mathcal{F} of $\mathcal{Q}(X)$ such that $\text{cl}(A) \in m(\mathcal{F})$.

Theorem 2.5 (Shen et al. 2019; Xu et al. 2020a). A T_0 space X is well-filtered if and only if for each KF-set A , there exists $x \in X$ such that $\text{cl}(A) = \text{cl}(\{x\})$.

Lemma 2.6 (Shen et al. 2019). Let X and Y be two T_0 spaces, and $f : X \rightarrow Y$ be a continuous mapping. If A is a KF-set in X , then $f(A)$ is a KF-set in Y .

Definition 2.7 (Gierz et al. 2003; Goubault-Larrecq 2003). A T_0 space X is called a *d-space*, if X is a dcpo and every open subset of X is Scott open in the specialization order.

Proposition 2.8 (Xu et al. 2020a, Proposition 3.3). A T_0 space X is a *d-space* if and only if for each directed subset D of X , there is $x \in X$ such that $\text{cl}(D) = \text{cl}(\{x\})$.

Remark 2.9. Every sober space is well-filtered, and every well-filtered space is a *d-space* (Gierz et al. 2003; Goubault-Larrecq 2003).

Definition 2.10 (Gierz et al. 2003; Goubault-Larrecq 2003). Let X be a T_0 space, and $\mathcal{F} \subseteq \mathcal{O}(X)$. Then, \mathcal{F} is called an *open filter*, if it is filtered and Scott open in $(\mathcal{O}(X), \subseteq)$. We denote by $\text{OF}(X)$ the family of all open filters of $\mathcal{O}(X)$.

Remark 2.11. For a T_0 space X , the following results can be checked easily.

- (1) For each compact saturated subset K of X , $\mathcal{N}(K) \in \text{OF}(X)$.
- (2) For each irreducible subset A of X , $\mathcal{M}(A) \in \text{OF}(X)$.
- (3) For each continuous mapping $f : X \rightarrow Y$ between T_0 spaces X and Y , if $\mathcal{F} \in \text{OF}(X)$, then $f_*(\mathcal{F}) = \{V \in \mathcal{O}(Y) : f^{-1}(V) \in \mathcal{F}\} \in \text{OF}(Y)$.

The following is a similar result to the Topological Rudin Lemma in Heckmann and Keimel (2013).

Lemma 2.12. *Let X be a T_0 space, $A \in \mathcal{C}(X)$ and $\mathcal{F} \in \text{OF}(X)$. Then, the following conditions hold:*

- (1) *every element of $m(\mathcal{F})$ is irreducible;*
- (2) *if $A \in \mathfrak{M}(\mathcal{F})$, then there is a closed subset C of A such that $C \in m(\mathcal{F})$.*

Proof. (1) Assume on the contrary there is $C \in m(\mathcal{F})$ that is not irreducible. Then, there exist closed sets C_1, C_2 such that $C = C_1 \cup C_2$ but $C \neq C_1$ and $C \neq C_2$. Since C_1, C_2 are proper subsets of C , by the minimality of C there exist $U_1, U_2 \in \mathcal{F}$ such that $C_1 \cap U_1 = \emptyset$ and $C_2 \cap U_2 = \emptyset$. Since \mathcal{F} is a filter, there exists $U_3 \in \mathcal{F}$ such that $U_3 \subseteq U_1 \cap U_2$. It follows that $C_1 \cap U_3 = C_2 \cap U_3 = \emptyset$. Thus, $C \cap U_3 = (C_1 \cup C_2) \cap U_3 = (C_1 \cap U_3) \cup (C_2 \cap U_3) = \emptyset$, contradicting the fact that $C \in m(\mathcal{F})$. Therefore, C is irreducible.

(2) Note that an open set is not in \mathcal{F} if and only if its complement is in $\mathfrak{M}(\mathcal{F})$. Then, using Proof (i) of Lemma II-1.19 in Gierz et al. (2003), condition (2) holds dually. \square

3. Ψ -Well-Filtered Spaces

For each $K \in \mathcal{Q}(X)$, it is trivial that $\mathcal{N}(K) \in \text{OF}(X)$ and $K = \bigcap \mathcal{N}(K)$. Then, the mapping $\mathcal{N} : (\mathcal{Q}(X), \supseteq) \rightarrow (\text{OF}(X), \subseteq)$, $K \mapsto \mathcal{N}(K)$ is well-defined and clearly is an order-embedding. In domain theory, the single most important result about sober spaces is the Hofmann-Mislove Theorem (see Gierz et al. 2003, Theorems II-1.20, II-1.21).

Theorem 3.1 (Hofmann-Mislove Theorem). *For a T_0 space X , the following conditions are equivalent:*

- (1) *X is sober;*
- (2) *$\forall \mathcal{F} \in \text{OF}(X)$, there is a $K \in \mathcal{Q}(X)$ such that $\mathcal{F} = \mathcal{N}(K)$;*
- (3) *$\forall \mathcal{F} \in \text{OF}(X)$, $\mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$.*

As a corollary of the Hofmann-Mislove Theorem, the following result is clear.

Corollary 3.2. *For a T_0 space X , the following conditions are equivalent:*

- (1) *X is sober;*
- (2) *$\text{OF}(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$;*
- (3) *$\forall \mathcal{F} \in \text{OF}(X)$, $\forall U \in \mathcal{O}(X)$, $\bigcap \mathcal{F} \subseteq U$ implies $U \in \mathcal{F}$.*

In the following, we would like to provide a unified characterization of Hofmann-Mislove Theorem type for the classes of d -spaces and well-filtered spaces via open filters.

Definition 3.3. *A covariant functor $\Psi : \mathbf{Top}_0 \rightarrow \mathbf{Set}$ is called a Hofmann-Mislove system (an HM-system for short) on \mathbf{Top}_0 if the following two conditions are satisfied:*

- (HM1) *for each T_0 space X , $\{\mathcal{N}(K) : K \in \mathcal{Q}(X)\} \subseteq \Psi(X) \subseteq \text{OF}(X)$;*
- (HM2) *for each continuous mapping $f : X \rightarrow Y$ in \mathbf{Top}_0 , $\Psi(f)(\mathcal{F}) = f_*(\mathcal{F}) = \{V \in \mathcal{O}(Y) : f^{-1}(V) \in \mathcal{F}\} \in \Psi(Y)$ for each $\mathcal{F} \in \Psi(X)$.*

Definition 3.4. *Let Ψ be an HM-system and X be a T_0 space. We call X a Ψ -well-filtered space, if $\forall \mathcal{F} \in \Psi(X)$, $\forall U \in \mathcal{O}(X)$,*

$$\bigcap \mathcal{F} \subseteq U \text{ implies } U \in \mathcal{F}.$$

Lemma 3.5. *Let Ψ be an HM-system, X be a Ψ -well-filtered space, and $\mathcal{F} \in \Psi(X)$. Then $\bigcap \mathcal{F} \in \mathcal{Q}(X)$, and $\emptyset \notin \mathcal{F}$ implies $\bigcap \mathcal{F} \neq \emptyset$.*

Proof. Let $\{U_i : i \in I\}$ be a directed family of $\mathcal{O}(X)$ such that $\bigcap \mathcal{F} \subseteq \bigcup_{i \in I} U_i$. Since X is Ψ -well-filtered, $\bigcup_{i \in I} U_i \in \mathcal{F}$, and since \mathcal{F} is Scott open, there exists $i_0 \in I$ such that $U_{i_0} \in \mathcal{F}$, which implies $\bigcap \mathcal{F} \subseteq U_{i_0}$. Hence, $\bigcap \mathcal{F}$ is compact and clearly is saturated. In addition, if $\bigcap \mathcal{F} = \emptyset$, then $\emptyset \in \mathcal{F}$ because X is Ψ -well-filtered, completing the proof. \square

Theorem 3.6. *Let Ψ be an HM-system and X be a T_0 space. Then, the following conditions are equivalent:*

- (1) X is Ψ -well-filtered;
- (2) $\forall \mathcal{F} \in \Psi(X), \mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$;
- (3) $\Psi(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$.

Proof. (1) \Rightarrow (2). Let $\mathcal{F} \in \Psi(X)$. It is clear that $\mathcal{F} \subseteq \mathcal{N}(\bigcap \mathcal{F})$. Conversely, if $U \in \mathcal{N}(\bigcap \mathcal{F})$, then $\bigcap \mathcal{F} \subseteq U$ and $U \in \mathcal{O}(X)$, and since X is Ψ -well-filtered, we have that $U \in \mathcal{F}$. This shows that $\mathcal{N}(\bigcap \mathcal{F}) \subseteq \mathcal{F}$. Hence, $\mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$.

(2) \Rightarrow (3). By the definition of Ψ , it is clear that $\{\mathcal{N}(K) : K \in \mathcal{Q}(X)\} \subseteq \Psi(X)$. Now suppose $\mathcal{F} \in \Psi(X)$. By Lemma 3.5, $K_0 = \bigcap \mathcal{F} \in \mathcal{Q}(X)$, and by condition (2) $\mathcal{F} = \mathcal{N}(K_0)$. This shows that $\Psi(X) \subseteq \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$. Thus, condition (3) holds.

(3) \Rightarrow (1). Let $\mathcal{F} \in \Psi(X)$ and $U \in \mathcal{O}(X)$ such that $\bigcap \mathcal{F} \subseteq U$. By (3), there exists $K \in \mathcal{Q}(X)$ such that $\mathcal{F} = \mathcal{N}(K)$. Then, $K = \bigcap \mathcal{N}(K) = \bigcap \mathcal{F} \subseteq U$, which implies that $U \in \mathcal{N}(K) = \mathcal{F}$. Therefore, X is Ψ -well-filtered. \square

The following result shows that the Ψ -well-filteredness is a topological property for each HM-system Ψ .

Proposition 3.7. *Let Ψ be an HM-system, X be a Ψ -well-filtered space, and Y be a T_0 space. If Y is homeomorphic to X , then Y is a Ψ -well-filtered space.*

Proof. Suppose $h : Y \rightarrow X$ is a homeomorphism. Let $\mathcal{F} \in \Psi(Y)$ and $W \in \mathcal{O}(Y)$ such that $\bigcap \mathcal{F} \subseteq W$. Since h is a homeomorphism, one can easily obtain that $h_*(\mathcal{F}) = \{h(U) : U \in \mathcal{F}\}$, which implies that

$$\bigcap h_*(\mathcal{F}) = \bigcap \{h(U) : U \in \mathcal{F}\} = h\left(\bigcap \mathcal{F}\right) \subseteq h(W) \in \mathcal{O}(X).$$

By (HM2), $h_*(\mathcal{F}) \in \Psi(X)$, and since X is Ψ -well-filtered, we have that $h(W) \in h_*(\mathcal{F})$, so $W \in \mathcal{F}$. Therefore, X is Ψ -well-filtered. \square

In the following, we will show that all the classes of sober spaces, well-filtered spaces, and d -spaces can be characterized via HM-systems.

Definition 3.8. Define $\Psi_{\text{sob}}, \Psi_{\text{wf}}, \Psi_d : \mathbf{Top}_0 \rightarrow \mathbf{Set}$ as follows: for each T_0 space X ,

$$\Psi_{\text{sob}}(X) = \text{OF}(X),$$

$$\Psi_{\text{wf}}(X) = \left\{ \bigcup_{K \in \mathcal{G}} \mathcal{N}(K) : \mathcal{G} \text{ is a filtered family of } \mathcal{Q}(X) \right\},$$

$$\Psi_d(X) = \left\{ \bigcup_{x \in D} \mathcal{N}(\uparrow x) : D \text{ is a directed subset of } X \right\} \cup \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}.$$

Then, it is trivial to check that $\Psi_{\text{sob}}, \Psi_{\text{wf}}, \Psi_d$ are all HM-systems.

Theorem 3.9. *Let X be a T_0 space. The following conditions are equivalent:*

- (1) X is sober;
- (2) X is Ψ_{sob} -well-filtered;
- (3) $\forall \mathcal{F} \in \Psi_{\text{sob}}(X)$, $\mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$;
- (4) $\Psi_{\text{sob}}(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$.

Proof. It is straightforward by the Hofmann-Mislove Theorem. □

Remark 3.10. Let $\Psi_0(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$ for each T_0 space X . Then, it is easy to check that Ψ_0 is an HM-system, and it is clear that each T_0 space X is Ψ_0 -well-filtered. From Theorem 3.9, for each HM-system Ψ , we have the following relations:

$$\text{sober } (\Psi_{\text{sob}}\text{-well-filtered}) \text{ space} \implies \Psi\text{-well-filtered space} \implies T_0 \text{ } (\Psi_0\text{-well-filtered}) \text{ space}.$$

In another words, the sober space is the strongest Ψ -well-filtered space, and the T_0 space is the weakest one.

Theorem 3.11. *Let X be a T_0 space. The following conditions are equivalent:*

- (1) X is well-filtered;
- (2) X is Ψ_{wf} -well-filtered;
- (3) $\forall \mathcal{F} \in \Psi_{\text{wf}}(X)$, $\mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$;
- (4) $\Psi_{\text{wf}}(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$.

Proof. That (2) \Leftrightarrow (3) \Leftrightarrow (4) follows immediately from Theorem 3.6.

(1) \Rightarrow (2). Suppose X is well-filtered. Let $\mathcal{F} \in \Psi_{\text{wf}}(X)$ and $U \in \mathcal{O}(X)$ such that $\bigcap \mathcal{F} \subseteq U$. Then, there exists a filtered family $\{K_i : i \in I\} \subseteq \mathcal{Q}(X)$ such that $\mathcal{F} = \bigcup_{i \in I} \mathcal{N}(K_i)$. Note that $\bigcap_{i \in I} K_i = \bigcap \mathcal{F} \subseteq U$, which implies that $K_{i_0} \subseteq U$ for some $i_0 \in I$ because X is well-filtered. It follows that $U \in \mathcal{N}(K_{i_0}) \subseteq \mathcal{F}$. Hence, X is Ψ_{wf} -well-filtered.

(2) \Rightarrow (1). Let $\{K_i : i \in I\}$ be a filtered family of $\mathcal{Q}(X)$ and $U \in \mathcal{O}(X)$ such that $\bigcap_{i \in I} K_i \subseteq U$. Then, $\mathcal{F} = \bigcup_{i \in I} \mathcal{N}(K_i) \in \Psi_{\text{wf}}(X)$. Note that $\bigcap \mathcal{F} = \bigcap_{i \in I} K_i \subseteq U$, and since X is Ψ_{wf} -well-filtered, it follows that $U \in \mathcal{F}$. By the definition of \mathcal{F} , there exists $i_0 \in I$ such that $U \in \mathcal{N}(K_{i_0})$, that is, $K_{i_0} \subseteq U$. Hence, X is well-filtered. □

Theorem 3.12. *Let X be a T_0 space. The following conditions are equivalent:*

- (1) X is a d -space;
- (2) X is Ψ_d -well-filtered;
- (3) $\forall \mathcal{F} \in \Psi_d(X)$, $\mathcal{F} = \mathcal{N}(\bigcap \mathcal{F})$;
- (4) $\Psi_d(X) = \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}$.

Proof. That (2) \Leftrightarrow (3) \Leftrightarrow (4) follows immediately from Theorem 3.6.

(1) \Rightarrow (2). Let $\mathcal{F} \in \Psi_d(X)$ and $U \in \mathcal{O}(X)$. If $\mathcal{F} = \mathcal{N}(K)$ for some $K \in \mathcal{Q}(X)$, then it is clear that $K = \bigcap \mathcal{N}(K) \subseteq U$ implies $U \in \mathcal{N}(K)$. Now assume there is a directed subset D of X such that $\mathcal{F} = \bigcup_{x \in D} \mathcal{N}(\uparrow x)$ and $\bigcap \mathcal{F} \subseteq U$. Since X is a d -space, $\bigvee D$ exists. We have that $\bigcap \mathcal{F} = \bigcap_{d \in D} \uparrow d = \uparrow \bigvee D \subseteq U$, which implies $\bigvee D \in U$. Since every open set in a d -space is Scott open, $D \cap U \neq \emptyset$, and take $x_0 \in D \cap U$. It follows that $U \in \mathcal{N}(\uparrow x_0) \subseteq \mathcal{F}$. Hence, X is Ψ_d -well-filtered.

(2) \Rightarrow (1). Let D be a directed subset of X , and $\mathcal{F} = \bigcup_{x \in D} \mathcal{N}(\uparrow x)$. We claim that $\text{cl}(D) \cap \bigcap \mathcal{F} \neq \emptyset$. Otherwise, $\bigcap \mathcal{F} \subseteq X \setminus \text{cl}(D)$, and since $\mathcal{F} \in \Psi_d(X)$ and X is Ψ_d -well-filtered, $X \setminus \text{cl}(D) \in \mathcal{F} = \bigcup_{x \in D} \mathcal{N}(\uparrow x)$. Then, there exists $x_0 \in D$ such that $X \setminus \text{cl}(D) \in \mathcal{N}(\uparrow x_0)$, which follows that $x_0 \in X \setminus$

$\text{cl}(D)$, contradicting the fact that $x_0 \in D$. Hence, there is $y \in \text{cl}(D) \cap \bigcap \mathcal{F} \neq \emptyset$. Then, $y \in \bigcap \mathcal{F} = \bigcap_{x \in D} \uparrow x$, so y is an upper bound of D . It follows that $D \subseteq \downarrow y = \text{cl}(\{y\})$, and since $y \in \text{cl}(D)$, it follows that $\text{cl}(D) = \text{cl}(\{y\})$. Hence, by Proposition 2.8 X is a d -space. \square

Remark 3.13. Note that $\Psi_d(X) \subseteq \Psi_{\text{wf}}(X) \subseteq \Psi_{\text{sob}}(X)$ for each T_0 space X . Then by Theorems 3.9, 3.11 and 3.12, the following relations are clear:

$$\text{sober space} \implies \text{well-filtered space} \implies d\text{-space}.$$

4. A Characterization for Ψ -Well-Filtered Spaces

In this section, we will show that a Ψ -well-filtered space X is determined by a class of subsets of X , called Ψ -sets.

Definition 4.1. Let Ψ be an HM-system and X be a T_0 space. A nonempty subset A of X is called a Ψ -set (relative to \mathcal{F}), if there exists $\mathcal{F} \in \Psi(X)$ such that $\text{cl}(A) \in m(\mathcal{F})$.

Remark 4.2. Let Ψ be an HM-system, X be a T_0 space, and $A \subseteq X$.

- (1) It is clear that A is a Ψ -set if and only if $\text{cl}(A)$ is a Ψ -set.
- (2) Every Ψ -set is irreducible by Lemma 2.12.

Lemma 4.3. Let X be a T_0 space, $K \in \mathcal{Q}(X)$, and $A \subseteq X$. The following two conditions are equivalent:

- (1) $\text{cl}(A) \cap K \neq \emptyset$;
- (2) $\forall U \in \mathcal{N}(K), A \cap U \neq \emptyset$.

Proof. That (1) \implies (2) is trivial. Conversely, if $\text{cl}(A) \cap K = \emptyset$, then $X \setminus \text{cl}(A) \in \mathcal{N}(K)$, but $A \cap (X \setminus \text{cl}(A)) = \emptyset$, contradicting the assumption (2). This shows that (2) implies (1). \square

Lemma 4.4. Let X be a T_0 space and $K \in \mathcal{Q}(X)$. Then,

$$m(\mathcal{N}(K)) = \{\text{cl}(\{x\}) : x \in \min K\},$$

where $\min K$ is the set of all minimal elements of K in the specialization order of X .

Proof. First, it is well-known that $K = \uparrow \min K$ (see, e.g., Ern  2009, pp. 2068). Suppose $x \in \min K$. It is clear that $\text{cl}(\{x\}) \in \mathcal{M}(\mathcal{N}(K))$. Now assume C is a closed subset of $\text{cl}(\{x\})$ such that $C \in \mathcal{M}(\mathcal{N}(K))$. By Lemma 4.3, there is $a \in C \cap K \neq \emptyset$. Then, $a \in C \subseteq \text{cl}(\{x\})$, so $a \leq x$. Since x is minimal in K , we have that $a = x \in C$, so $\text{cl}(\{x\}) \subseteq C$. Thus, $C = \text{cl}(\{x\})$. All this shows that $\text{cl}(\{x\}) \in m(\mathcal{N}(K))$.

Now assume $A \in m(\mathcal{N}(K))$. By Lemma 4.3, there is $a \in A \cap K \neq \emptyset$. Since $a \in K = \uparrow \min K$, there exists $x \in \min K$ such that $x \leq a$, which follows that $x \in \text{cl}(\{a\}) \subseteq A$. Then, $\text{cl}(\{x\})$ is a closed subset of A such that $\text{cl}(\{x\}) \in \mathcal{M}(\mathcal{N}(K))$. By the minimality of A , we have that $A = \text{cl}(\{x\})$. This shows that $m(\mathcal{N}(K)) \subseteq \{\text{cl}(\{x\}) : x \in \min K\}$, completing the proof. \square

Using Lemma 4.4, we deduce that $m(\mathcal{N}(\uparrow x)) = \{\text{cl}(\{x\})\}$ for each point x of a T_0 space X , and since $\mathcal{N}(\uparrow x) \in \Psi(X)$ for each HM-system Ψ , we have the following result.

Proposition 4.5. Let Ψ be an HM-system and X be a T_0 space. Every singleton of X is a Ψ -set.

Theorem 4.6. *Let Ψ be an HM-system and X be a T_0 space. The following conditions are equivalent:*

- (1) X is Ψ -well-filtered;
- (2) for each Ψ -set $A \subseteq X$, there exists $x \in X$ such that $\text{cl}(A) = \text{cl}(\{x\})$;
- (3) for each Ψ -set $A \subseteq X$ and $U \in \mathcal{O}(X)$, $\bigcap \mathcal{M}(A) \subseteq U$ implies $U \in \mathcal{M}(A)$;
- (4) for each Ψ -set $A \subseteq X$ and $U \in \mathcal{O}(X)$, $\bigcap_{a \in A} \uparrow a \subseteq U$ implies $\uparrow a \subseteq U$ for some $a \in A$.

Proof. (1) \Rightarrow (2). Let A be a Ψ -set in X . Then, there exists $\mathcal{F} \in \Psi(X)$ such that $\text{cl}(A) \in m(\mathcal{F})$. Since X is Ψ -well-filtered, it follows that $(\bigcap \mathcal{F}) \cap \text{cl}(A) \neq \emptyset$. Take $x \in (\bigcap \mathcal{F}) \cap \text{cl}(A) \neq \emptyset$. Then, $\text{cl}(\{x\})$ is a subset of $\text{cl}(A)$ such that $\text{cl}(\{x\}) \in \mathfrak{M}(\mathcal{F})$. By the minimality of $\text{cl}(A)$, we deduce that $\text{cl}(A) = \text{cl}(\{x\})$.

(2) \Rightarrow (3). It is trivial since $\mathcal{M}(\text{cl}(\{x\})) = \mathcal{N}(\{x\})$ for each $x \in X$.

(3) \Rightarrow (1). Let $\mathcal{F} \in \Psi(X)$ and $O \in \mathcal{O}(X)$ such that $\bigcap \mathcal{F} \subseteq O$. We need to show that $O \in \mathcal{F}$, or equivalently, $U \subseteq O$ for some $U \in \mathcal{F}$, because \mathcal{F} is an upper set. On the contrary, we assume $U \not\subseteq O$, that is, $U \cap (X \setminus O) \neq \emptyset$ for each $U \in \mathcal{F}$. By Lemma 2.12, there exists a closed subset A of $X \setminus O$ such that $A \in m(\mathcal{F})$. Then, A is a Ψ -set in X and note that $\mathcal{F} \subseteq \mathcal{M}(A)$, so $\bigcap \mathcal{M}(A) \subseteq \bigcap \mathcal{F} \subseteq O$. From condition (3), it follows that $O \in \mathcal{M}(A)$, that is, $O \cap A \neq \emptyset$, contradicting the fact that $A \subseteq X \setminus O$. This implies that $O \in \mathcal{F}$. Therefore, X is Ψ -well-filtered.

(2) \Rightarrow (4). Suppose condition (2) is satisfied. Then, there exists $x \in X$ such that $\text{cl}(A) = \text{cl}(\{x\}) = \downarrow x$, which follows that

$$\uparrow x = \bigcap_{a \in \downarrow x} \uparrow a = \bigcap_{a \in \text{cl}(A)} \uparrow a \subseteq \bigcap_{a \in A} \uparrow a \subseteq U.$$

This shows that U is an open neighborhood of x , and since $x \in \text{cl}(A)$, there is $a_0 \in U \cap A \neq \emptyset$, so $\uparrow a_0 \subseteq U$. This gives (4).

(4) \Rightarrow (2). Suppose A is a Ψ -set in X . By Remark 4.2, $\text{cl}(A)$ is also a Ψ -set. Since $\uparrow a \not\subseteq X \setminus \text{cl}(A)$ for each $a \in \text{cl}(A)$, by condition (4) there exists $x \in \text{cl}(A) \cap \bigcap_{a \in \text{cl}(A)} \uparrow a \neq \emptyset$. Then, we can easily obtain that $\text{cl}(A) = \text{cl}(\{x\})$. \square

Lemma 4.7. *Let X and Y be two T_0 spaces and A be an irreducible subset of X .*

- (1) $\mathcal{M}(A) \in \text{OF}(X)$ and $\text{cl}(A) \in m(\mathcal{M}(A))$.
- (2) For each $B \in m(\mathcal{M}(A))$, $\text{cl}(A) = B$. Hence, $\mathcal{M}(B) = \mathcal{M}(A)$.
- (3) If $f : X \rightarrow Y$ is a continuous mapping, then $f_*(\mathcal{M}(A)) = \mathcal{M}(f(A))$, where

$$f_*(\mathcal{M}(A)) = \{V \in \mathcal{O}(Y) : f^{-1}(V) \in \mathcal{M}(A)\}.$$

Hence, $\text{cl}_Y(f(A)) \in m(f_*(\mathcal{M}(A)))$.

Proof. (1) It is trivial that $\mathcal{M}(A) \in \text{OF}(X)$ and $\text{cl}(A) \in \mathfrak{M}(\mathcal{M}(A))$. To show $\text{cl}(A) \in m(\mathcal{M}(A))$, assume C is a closed subset of $\text{cl}(A)$ such that $C \in \mathfrak{M}(\mathcal{M}(A))$. We need to show that $A \subseteq C$. Otherwise, $A \cap (X \setminus C) \neq \emptyset$. Then, $X \setminus C \in \mathcal{M}(A)$. Since $C \in \mathfrak{M}(\mathcal{M}(A))$, it follows that $C \cap (X \setminus C) \neq \emptyset$, a contradiction. Thus, $C = \text{cl}(A)$. This shows that $\text{cl}(A) \in m(\mathcal{M}(A))$.

(2) Suppose $B \in m(\mathcal{M}(A))$. Let $x \in A$. For each open neighborhood U of x , $U \in \mathcal{M}(A)$, hence $B \cap U \neq \emptyset$. This implies $x \in \text{cl}(B) = B$. We then deduce that $A \subseteq B$, so $\text{cl}(A) \subseteq B$. From the minimality of B , it follows that $\text{cl}(A) = B$.

(3) For each $V \in \mathcal{O}(Y)$, we have that $V \in f_*(\mathcal{M}(A))$ iff $f^{-1}(V) \cap A \neq \emptyset$ iff $V \cap f(A) \neq \emptyset$ iff $V \in \mathcal{M}(f(A))$, which implies that $f_*(\mathcal{M}(A)) = \mathcal{M}(f(A))$. Since f is continuous, it follows that $f(A)$ is an irreducible set in Y . Hence, by (1), we have $\text{cl}_Y(f(A)) \in m(\mathcal{M}(f(A))) = m(f_*(\mathcal{M}(A)))$. \square

Proposition 4.8. *Let X be a T_0 space.*

- (1) The Ψ_{sob} -sets are exactly the irreducible sets in X .

- (2) The Ψ_{wf} -sets are exactly the KF-sets in X .
 (3) The closed Ψ_{d} -sets are exactly the closure of directed sets in X .

Proof. (1) Suppose A is an irreducible subset of X . By Lemma 4.7, $\mathcal{M}(A) \in \Psi_{\text{sob}}(X) = \text{OF}(X)$ and $\text{cl}(A) \in m(\mathcal{M}(A))$, so A is a Ψ_{sob} -set in X . The converse is trivial by Remark 4.2.

(2) Suppose $\mathcal{F} \in \Psi_{\text{wf}}(X)$. Then, there exists a filtered family $\{K_i : i \in I\} \subseteq \mathcal{Q}(X)$ such that $\mathcal{F} = \bigcup_{i \in I} \mathcal{N}(K_i)$. By Lemma 4.3, for each subset $A \subseteq X$, $\text{cl}(A) \in \mathfrak{M}(\mathcal{F})$ iff $\text{cl}(A) \cap K_i \neq \emptyset$ for each $i \in I$. Then, we deduce that the Ψ_{wf} -sets are exactly the KF-sets.

(3) Suppose A is a closed Ψ_{d} -set in X . Then, there exists a directed subset D of X such that $A \in m(\mathcal{M}(D))$. By Lemma 4.7, we have $\text{cl}(D) = A$. Now suppose E is a directed subset of X . Then, by Lemma 4.7, $\text{cl}(E) \in m(\mathcal{M}(E))$. This means that $\text{cl}(E)$ is a closed Ψ_{d} -set in X . \square

5. Relations between Ψ -Well-Filtered Spaces and H -Sober Spaces

In the paper Xu (2021), Xu introduces the notions of R -subset system and H -sober space, which provides a uniform approach to d -spaces, well-filtered spaces, and sober spaces. In this section, we study the relations between H -sober spaces and Ψ -well-filtered spaces.

Definition 5.1 (Xu 2021). A covariant functor $H : \mathbf{Top}_0 \longrightarrow \mathbf{Set}$ is called an R -subset system on \mathbf{Top}_0 if it satisfies the following two conditions:

- (H1) for each T_0 space X , $\{\{x\} : x \in X\} \subseteq H(X) \subseteq \text{Irr}(X)$;
 (H2) For each continuous mapping $f : X \longrightarrow Y$ in \mathbf{Top}_0 and each $A \in H(X)$, $H(f)(A) = f(A) \in H(Y)$.

For an R -subset system H and a T_0 space X , we call $A \subseteq X$ an H -set if $A \in H(X)$.

Definition 5.2 (Xu 2021). Let H be an R -subset system. A T_0 space X is called H -sober if for each $A \in H(X)$, there is a (unique) point $x \in X$ such that $\text{cl}(A) = \text{cl}(\{x\})$.

Next, we study the relationship between Ψ -well-filtered spaces and H -sober spaces, the following concept is needed.

Definition 5.3. An HM-system Ψ is called complete if for each continuous mapping $f : X \longrightarrow Y$ between T_0 spaces X and Y , and each Ψ -set A in X , $f(A)$ is a Ψ -set in Y .

Lemma 5.4. An HM-system Ψ is complete if and only if for each continuous mapping $f : X \longrightarrow Y$ between T_0 spaces X and Y , and each closed Ψ -set A in X , $f(A)$ is a Ψ -set in Y .

Proof. It suffices to prove the sufficiency. Suppose A is a Ψ -set in X . By assumption, $f(\text{cl}_X(A))$ is a Ψ -set in Y . By Remark 4.2, $\text{cl}_Y(f(\text{cl}_X(A))) = \text{cl}_Y(f(A))$ is a Ψ -set, so is $f(A)$, completing the proof. \square

Proposition 5.5. The HM-systems Ψ_{sob} , Ψ_{wf} , and Ψ_{d} are all complete.

Proof. By Proposition 4.8 and Lemma 2.6, Ψ_{wf} is complete. The completeness of Ψ_{sob} and Ψ_{d} is trivial (see Gierz et al. 2003) by Proposition 4.8. \square

Lemma 5.6. Let X and Y be two T_0 spaces, K be a compact saturated subset of X , and $f : X \longrightarrow Y$ be a continuous mapping.

- (1) $f_*(\mathcal{N}(K)) = \mathcal{N}(f(K))$.
 (2) If $A \in m(\mathcal{N}(K))$, then $\text{cl}_Y(f(A)) \in m(\mathcal{N}(\uparrow f(K \cap A)))$.

Proof. (1) It is trivial.

(2) First, by Lemma 4.4 there exists $x \in \min K$ such that $A = \text{cl}_X(\{x\})$. It follows that $A \cap K = \{x\}$, so $\uparrow f(K \cap A) = \uparrow f(x)$. In addition, using Lemma 4.3, one can easily obtain that $\text{cl}_Y(f(A)) \in \mathfrak{M}(\mathcal{N}(\uparrow f(x)))$. Suppose C is a closed subset of $\text{cl}_Y(f(A))$ such that $C \in \mathfrak{M}(\mathcal{N}(\uparrow f(x)))$. By Lemma 4.3, $C \cap \uparrow f(x) \neq \emptyset$, and since C is a lower set, we have that $f(x) \in C$, so $A = \text{cl}_X(\{x\}) \subseteq f^{-1}(C)$. It follows that $\text{cl}_Y(f(A)) \subseteq C$. Thus, $C = \text{cl}_Y(f(A))$. Therefore, $\text{cl}_Y(f(A)) \in m(\mathcal{N}(\uparrow f(K \cap A)))$. \square

Theorem 5.7. Let $H : \mathbf{Top}_0 \longrightarrow \mathbf{Set}$ be an R -subset system. Define $\Psi_H : \mathbf{Top}_0 \longrightarrow \mathbf{Set}$ by

$$\Psi_H(X) = \{\mathcal{M}(A) : A \in H(X)\} \cup \{\mathcal{N}(K) : K \in \mathcal{Q}(X)\}.$$

for each T_0 space X , and for each continuous mapping $f : X \longrightarrow Y$ between T_0 spaces X and Y , we define $\Psi_H(f) : \Psi_H(X) \longrightarrow \Psi_H(Y)$ by

$$\Psi_H(f)(\mathcal{F}) = f_*(\mathcal{F}) = \{V \in \mathcal{O}(Y) : f^{-1}(V) \in \mathcal{F}\}$$

for each $\mathcal{F} \in \Psi_H(X)$.

- (1) Ψ_H is a complete HM-system.
- (2) A T_0 space X is H -sober if and only if it is Ψ_H -well-filtered.

Proof. (1) We first prove that Ψ_H is an HM-system.

Note that each member of $H(X)$ is irreducible in X , so it is clear that Ψ_H satisfies (H1). To show (HM2), let $f : X \longrightarrow Y$ be a continuous mapping between T_0 spaces X and Y , and $A \in H(X)$, that is, $\mathcal{M}(A) \in \Psi_H(X)$. By (H2), $f(A) \in H(Y)$, and it follows from Lemma 4.7 that $f_*(\mathcal{M}(A)) = \mathcal{M}(f(A)) \in \Psi_H(Y)$, and since $f_*(\mathcal{N}(K)) = \mathcal{N}(f(K)) \in \Psi_H(Y)$, (H2) holds. In addition, it is trivial to check that Ψ_H is a covariant functor. Hence, Ψ_H is an HM-system.

Now we prove that Ψ_H is complete. Suppose A is a closed Ψ_H -set in a T_0 space X and $f : X \longrightarrow Y$ is a continuous mapping to a T_0 space Y . We need to prove that $f(A)$ is a Ψ_H -set in Y . We consider the following cases:

(c1) there exists $K \in \mathcal{Q}(X)$ such that $A \in m(\mathcal{N}(K))$. Note that the intersection $K \cap A$ is compact and since f is continuous, $f(K \cap A)$ is compact in Y , so $\uparrow f(K \cap A) \in \mathcal{Q}(Y)$. By Lemma 5.6, $\text{cl}_Y(f(A)) \in m(\mathcal{N}(\uparrow f(K \cap A)))$. This shows that $f(A)$ is a Ψ_H -set in Y .

(c2) there exists a $B \in H(X)$ such that $A \in m(\mathcal{M}(B))$. By Lemma 4.7, $A = \text{cl}_X(B)$ and $\text{cl}_Y(f(A)) \in m(\mathcal{M}(f(A)))$. Since $f(A) \in H(Y)$ by (H2), it follows that $\mathcal{M}(f(A)) = \mathcal{M}(f(\text{cl}_X(B))) = \mathcal{M}(f(B)) \in \Psi_H(Y)$. Thus $f(A)$ is a Ψ_H -set in Y .

By Lemma 5.4, Ψ is a complete HM-system.

(2) (\Rightarrow). Assume X is H -sober. Let A be a Ψ_H -set in X . There are two cases:

(c1) $A \in m(\mathcal{M}(B))$ for some $B \in H(X)$. By Lemma 4.7, $\text{cl}(B) = \text{cl}(A)$, and since X is H -sober, $\text{cl}(B) = \text{cl}(A) = \text{cl}(\{x\})$ for some $x \in X$.

(c2) $A \in m(\mathcal{N}(K))$ for some $K \in \mathcal{Q}(X)$. By Lemma 4.4, $\text{cl}(A) = \text{cl}(\{x\})$ for some $x \in \min K$.

Then by Theorem 4.6, X is Ψ_H -well-filtered.

(\Leftarrow). Assume X is a Ψ_H -well-filtered space. Let $A \in H(X)$. Then by Lemma 4.7, $\text{cl}(A) \in m(\mathcal{M}(A)) \in \Psi_H(X)$, so A is a Ψ_H -set in X . Since X is Ψ_H -well-filtered, by Theorem 4.6 $\text{cl}(A) = \text{cl}(\{x\})$ for some $x \in X$. Therefore, X is H -sober. \square

Theorem 5.8. Let $\Psi : \mathbf{Top}_0 \longrightarrow \mathbf{Set}$ be a complete HM-system. Define $H_\Psi : \mathbf{Top}_0 \longrightarrow \mathbf{Set}$ by

$$H_\Psi(X) = \{A \subseteq X : A \text{ is a } \Psi\text{-set in } X\}$$

for each T_0 space X , and for each continuous mapping $f : X \longrightarrow Y$ between T_0 spaces X and Y , define $H_\Psi(f) : H_\Psi(X) \longrightarrow H_\Psi(Y)$ by

$$H_\Psi(f)(A) = f(A)$$

for each $A \in H_\Psi(X)$.

- (1) H_Ψ is an R -system.
- (2) A T_0 space X is Ψ -well-filtered if and only if it is H_Ψ -sober.

Proof. (1) We first prove that H_Ψ satisfies (H1) and (H2). For (H1), it follows from Proposition 4.5 that $\{\{x\} : x \in X\} \subseteq H_\Psi(X)$. By Lemma 2.12, every Ψ -set is irreducible; hence, $H_\Psi(X) \subseteq \text{Irr}(X)$. Thus, (H1) holds. Condition (H2) holds immediately since Ψ is complete. It is trivial to check that H is a covariant functor. Therefore, H_Ψ is an R -system.

- (2) It is straightforward by Theorem 4.6. □

For an HM-system Ψ , we use Ψ -WF to denote the category of all Ψ -well-filtered spaces with continuous mappings. In Xu (2021), Xu proved that the category of all H -sober spaces with continuous mappings is reflective in the category \mathbf{Top}_0 of T_0 spaces. Then by Theorem 5.8, we have the following corollary.

Corollary 5.9. *For a complete HM-system Ψ , the category Ψ -WF is a reflective subcategory of \mathbf{Top}_0 .*

6. The Smyth Power Space of a Ψ -Well-Filtered Space

For a topological space X , the upper Vietoris topology on $\mathcal{Q}^*(X) = \mathcal{Q}(X) \setminus \{\emptyset\}$ is generated by the following family (as a base)

$$\square U = \{K \in \mathcal{Q}^*(X) : K \subseteq U\},$$

where U ranges over the open subsets of X . The resulting space, denoted by $P_s(X)$, is called the Smyth power space or the upper space.

Remark 6.1 (Goubault-Larrecq 2003; Jia and Jung 2016; Schalk 1993). Let X be a T_0 space.

- (1) The specialization order of $P_s(X)$ is \supseteq . Hence, for each $\mathcal{A} \subseteq \mathcal{Q}^*(X)$,

$$\uparrow_{P_s(X)} \mathcal{A} = \{K \in \mathcal{Q}^*(X) : K \subseteq G \text{ for some } G \in \mathcal{A}\}$$

in the specialization order of $P_s(X)$.

- (2) Define $\xi : X \rightarrow P_s(X)$, $x \mapsto \uparrow x$. Then $\xi^{-1}(\square U) = U$ for each $U \in \mathcal{O}(X)$, and hence ξ is continuous.
- (3) If \mathcal{K} is compact in $P_s(X)$, then $\bigcup \mathcal{K}$ is compact in X .

Definition 6.2 (Xu 2021). Let X be a T_0 space. An R -subset system H is said to satisfy property Q for X if for each $\mathcal{A} \in H(P_s(X))$ and each $C \in \mathfrak{M}(\mathcal{A})$, there is an H -subset F of C such that $\text{cl}(F) \in \mathfrak{M}(\mathcal{A})$.

Lemma 6.3. Let X be a T_0 space, $\mathcal{A} \subseteq \mathcal{Q}(X)$. Then, $\mathfrak{M}(\mathcal{A}) = \mathfrak{M}(\text{cl}_{P_s(X)}(\mathcal{A}))$. Hence, $m(\mathcal{A}) = m(\text{cl}_{P_s(X)}(\mathcal{A}))$.

Proof. We only need to prove that $\mathfrak{M}(\mathcal{A}) \subseteq \mathfrak{M}(\text{cl}_{P_s(X)}(\mathcal{A}))$. Assume on the contrary there exists $C \in \mathfrak{M}(\mathcal{A})$ such that $C \notin \mathfrak{M}(\text{cl}_{P_s(X)}(\mathcal{A}))$. Then, there exists $K \in \text{cl}_{P_s(X)}(\mathcal{A})$ such that $C \cap K = \emptyset$. This implies that $\square(X \setminus C)$ is an open neighborhood of K in $P_s(X)$, and since $K \in \text{cl}_{P_s(X)}(\mathcal{A})$, there is $G \in \mathcal{A} \cap \square(X \setminus C) \neq \emptyset$. Thus, G is a member of \mathcal{A} such that $G \cap C = \emptyset$, contradicting the fact that $C \in \mathfrak{M}(\mathcal{A})$. Hence, $\mathfrak{M}(\mathcal{A}) \subseteq \mathfrak{M}(\text{cl}_{P_s(X)}(\mathcal{A}))$. □

The following theorem strengthens a result in Xu (2021).

Theorem 6.4. *Let H be an R -subset system and X be a T_0 space. The following two conditions are equivalent:*

- (1) $P_s(X)$ is H -sober;
- (2) X is H -sober and H has property Q for X .

Proof. By Xu (2021, Theorem 5.12), we only need to check that condition (1) implies that H has property Q for X . Suppose $\mathcal{A} \in H(P_s(X))$ and $C \in \mathfrak{M}(\mathcal{A})$. Since $P_s(X)$ is H -sober, there exists $K \in \mathcal{Q}^*(X)$ such that $\text{cl}_{P_s(X)}(\mathcal{A}) = \text{cl}_{P_s(X)}(\{K\})$. By Lemma 6.3, $C \in \mathfrak{M}(\mathcal{A}) = \mathfrak{M}(\text{cl}_{P_s(X)}(\mathcal{A})) = \mathfrak{M}(\text{cl}_{P_s(X)}(\{K\})) = \mathfrak{M}(\{K\})$, which follows that $C \cap K \neq \emptyset$. Take $x \in C \cap K$. Then, $\{x\}$ is an H -set, and it is clear that $\text{cl}(\{x\}) \in \mathfrak{M}(\{K\}) = \mathfrak{M}(\mathcal{A})$. \square

Definition 6.5. *Let X be a T_0 space. An HM-system Ψ is said to satisfy property Q for X if for each Ψ -set \mathcal{A} in $P_s(X)$ and each $C \in \mathfrak{M}(\mathcal{A})$, there is a Ψ -subset F of C such that $\text{cl}(F) \in \mathfrak{M}(\mathcal{A})$.*

From Theorems 5.7 and 5.8, it turns out that there is a one to one correspondence between H -sober spaces and Ψ -well-filtered spaces when Ψ is complete. In other words, the class of Ψ -well-filtered spaces are more general than H -sober spaces. As a generalized result of Theorem 6.4, we have the following result.

Theorem 6.6. *Let Ψ be an HM-system and X be a T_0 space. Then, the following conditions are equivalent:*

- (1) $P_s(X)$ is Ψ -well-filtered;
- (2) X is Ψ -well-filtered and Ψ has property Q for X ;
- (3) for each Ψ -set \mathcal{A} in $P_s(X)$ and each $U \in \mathcal{O}(X)$, $\bigcap \mathcal{A} \subseteq U$ implies $K \subseteq U$ for some $K \in \mathcal{A}$.

Proof. (1) \Rightarrow (2). We first prove that X is Ψ -well-filtered. Let $\mathcal{F} \in \Psi(X)$ and $U \in \mathcal{O}(X)$ with $\bigcap \mathcal{F} \subseteq U$. Consider the continuous mapping $\xi : X \rightarrow P_s(X)$, $x \mapsto \uparrow x$. Then, $\xi_*(\mathcal{F}) = \{\mathcal{U} \in \mathcal{O}(P_s(X)) : \xi^{-1}(\mathcal{U}) \in \mathcal{F}\} \in \Psi(P_s(X))$ by the definition of Ψ . Let

$$\mathfrak{F} = \xi_*(\mathcal{F}) \cap \{\square V : V \in \mathcal{O}(X)\}.$$

For each $V \in \mathcal{O}(X)$, observe that $\square V \in \mathfrak{F}$ iff $\xi^{-1}(\square V) = V \in \mathcal{F}$, which implies that $\mathfrak{F} = \{\square V : V \in \mathcal{F}\}$. If $K \in \bigcap \mathfrak{F}$, then $K \in \square V$ (i.e., $K \subseteq V$) for all $V \in \mathcal{F}$, so $K \subseteq \bigcap \mathcal{F} \subseteq U$, which implies $K \in \square U$. Thus, $\bigcap \mathfrak{F} \subseteq \square U$. Since $P_s(X)$ is Ψ -well-filtered and $\bigcap \xi_*(\mathcal{F}) \subseteq \bigcap \mathfrak{F} \subseteq \square U$, we have that $\square U \in \xi_*(\mathcal{F})$, so $\xi^{-1}(\square U) = U \in \mathcal{F}$. Hence, X is Ψ -well-filtered. Using a similar method of Theorem 6.4, one can prove that Ψ has property Q for X .

(2) \Rightarrow (3). Assume on the contrary that $K \cap (X \setminus U) \neq \emptyset$ for each $K \in \mathcal{A}$. It follows that $X \setminus U \in \mathfrak{M}(\mathcal{A})$. Since Ψ satisfies property Q for X , there exists a Ψ -set $F \subseteq X \setminus U$ such that $\text{cl}(F) \in \mathfrak{M}(\mathcal{A})$. Since X is Ψ -well-filtered, by Theorem 4.6 there exists $x \in X$ such that $\text{cl}(F) = \text{cl}(\{x\})$. For each $K \in \mathcal{A}$, since $K = \uparrow K$ and $\text{cl}(F) \cap K = \text{cl}(\{x\}) \cap K \neq \emptyset$, we deduce that $x \in K$. Thus, $x \in \bigcap \mathcal{A} \subseteq U$, contradicting the fact that $x \in \text{cl}(F) \subseteq X \setminus U$. This shows that $K \subseteq U$ for some $K \in \mathcal{A}$, which gives (3).

(3) \Rightarrow (1). Suppose \mathcal{A} is a Ψ -set in $P_s(X)$ and \mathcal{U} is an open set in $P_s(X)$ such that $\bigcap_{K \in \mathcal{A}} \uparrow_{P_s(X)} K \subseteq \mathcal{U}$. Then, there exists a family $\{U_i : i \in I\} \subseteq \mathcal{O}(X)$ such that $\mathcal{U} = \bigcup_{i \in I} \square U_i$. Using a similar proof of Lemma 3.5, one can obtain that $K_0 = \bigcap \mathcal{A} \in \mathcal{Q}^*(X)$. Note that $\uparrow_{P_s(X)} K = \{G \in \mathcal{Q}^*(X) : G \subseteq K\}$ for each $K \in \mathcal{A}$, so $K_0 \in \bigcap_{K \in \mathcal{A}} \uparrow_{P_s(X)} K \subseteq \bigcup_{i \in I} \square U_i$. Then, there exists $i_0 \in I$ such that $K_0 = \bigcap \mathcal{A} \subseteq \square U_{i_0}$, so $\bigcap \mathcal{A} \subseteq U_{i_0}$. By condition (3), there exists $K_1 \in \mathcal{A}$ such that $K_1 \subseteq U_{i_0}$. This implies that $\uparrow_{P_s(X)} K_1 \subseteq \square U_{i_0} \subseteq \mathcal{U}$. By Theorem 4.6, we deduce that $P_s(X)$ is Ψ -well-filtered. \square

Definition 6.7. Let X be a T_0 space. An HM-system Ψ is said to satisfy property T for X if for any Ψ -set \mathcal{A} in $P_s(X)$, $\bigcup_{K \in \mathcal{A}} \mathcal{N}(K) \in \Psi(X)$.

Remark 6.8. Using Lemma 4.3, one can easily deduce that $\mathfrak{M}(\mathcal{A}) = \mathfrak{M}(\mathcal{F}_{\mathcal{A}})$ in Definition 6.7, where $\mathcal{F}_{\mathcal{A}} = \bigcup_{K \in \mathcal{A}} \mathcal{N}(K)$. Then by Lemma 2.12, if an HM-system Ψ satisfies property T, then it must satisfy Q.

Lemma 6.9. Both Ψ_{sob} and Ψ_{wf} satisfy property T for each T_0 space X .

Proof. Let X be a T_0 space. It is trivial that Ψ_{sob} satisfies property T for X . Next, we verify that Ψ_{wf} satisfies property T for X . To show this, let $\mathfrak{F} \in \Psi_{\text{wf}}(P_s(X))$ and $\mathcal{A} \in m(\mathfrak{F})$. Then, there exists a filtered family $\{\mathcal{K}_i : i \in I\}$ of compact saturated subsets of $P_s(X)$ such that $\mathfrak{F} = \{\mathcal{U} \in \mathcal{O}(P_s(X)) : \exists i \in I, \mathcal{K}_i \subseteq \mathcal{U}\}$. It suffices to prove $\bigcup_{K \in \mathcal{A}} \mathcal{N}(K) \in \Psi_{\text{wf}}(X)$.

Claim: $\bigcup_{K \in \mathcal{A}} \mathcal{N}(K) = \bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i))$.

On the one hand, suppose $U \in \bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i))$. Then, there exists $i_0 \in I$ such that $\bigcup (\mathcal{A} \cap \mathcal{K}_{i_0}) \subseteq U$. Choose one $K_{i_0} \in \mathcal{A} \cap \mathcal{K}_{i_0} \neq \emptyset$. Then, $K_{i_0} \subseteq \bigcup (\mathcal{A} \cap \mathcal{K}_{i_0}) \subseteq U$, which implies that $U \in \mathcal{N}(K_{i_0}) \subseteq \bigcup_{K \in \mathcal{A}} \mathcal{N}(K)$. This shows that $\bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i)) \subseteq \bigcup_{K \in \mathcal{A}} \mathcal{N}(K)$. On the other hand, suppose $U \notin \bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i))$. For each $i \in I$, $\bigcup (\mathcal{A} \cap \mathcal{K}_i) \not\subseteq U$, so there exists $G_i \in \mathcal{A} \cap \mathcal{K}_i$ such that $G_i \not\subseteq U$; that is, $G_i \notin \square U$. Then, $G_i \in (P_s(X) \setminus \square U) \cap \mathcal{A} \cap \mathcal{K}_i \neq \emptyset$ for all $i \in I$. Thus, $(P_s(X) \setminus \square U) \cap \mathcal{A}$ is a closed subset of \mathcal{A} such that $(P_s(X) \setminus \square U) \cap \mathcal{A} \in \mathfrak{M}(\mathfrak{F})$. Since $\mathcal{A} \in m(\mathfrak{F})$, $\mathcal{A} \cap (P_s(X) \setminus \square U) = \mathcal{A}$, that is, $\mathcal{A} \subseteq P_s(X) \setminus \square U$. It follows that $K \not\subseteq U$ for all $K \in \mathcal{A}$; hence, $U \notin \bigcup_{K \in \mathcal{A}} \mathcal{N}(K)$. Therefore, $\bigcup_{K \in \mathcal{A}} \mathcal{N}(K) \subseteq \bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i))$.

Recall that the intersection of a closed set and a compact set is always compact, so $\mathcal{A} \cap \mathcal{K}_i$ is compact in $P_s(X)$ for each $i \in I$. Since $\{\mathcal{K}_i : i \in I\}$ is a filtered family, by Remark 6.1 $\{\uparrow \bigcup (\mathcal{A} \cap \mathcal{K}_i) : i \in I\}$ is a filtered family of $\mathcal{Q}^*(X)$. Note that for each open subset U of X , $\bigcup (\mathcal{A} \cap \mathcal{K}_i) \subseteq U$ if and only if $\uparrow \bigcup (\mathcal{A} \cap \mathcal{K}_i) \subseteq U$, we have that $\bigcup_{K \in \mathcal{A}} \mathcal{N}(K) = \bigcup_{i \in I} \mathcal{N}(\bigcup (\mathcal{A} \cap \mathcal{K}_i)) = \bigcup_{i \in I} \mathcal{N}(\uparrow \bigcup \mathcal{A} \cap \mathcal{K}_i) \in \Psi_{\text{wf}}(X)$. Therefore, Ψ_{wf} satisfies property T. \square

As a direct consequence of Theorem 6.6, Remark 6.8 and Lemma 6.9, we have the following corollaries.

Corollary 6.10 (Heckmann and Keimel 2013). Let X be a T_0 space. The following conditions are equivalent:

- (1) X is sober;
- (2) $P_s(X)$ is sober.

Corollary 6.11 (Xu et al. 2021). Let X be a T_0 space. The following conditions are equivalent:

- (1) X is well-filtered;
- (2) $P_s(X)$ is well-filtered.

7. Conclusion

Motivated by the Hofmann-Mislove Theorem on sober spaces, we introduce the Ψ -well-filtered spaces, where $\Psi : \mathbf{Top}_0 \rightarrow \mathbf{Set}$ is a covariant functor that assigns each T_0 space X a family of open filters of $\mathcal{O}(X)$. The classes of d -spaces, well-filtered spaces, and sober spaces are all special types of Ψ -well-filtered spaces. The \mathcal{U}_s -admitting spaces (Heckmann 1991), and the recently introduced open well-filtered spaces (Shen et al. 2020) and the ω -well-filtered spaces (Xu et al. 2020b) can also be viewed as special types of Ψ -well-filtered spaces. The results in this paper reveal some features of such spaces similar to that of sober space as shown by the Hofmann-Mislove Theorem. We

hope that based on this general notion, some new classes of interesting spaces can be identified in the future.

Acknowledgements. The authors would like to thank the editor and referee for the numerous and very helpful suggestions that have improved this paper substantially.

References

- Engelking, R. (1989). *General Topology*, Sigma Series in Pure Mathematics, vol. 6, Berlin, Heldermann Verlag.
- Erné, M. (2009). Infinite distributive laws versus local connectedness and compactness properties. *Topology and its Applications* **156** 2054–2069.
- Eršov, Yu. L. (1999). On d -spaces. *Theoretical Computer Science* **224** 59–72.
- Gierz, G., Hofmann, K., Keimel, K., Lawson, J., Mislove, M. and Scott, D. (2003). *Continuous lattices and Domains*, Encyclopedia of Mathematics and Its Applications, vol. 93, Cambridge, Cambridge University Press.
- Goubault-Larrecq, J. (2013). *Non-Hausdorff Topology and Domain Theory*, Cambridge, Cambridge University Press.
- Heckmann, R. (1991). An upper power domain construction in terms of strongly compact sets. In: *International Conference on Mathematical Foundations of Programming Semantics*, Berlin, Heidelberg, Springer, 272–293.
- Heckmann, R. and Keimel, K. (2013). Quasicontinuous domains and the Smyth powerdomain. *Electronic Notes in Theoretical Computer Science* **298** 215–232.
- Hoffmann, R. E. (1981). Projective sober spaces. In: Banaschewski, B. and Hoffmann, R. E. (eds.) *Continuous Lattices*, Lecture Notes in Mathematics, vol. 871, Berlin, Heidelberg, Springer, **105** 125–158.
- Hoffmann, R. (1979). On the sobrification remainder $X^s - X$. *Pacific Journal of Mathematics* **83** 145–156.
- Jia, X. and Jung, A. (2016). A note on coherence of dcpos. *Topology and its Applications* **209** 235–238.
- Keimel, K. and Lawson, J. (2009). D -topology and d -completion. *Annals of Pure and Applied Logic* **159** 292–306.
- Li, Q., Yuan, Z. and Zhao, D. (2021). A unified approach to some non-Hausdorff topological properties. *Mathematical Structures in Computer Science* **30** (9) 997–1010.
- Liu, B., Li, Q. and Wu, G. (2020). Well-filterifications of topological spaces. *Topology and its Applications* **279** 107245.
- Schalk, A. (1993). *Algebras for Generalized Power Constructions*. Phd thesis, Technische Hochschule Darmstadt.
- Shen, C., Xi, X., Xu, X. and Zhao, D. (2019). On well-filtered reflections of T_0 spaces. *Topology and its Applications* **267** 106869.
- Shen, C., Xi, X., Xu, X. and Zhao, D. (2020). On open well-filtered spaces. *Logical Methods in Computer Science* **16** (4) 4–18.
- Wu, G., Xi, X., Xu, X. and Zhao, D. (2020). Existence of well-filtered reflections of T_0 topological spaces. *Topology and its Applications* **270** 107044.
- Wyler, O. (1981). Dedekind complete posets and Scott topologies. In: *Continuous Lattices*, Berlin, Heidelberg, Springer, 384–389.
- Xu, X. (2021). On H -sober spaces and H -sobrifications of T_0 spaces. *Topology and its Applications* **289** 107548.
- Xu, X., Shen, C., Xi, X. and Zhao, D. (2020a). On T_0 spaces determined by well-filtered spaces. *Topology and its Applications* **282** 107323.
- Xu, X., Shen, C., Xi, X. and Zhao, D. (2020b). First countability, ω -well-filtered spaces and reflections. *Topology and its Applications* **279** 107255.
- Xu, X., Xi, X. and Zhao, D. (2021). A complete Heyting algebra whose Scott space is non-sober. *Fundamenta Mathematicae* **252** 315–323.