

THE NON-REFLECTIVITY OF OPEN WELL-FILTERED SPACES VIA b -TOPOLOGY

CHONG SHEN*, XIAOYONG XI, AND DONGSHENG ZHAO

Communicated by Yasunao Hattori

ABSTRACT. Results have shown that Skula's b -topology plays a core role in studying sober spaces (especially in studying the sobrification). In this paper, following Skula's method of b -embedding a subcategory into the category **Top** of all topological spaces, we restrict our attention to T_0 spaces, and mainly prove that if \mathbf{K} is a reflective subcategory of the category **Top**₀ of all T_0 spaces such that \mathbf{K} contains a non- T_1 space, then (i) each \mathbf{K} -reflection is a b -dense embedding; (ii) \mathbf{K} is saturated-hereditary. As a corollary, we deduce that the category **Owf** of all open well-filtered spaces is not reflective in **Top**₀, which demonstrates a new approach to the reflectivity of subcategories of **Top**₀.

1. INTRODUCTION

The reflective subcategories of the category **Top** of all topological spaces have been extensively studied. In the paper [9], Kennison gave three types of (full) reflective subcategories of **Top**, called simple, identifying and embedding, and provided a characterization for each of these reflections. He then asked whether these three types include all the full reflective subcategories of **Top**. Skula [11] then introduced another type, called b -embedding type, not mentioned by Kennison in [9], and showed that if \mathbf{K} is a reflective subcategory of **Top** that contains a non- T_1 space, then \mathbf{K} is a subcategory of one of the above-mentioned types.

2020 *Mathematics Subject Classification.* 06B35; 06B30; 54A05.

Key words and phrases. Epimorphism, sober space, b -topology, strong topology, reflection.

This work was supported by the National Natural Science Foundation of China (No. 1210010153, 12071188), Jiangsu Provincial Department of Education (No. 21KJB110008), International Postdoctoral Exchange Fellowship Program (No. PC2022013).

*Corresponding author.

In domain theory, many important classes of topological spaces are generally T_0 but not T_1 . Sobriety is one of the most well studied topological properties. One of the heavily used structures in study sober spaces is the b -topology (also called the Skula topology, or the strong topology), which was introduced by Skula in [11]. Some major results on b -topology include:

- (i) for a sober space X , the subsets of X that are sober as a subspace are exactly the b -closed subsets [3, 8];
- (ii) for a sober space X and $A \subseteq X$, the sobrification of the subspace A is homeomorphic to its b -closure [3, 8];
- (iii) for a T_0 space X and a sober space Y , $f : X \rightarrow Y$ is a sobrification mapping if and only if f is an embedding and $f(X)$ is b -dense in Y [3, 5, 8];
- (iv) if $f : X \rightarrow Y$ is a continuous mapping between T_0 spaces, then f is an epimorphism in **Top**₀ (the category of all T_0 spaces) if and only if $f(X)$ is b -dense in Y [1].

In this paper, we shall focus on T_0 spaces as this entails no essential loss of generality from the viewpoint of domain theory. Let **K** be a reflective subcategory of **Top**₀ that contains a non- T_1 space. The main results are

- (1) each **K**-reflection is a b -dense embedding, hence **K** is an epireflection;
- (2) **K** is saturated-hereditary: for any $X \in \mathbf{K}$, every saturated subset A of X , as a subspace, is in **K**.

Johnstone's \mathbb{J} and the space \mathbb{N}_{cof} of the set of natural numbers with the cofinite topology are commonly used for giving counterexamples. Note that \mathbb{N}_{cof} is homeomorphic to the maximal points space $\text{Max}(\mathbb{J})$ relative to the Scott space $\Sigma\mathbb{J}$, and as an immediate corollary of the above result (2), we have the following result:

- (3) If $\Sigma\mathbb{J} \in \mathbf{K}$, then $\mathbb{N}_{\text{cof}} \in \mathbf{K}$.

The notion of open well-filtered spaces, proposed by Shen, Xi, Xu and Zhao [12] recently, forms a strictly larger class than that of the well-filtered spaces. One notable result on open well-filtered spaces is that every core-compact open well-filtered space is sober, which strengthens the result of Jia-Jung problem: whether core-compact well-filtered spaces are sober [6] (a positive answer is given by Lawson, Wu and Xi [10]). However, it is still unknown

- whether the category **Owf** of all open well-filtered spaces is reflective in **Top**₀.

To solve this problem, we prove that $\Sigma\mathbb{J} \in \mathbf{Owf}$ but $\mathbb{N}_{\text{cof}} \notin \mathbf{Owf}$, and then by (3), we deduce the following result:

(4) The category **Owf** is not reflective in **Top**₀.

2. PRELIMINARIES

Let P be a poset. For $A \subseteq P$, let $\downarrow A = \{x \in P : x \leq a \text{ for some } a \in A\}$ and $\uparrow A = \{x \in P : x \geq a \text{ for some } a \in A\}$. For $x \in P$, $\downarrow\{x\}$ and $\uparrow\{x\}$ will be abbreviated as $\downarrow x$ and $\uparrow x$, respectively.

For a T_0 space X , the specialization order \leq is defined as $x \leq y$ iff $x \in \text{cl}_X(\{y\})$, where cl_X is the closure operator of X . In the following, when we consider a T_0 space X as a poset, it is always equipped with the specialization order.

For a T_0 space X , we use $\mathcal{O}(X)$ to denote the topology of X . For any subset A of X , the *saturation* of A , denoted by $\text{Sat}_X(A)$, is defined to be

$$\text{Sat}_X(A) = \bigcap \{U \subseteq \mathcal{O}(X) : A \subseteq U\}.$$

A subset A of a T_0 space X is called *saturated* if $A = \text{Sat}_X(A)$. In the following, we use $\text{Sat}(A)$ and $\text{cl}(A)$ for $\text{Sat}_X(A)$ and $\text{cl}_X(A)$ respectively if no ambiguity occurs.

Remark 2.1 ([3, 4]). Let X be a T_0 space.

- (1) For any subset A of X , $\text{Sat}(A) = \uparrow A$ with respect to the specialization order.
- (2) For any $x \in X$, $\downarrow x = \text{cl}(\{x\})$, and $x \in \text{Sat}(A)$ if and only if $\downarrow x \cap A \neq \emptyset$.
- (3) For any open subset U of X , $U = \text{Sat}(U) = \uparrow U$, and dually for any closed subset F of X , $F = \downarrow F$.

Definition 2.2 ([3, 4]). Let X be a T_0 space.

- (1) A nonempty subset $A \subseteq X$ is called *irreducible* if for any closed sets F_1, F_2 of X , $A \subseteq F_1 \cup F_2$ implies $A \subseteq F_1$ or $A \subseteq F_2$.
- (2) The space X is called *sober*, if for any irreducible closed set F of X there is a unique point $x \in X$ such that $F = \downarrow x$.

An essential tool for studying sober spaces is the b -topology introduced by L. Skula [11] (see also [2]).

Definition 2.3 ([2, 11]). Let X be a T_0 space. The b -topology associated with X is the topology which has the family

$$\{U \cap \downarrow x : x \in U \in \mathcal{O}(X)\}$$

as a base, and the resulting space is denoted by bX . We call a subset B of X *b-dense* in X , if it is dense in bX .

Lemma 2.4. *Let X be a T_0 space, and Y be a b -dense subspace of X .*

- (1) For each $V \in \mathcal{O}(X)$, $V = \uparrow(V \cap Y)$.
 (2) For each $U \in \mathcal{O}(Y)$, $\uparrow U \in \mathcal{O}(X)$.

PROOF. (1) Since every open set is saturated, it follows that $\uparrow(V \cap Y) \subseteq V$. Let $x \in V$. Then $\downarrow_X x \cap V$ is b -open in X , and since Y is b -dense in X , it follows that $\downarrow_X x \cap V \cap Y \neq \emptyset$, which implies that $x \in \uparrow(V \cap Y)$. Hence, $V \subseteq \uparrow(V \cap Y)$.

(2) Since U is open in Y , there exists an open set V in X such that $U = V \cap Y$. By (1), we have that $V = \uparrow(V \cap Y) = \uparrow U$. Hence, $\uparrow U$ is open in X . \square

The following theorem is trivial by Lemma 2.4.

Theorem 2.5. Let X be a T_0 space, and Y be a b -dense subspace of X . Define $\varphi : \mathcal{O}(Y) \rightarrow \mathcal{O}(X)$ and $\phi : \mathcal{O}(X) \rightarrow \mathcal{O}(Y)$ by

$$\varphi(U) = \uparrow U \text{ and } \phi(V) = V \cap Y,$$

for any $U \in \mathcal{O}(Y)$ and $V \in \mathcal{O}(X)$. Then φ and ϕ are order-isomorphisms under the inclusion order, and they are inverse to each other.

In what follows, we denote by \mathbf{Top}_0 (resp., \mathbf{Top}_1 , \mathbf{Sob}) the category of all T_0 spaces (resp., T_1 spaces, sober spaces) with continuous mappings as morphisms, and when we say \mathbf{K} a subcategory of \mathbf{Top}_0 it is always assumed to be full and replete (i.e., \mathbf{K} is closed under the formation of homeomorphic objects).

Definition 2.6. A subcategory \mathbf{K} of \mathbf{Top}_0 is *reflective*, if for each $X \in \mathbf{Top}_0$, there exists $X^k \in \mathbf{K}$ (the \mathbf{K} -completion for X) and a continuous mapping $\mu_X : X \rightarrow X^k$ (the \mathbf{K} -reflection for X) satisfying the universal property: for any continuous mapping $f : X \rightarrow Z$ to any space $Z \in \mathbf{K}$, there exists a unique continuous mapping $g : X^k \rightarrow Z$ such that $g \circ \mu_X = f$:

$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow f & \downarrow g \\ & & Z \end{array}$$

Definition 2.7. Let $e : X \rightarrow Y$ be a mapping between topological spaces. We call e a b -dense embedding, if it is an embedding such that $e(X)$ is b -dense in Y .

The following result, due to Skula [11], will be used. Here, for the sake of completeness, we provide a proof by using Definition 2.3 directly.

Lemma 2.8 (compare to [11, I-2.5]). Let $X, Y \in \mathbf{Top}_0$ and $e : X \rightarrow Y$ be a b -dense embedding. Then for any continuous mapping $f : X \rightarrow Z$ to a T_0 space Z , there exists at most one continuous mapping $g : Y \rightarrow Z$ such that $g \circ e = f$.

PROOF. Suppose there exist two continuous mappings $g_1, g_2 : Y \longrightarrow Z$ such that $g_1 \circ e = g_2 \circ e = f$:

$$\begin{array}{ccc} X & \xrightarrow{e} & Y \\ & \searrow f & \downarrow g_1, g_2 \\ & & Z \end{array}$$

Let $y \in Y$. Suppose $V \in \mathcal{O}(Z)$ such that $g_1(y) \in V$. Then $y \in g_1^{-1}(V) \in \mathcal{O}(Y)$, which follows that $\downarrow_Y y \cap g_1^{-1}(V)$ is b -open in Y . Since $e(X)$ is b -dense in Y , $\downarrow_Y y \cap g_1^{-1}(V) \cap e(X) \neq \emptyset$, and from $g_1 \circ e = g_2 \circ e = f$, we deduce that $g_1^{-1}(V) \cap e(X) = g_2^{-1}(V) \cap e(X)$, and hence $\downarrow_Y y \cap g_2^{-1}(V) \cap e(X) \neq \emptyset$. It follows that $y \in \uparrow_Y g_2^{-1}(V) = g_2^{-1}(V)$, so $g_2(y) \in V$. All this shows that each open neighborhood of $g_1(y)$ contains $g_2(y)$, that is, $g_1(y) \in \text{cl}_Z(\{g_2(y)\})$. Dually, $g_2(y) \in \text{cl}_Z(\{g_1(y)\})$. Since Z is a T_0 space, we have that $g_1(y) = g_2(y)$. Therefore, $g_1 = g_2$. \square

Definition 2.9 ([3, 4]). Let X and Y be two topological spaces. X is called a *retract* of Y if there are two continuous mappings $s : X \longrightarrow Y$ (the *section*) and $r : Y \longrightarrow X$ (the *retraction*) such that $r \circ s = \text{id}_X$, the identity mapping on X .

We call X a *b-retract* of Y , if X is a retract of Y such that $s(X)$ is b -dense in Y .

Remark 2.10 ([3, 4]). (1) Every section $s : X \longrightarrow Y$ is an embedding.
(2) Every retraction $r : Y \longrightarrow X$ is quotient.

Proposition 2.11. *Let $X, Y \in \mathbf{Top}_0$. If X is a b -retract of Y , then X is homeomorphic to Y .*

PROOF. Suppose X is a b -retract of Y . Then there are two continuous mappings $s : X \longrightarrow Y$ and $r : Y \longrightarrow X$ such that $r \circ s = \text{id}_X$. We have that $(s \circ r) \circ s = s \circ (r \circ s) = s \circ \text{id}_X = s = \text{id}_Y \circ s$, i.e., the following two diagrams commute:

$$\begin{array}{ccc} X & \xrightarrow{s} & Y \\ & \searrow s & \swarrow \text{id}_Y \\ & & Y \end{array} \quad \begin{array}{ccc} X & \xrightarrow{s} & Y \\ & \searrow \text{id}_X = r \circ s & \swarrow r \\ & & X \\ & \searrow s & \swarrow s \circ r \\ & & Y \end{array}$$

By Remark 2.10, s is a b -dense embedding, and by Lemma 2.8, we have $s \circ r = \text{id}_Y$. Since $r \circ s = \text{id}_X$, s is a homeomorphism. Therefore, X is homeomorphic to Y . \square

3. MAIN RESULTS

Lemma 3.1. *Let X, Y be T_0 spaces, U be an open subset of X , and $y_1, y_2 \in Y$. If $y_1 < y_2$ (i.e., $y_1 \leq y_2$ and $y_1 \neq y_2$) in Y , then the mapping $f : X \rightarrow Y$ defined by*

$$f(x) = \begin{cases} y_1 & x \notin U \\ y_2 & x \in U \end{cases}$$

is continuous.

PROOF. In fact, as $y_1 < y_2$, every open set in Y containing y_1 contains y_2 . Then, for each open subset V of Y , we have that

$$f^{-1}(V) = \begin{cases} X & y_1 \in V (\Rightarrow y_2 \in V) \\ U & y_1 \notin V, y_2 \in V \\ \emptyset & \text{otherwise,} \end{cases}$$

which implies that $f^{-1}(V)$ is open in X . Therefore, f is continuous. \square

Theorem 3.2. *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then each \mathbf{K} -reflection is a b -dense embedding.*

PROOF. Let $X \in \mathbf{Top}_0$ and $\mu_X : X \rightarrow X^k$ be a \mathbf{K} -reflection for X . Since $\mathbf{K} \subseteq \mathbf{Top}_0$ and $\mathbf{K} \not\subseteq \mathbf{Top}_1$, there exists a T_0 space $Y \in \mathbf{K}$ that is not T_1 . Then there exists a point $y_2 \in Y$ such that $\{y_2\}$ is not closed, i.e., $\downarrow y_2 \neq \{y_2\}$. Choose a point $y_1 \in \downarrow y_2 \setminus \{y_2\}$, i.e., $y_1 < y_2$. Note that $y_2 \in Y \setminus \downarrow y_1$. Next, we prove our conclusion in a few steps.

Step 1: μ_X is injective.

Let $x_1, x_2 \in X$ with $x_1 \neq x_2$. Since X is T_0 , either $x_1 \not\leq x_2$ or $x_2 \not\leq x_1$. Without loss of generality, we assume $x_2 \not\leq x_1$. Then there exists an open subset U of X such that $x_2 \in U$ and $x_1 \notin U$. Define $f : X \rightarrow Y$ by

$$f(x) = \begin{cases} y_1 & x \notin U \\ y_2 & x \in U. \end{cases}$$

Then by Lemma 3.1, f is a continuous mapping, and clearly $f(x_1) = y_1$ and $f(x_2) = y_2$. Since μ_X is a \mathbf{K} -reflection, there exists a unique continuous mapping $g : X^k \rightarrow Y$ such that $g \circ \mu_X = f$:

$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow f & \vdots g \\ & & Y \end{array}$$

We have that $g(\mu_X(x_1)) = f(x_1) = y_1$ and $g(\mu_X(x_2)) = f(x_2) = y_2$, so that $g(\mu_X(x_1)) \neq g(\mu_X(x_2))$, which implies that $\mu_X(x_1) \neq \mu_X(x_2)$. Hence, μ_X is injective.

Step 2: μ_X is an embedding.

Let V be any open subset of X . Define a mapping $\tilde{f}: X \rightarrow Y$ as follows:

$$\tilde{f}(x) = \begin{cases} y_1 & x \notin V \\ y_2 & x \in V. \end{cases}$$

By Lemma 3.1, \tilde{f} is a continuous mapping. Then there exists a unique continuous mapping $\tilde{g}: X^k \rightarrow Y$ such that $\tilde{g} \circ \mu_X = \tilde{f}$:

$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow \tilde{f} & \downarrow \tilde{g} \\ & & Y \end{array}$$

Claim 1: $\mu_X(V) = \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$.

If $x \in V$, then $\tilde{g}(\mu_X(x)) = \tilde{f}(x) = y_2 \in Y \setminus \downarrow y_1$, which implies that $\mu_X(x) \in \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$, and this proves $\mu_X(V) \subseteq \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$. Conversely, if $\mu_X(x) \in \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$, then $\tilde{f}(x) = \tilde{g}(\mu_X(x)) \in Y \setminus \downarrow y_1$. It follows that $\tilde{f}(x) = y_2$, which is equivalent to $x \in V$, so $\mu_X(x) \in \mu_X(V)$. This implies that $\tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X) \subseteq \mu_X(V)$. Therefore, $\mu_X(V) = \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$.

Note that $Y \setminus \downarrow y \in \mathcal{O}(Y)$ and \tilde{g} is continuous, so $\tilde{g}^{-1}(Y \setminus \downarrow y_1) \in \mathcal{O}(X^k)$. By Claim 1, it follows that $\mu_X(V) = \tilde{g}^{-1}(Y \setminus \downarrow y_1) \cap \mu_X(X)$ is open in the subspace $\mu_X(X)$ of X^k . Thus, together with Step 1, we have that μ_X is an embedding, proving Step 2.

Step 3: $\mu_X(X)$ is b -dense in X^k .

Suppose, on the contrary, $\mu_X(X)$ is not b -dense in X^k . Then there exist a point $z_0 \in X^k$ and an open subset W of X^k such that $z_0 \in W$, and $\downarrow z_0 \cap W \cap \mu_X(X) = \emptyset$. Define two mappings $g_1, g_2: X^k \rightarrow Y$ by

$$g_1(z) = \begin{cases} y_1 & z \notin W \\ y_2 & z \in W, \end{cases} \quad g_2(z) = \begin{cases} y_1 & z \notin W \setminus \downarrow z_0 \\ y_2 & z \in W \setminus \downarrow z_0. \end{cases}$$

Then by Lemma 3.1, g_1, g_2 are both continuous mappings. It should note that $g_1 \neq g_2$ as $g_1(z_0) = y_2$ and $g_2(z_0) = y_1$.

Claim 2: $g_1 \circ \mu_X = g_2 \circ \mu_X$.

Let $x \in X$. If $g_1(\mu_X(x)) = y_2$, then $\mu_X(x) \in W$, and since $\downarrow z_0 \cap W \cap \mu_X(X) = \emptyset$, it follows that $\mu_X(x) \notin \downarrow z_0$, and hence $\mu_X(x) \in W \setminus \downarrow z_0$, which implies

$g_2(\mu_X(x)) = y_2$. If $g_1(\mu_X(x)) = y_1$, then $\mu_X(x) \notin W$, and hence $\mu_X(x) \notin W \setminus \downarrow z_0$, which implies $g_2(\mu_X(x)) = y_1$. Then we have that $g_1 \circ \mu_X = g_2 \circ \mu_X$:

$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow g_1 \circ \mu_X = g_2 \circ \mu_X & \downarrow g_1, g_2 \\ & & Z \end{array}$$

Since μ_X is a \mathbf{K} -reflection for X , we have that $g_1 = g_2$, which contradicts that $g_1 \neq g_2$. Therefore, $\downarrow z_0 \cap W \cap \mu_X(X) \neq \emptyset$, which implies that $\mu_X(X)$ is b -dense in X^k .

All these show that the \mathbf{K} -reflection μ_X is a b -dense embedding. \square

Remark 3.3. It is noteworthy that there is a difference between b -dense embedding of Definition 2.7 and Skula's b -embedding, where the "embedding" condition is not required in the definition of Skula's b -embedding (see [11, Definition 3.1]). As a consequence, Theorem 3.2 above cannot be deduced directly from the result in [11, Theorem 3.5]. Undoubtedly, Skula's method plays an important role in the proof of Theorem 3.2, while for the sake of completeness, some parts of the proof overlap slightly.

Lemma 3.4 ([1]). *Let $f : X \rightarrow Y$ be a continuous mapping between T_0 spaces. Then f is an epimorphism if and only if $f(X)$ is b -dense.*

By Theorem 3.2 and Lemma 3.4, we have the following result.

Corollary 3.5. *If \mathbf{K} is a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$, then it is epireflective.*

The following lemma follows directly from Lemma 2.4.

Lemma 3.6. *Let X be a T_0 space and Y be a subset of X . If Y is b -dense in X , then $X = \uparrow Y$.*

Theorem 3.7. *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. If $X \in \mathbf{K}$ and Y is a saturated subspace of X , then $Y \in \mathbf{K}$.*

PROOF. Let $\mu_Y : Y \rightarrow Y^k$ be the \mathbf{K} -reflection for Y . By Theorem 3.2, $\mu_Y(Y)$ is b -dense in Y^k , and by Lemma 3.6, we have $Y^k = \uparrow \mu_Y(Y)$. Let $e : Y \rightarrow X$ be the inclusion mapping. Then there exists a unique continuous mapping $f : Y^k \rightarrow X$

such that $f \circ \mu_Y = e$:

$$\begin{array}{ccc} Y & \xrightarrow{\mu_Y} & Y^k \\ & \searrow e & \downarrow f \\ & & X \end{array}$$

Claim: $f(Y^k) = Y$. On the one hand, $Y = e(Y) = f(\mu_Y(Y)) \subseteq f(Y^k)$. On the other hand, note that each continuous mapping is order-preserving and $Y^k = \uparrow \mu_Y(Y)$, so $f(Y^k) = f(\uparrow \mu_Y(Y)) \subseteq \uparrow f(\mu_Y(Y)) = \uparrow e(Y) = \uparrow Y = Y$. Hence, $f(Y^k) = Y$.

Now consider the co-restriction $\hat{f}: Y^k \rightarrow Y$ (i.e., $\hat{f}(x) = f(x)$) of f . Then it is clear that $\hat{f} \circ \mu_Y = \text{id}_Y$, the identity mapping on Y :

$$\begin{array}{ccc} Y & \xrightarrow{\mu_Y} & Y^k \\ & \searrow \hat{f} \circ \mu_Y = \text{id}_Y & \downarrow \hat{f} \\ & & Y \end{array}$$

We then deduce that Y is a b -retract of Y^k , hence by Proposition 2.11, Y is homeomorphic to Y^k . As \mathbf{K} is required to be replete, we obtain that $Y \in \mathbf{K}$. \square

4. THE NON-REFLECTIVITY OF OPEN WELL-FILTERED SPACES

Definition 4.1. [12] Let X be a T_0 space.

- (1) For any $U, V \in \mathcal{O}(X)$, we define $U \ll V$ if and only if each open cover of V has a finite subfamily that covers U .
- (2) A subfamily $\mathcal{F} \subseteq \mathcal{O}(X)$ is called a \ll -filtered family if for any $U_1, U_2 \in \mathcal{F}$, there exists $U_3 \in \mathcal{F}$ such that $U_3 \ll U_1$ and $U_3 \ll U_2$.
- (3) X is called *open well-filtered* if for each \ll -filtered family $\mathcal{F} \subseteq \mathcal{O}(X)$ and $U \in \mathcal{O}(X)$,

$$\bigcap \mathcal{F} \subseteq U \Rightarrow V \subseteq U \text{ for some } V \in \mathcal{F}.$$

The category of all open well-filtered spaces with continuous mappings is denoted by **Owf**. Then **Owf** is a full subcategory of **Top**₀.

On open well-filtered spaces, a natural question is

- whether the category **Owf** is reflective in **Top**₀?

In the following, we will give a negative answer to the above question.

Recall that a subset U of a poset P is *Scott open* if (i) $U = \uparrow U$ and (ii) for any directed subset D of P for which $\bigvee D$ exists, $\bigvee D \in U$ implies $D \cap U \neq \emptyset$. All Scott open subsets of P form a topology, and we call this topology the *Scott*

topology on P and denote it by $\sigma(P)$. The space $\Sigma P = (P, \sigma(P))$ is called the *Scott space* of P .

Let \mathbb{N}_{cof} be the set \mathbb{N} of natural numbers with the cofinite topology (the open sets are \emptyset , \mathbb{N} , and all the complements of finite subsets of \mathbb{N}). Let $\mathbb{J} = \mathbb{N} \times (\mathbb{N} \cup \{\infty\})$ be the Johnstone's dcpo [4, 7], which is ordered by $(m, n) \leq (m', n')$ iff either $m = m'$ and $n \leq n'$ or $n' = \infty$ and $n \leq m'$ (see Figure 1).




FIGURE 1. The Johnstone's dcpo \mathbb{J}

Remark 4.2. The following well-known results on \mathbb{N}_{cof} and $\Sigma\mathbb{J}$ will be used.

- (1) Every set in \mathbb{N}_{cof} is compact.
- (2) $\forall U, V \in \sigma(\mathbb{J})$, $U \ll V$ if and only if $U = \emptyset$ (see Exercise 5.2.15 in [4]).
- (3) The maximal points space $\text{Max}(\mathbb{J})$ of \mathbb{J} with the relative topology of $\Sigma\mathbb{J}$ is homeomorphic to \mathbb{N}_{cof} .

Theorem 4.3. *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 . If $\Sigma\mathbb{J} \in \mathbf{K}$, then $\mathbb{N}_{\text{cof}} \in \mathbf{K}$.*

PROOF. If $\Sigma\mathbb{J} \in \mathbf{K}$, then by Remark 4.2(3) the maximal points space $\text{Max}(\mathbb{J})$ is a saturated subspace of $\Sigma\mathbb{J}$, which is homeomorphic to \mathbb{N}_{cof} . Then by Theorem 3.7, $\mathbb{N}_{\text{cof}} \in \mathbf{K}$. \square

Proposition 4.4. *The Scott topology $\sigma(\mathbb{J})$ is open well-filtered.*

PROOF. Let \mathcal{F} be a \ll -filtered family of Scott open subsets of \mathbb{J} , and $U \in \sigma(\mathbb{J})$ such that $\bigcap \mathcal{F} \subseteq U$. Fix a member $V_0 \in \mathcal{F}$. Since \mathcal{F} is \ll -filtered, there exists

$V_1 \in \mathcal{F}$ such that $V_1 \ll V_0$, and from Remark 4.2(2), it follows that $V_1 = \emptyset \subseteq U$. Therefore, $\sigma(\mathbb{J})$ is open well-filtered. \square

Proposition 4.5. *The space \mathbb{N}_{cof} is not open well-filtered.*

PROOF. Since each set in \mathbb{N}_{cof} is compact, $U \ll V$ iff $U \subseteq V$ for any open subsets U, V of \mathbb{N}_{cof} . For each $n \in \mathbb{N}$, define $U_n = \{k \in \mathbb{N} : k \geq n\}$. Then $\{U_n : n \in \mathbb{N}\}$ is a \ll -filtered family of open sets in \mathbb{N}_{cof} such that $\bigcap_{n \in \mathbb{N}} U_n = \emptyset$, but $U_n \neq \emptyset$ for any $n \in \mathbb{N}$. This implies \mathbb{N}_{cof} is not open well-filtered. \square

From Propositions 4.4 and 4.5, we have that $\mathbb{N}_{\text{cof}} \notin \mathbf{Owf}$ and $\Sigma\mathbb{J} \in \mathbf{Owf}$. Hence, by Theorem 4.3, we have the following result.

Theorem 4.6. *The category \mathbf{Owf} is not reflective in \mathbf{Top}_0 .*

5. CONCLUSION

In this paper, we use Skula's b -topology to study the reflective subcategories of \mathbf{Top}_0 , and prove the non-reflectivity of open well-filtered spaces in \mathbf{Top}_0 . These results enrich the b -topology applications in domain theory, which fully illustrates that Skula's b -topology is an effective tool to study the reflectivity of subcategories of T_0 spaces. It also provides a possibility of solving some open problems on the reflectivity of some other T_0 spaces in domain theory (see [13] for example) by using the approach in this paper. We leave these as our future work.

Acknowledge The authors would like to thank the anonymous referee for the constructive suggestions that helped to improve the paper.

REFERENCES

- [1] S. Baron, Note on epi in T_0 , Canad. Math. Bull. 11 (1968) 503–504.
- [2] C. H. Dowker, D. Papert, Quotient frames and subspaces, Proc. London Math. Soc. 3 (16) (1966) 275–296.
- [3] G. Gierz, K. Hofmann, K. Keimel, J. Lawson, M. Mislove, D. Scott, Continuous lattices and Domains, Encyclopedia of Mathematics and Its Applications, Vol. 93, Cambridge University Press, 2003.
- [4] J. Goubault-Larrecq, Non-Hausdorff Topology and Domain Theory, Cambridge University Press, 2013.
- [5] R-E. Hoffmann, Topological functors admitting generalized Cauchy-completions, in Categorical Topology, Proc. of the conf. held at Mannheim 1975 ed. E. Binz and H. Herrlich, pp. 286–344, Lect. Notes in Math., 540, Berlin-Heidelberg-New York: Springer 1976.
- [6] X. Jia, Meet-continuity and locally compact sober dcpos, PhD thesis, University of Birmingham, 2018.

- [7] P.T. Johnstone, Scott is not always sober, In: Banaschewski B., Hoffmann RE. (eds) Continuous Lattices. Lecture Notes in Mathematics, vol. 871 (1981) 282-283. Springer, Berlin, Heidelberg, <https://doi.org/10.1007/BFb0089911>.
- [8] K. Keimel, J.D. Lawson, D -completions and the d -topology, Ann. Pure Appl. Logic 159 (2009) 292–306.
- [9] J.F. Kennison, Reflective functors in general topology and elsewhere, Trans. Amer. Math. Soc. 118 (1965) 303–315.
- [10] J. Lawson, G. Wu and X. Xi, Well-filtered spaces, compactness, and the lower topology, Houston J. Math. 46 (2020) 283–294.
- [11] L. Skula, On a reflective subcategory of the category of all topological spaces, Trans. Amer. Math. Soc. 142 (1969) 37–41.
- [12] C. Shen, X. Xi, X. Xu, D. Zhao, On open well-filtered spaces, Logic Meth. Computer Sci. 16 (4) (2020) 4–18.
- [13] X. Xu, D. Zhao, Some open problems on well-filtered spaces and sober spaces, Topol. Appl. (2020) 107540.

Received August 21, 2022

Revised version received November 20, 2022

CHONG SHEN, SCHOOL OF SCIENCE, BEIJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, BEIJING, CHINA

Email address: shenchong0520@163.com

XIAOYONG XI, SCHOOL OF MATHEMATICS AND STATISTICS, YANCHENG TEACHERS UNIVERSITY, YANCHENG, JIANGSU, CHINA

Email address: xixy@yctu.edu.cn

DONGSHENG ZHAO, MATHEMATICS AND MATHEMATICS EDUCATION, NATIONAL INSTITUTE OF EDUCATION, NANYANG TECHNOLOGICAL UNIVERSITY, 1 NANYANG WALK, SINGAPORE 637616

Email address: dongsheng.zhao@nie.edu.sg