

Chapter 11. Dynamic models and their simulation by Euler's method

Objectives

- ① Dynamic systems can often be represented in terms of *flows* between *compartments*. We will develop the concept of a **compartment model** for which we specify *rates* for the flows between compartments.
- ② We develop deterministic and stochastic interpretations of a compartment model.
- ③ We introduce Euler's method to simulate from dynamic models, and we apply it to both deterministic and stochastic compartment models.

Compartment models

- A **compartment model** is a model associated with a **flow diagram** specifying how objects move between categories, called **compartments**.
- It is often useful to represent systems by **flow diagrams**. We will need equations to specify formally what the flow diagram means.
- One major applications of compartment models is pharmacokinetics, the study of how pharmacological drugs enter the body, move between organs, metabolize, and leave. The compartments may be the organs; the flow is movement of the drug and its metabolites between organs.
- Another major application of compartment models is epidemiology. Compartments are groups of people; the flow is the movement of an infectious disease through the population.

Compartment models in epidemiology: the SIR model and its generalizations

We will develop deterministic and stochastic representations of a susceptible-infected-recovered (SIR) system, a fundamental class of models for disease transmission dynamics. We will do this using notation which generalizes to more complex systems

[[@breto09]](<http://dept.stat.lsa.umich.edu/ionides/pubs/breto09.pdf>).

General notation for compartment models

- To develop a systematic notation, it turns out to be convenient to keep track of the flows between compartments as well as the number of individuals in each compartment. Let

$$N_{SI}(t)$$

count the number of individuals who have transitioned from S to I by time t . We say that $N_{SI}(t)$ is a **counting process**. A similarly constructed process

$$N_{IR}(t)$$

counts individuals transitioning from I to R . To include demography, we could keep track of birth and death events by the counting processes $N_{\bullet S}(t)$, $N_{S\bullet}(t)$, $N_{I\bullet}(t)$, $N_{R\bullet}(t)$.

- For discrete population compartment models, the flow counting processes are non-decreasing and integer valued.
- For continuous population compartment models, the flow counting processes are non-decreasing and real valued.
- The numbers of people in each compartment can be computed via these counting processes. Ignoring demography, we have:

The ordinary differential equation (ODE) interpretation of the SIR model

Together with initial conditions specifying $S(0)$, $I(0)$ and $R(0)$, we just need to write down ODEs for the flow counting processes. These are,

$$dN_{SI}/dt = \mu_{SI}(t) S(t),$$

$$dN_{IR}/dt = \mu_{IR} I(t).$$

The simple continuous-time Markov chain interpretation of the SIR model

- Continuous-time Markov chains are the basic tool for building discrete population epidemic models.
- The Markov property lets us specify a model by the transition probabilities on small intervals (together with the initial conditions). For the SIR model, we have

$$\begin{aligned}\mathbb{P}[N_{SI}(t + \delta) = N_{SI}(t) + 1] &= \mu_{SI}(t) S(t)\delta + o(\delta) \\ \mathbb{P}[N_{SI}(t + \delta) = N_{SI}(t)] &= 1 - \mu_{SI}(t) S(t)\delta + o(\delta) \\ \mathbb{P}[N_{IR}(t + \delta) = N_{IR}(t) + 1] &= \mu_{IR}(t) I(t)\delta + o(\delta) \\ \mathbb{P}[N_{IR}(t + \delta) = N_{IR}(t)] &= 1 - \mu_{IR}(t) I(t)\delta + o(\delta)\end{aligned}$$

- Here, we are using [little o notation] We write

$$h(\delta) = o(\delta)$$

to mean

$$\lim_{\delta \rightarrow 0} \frac{h(\delta)}{\delta} = 0.$$

Question: What is the link between little o notation and the derivative?

- Explain why

$$f(x + \delta) = f(x) + \delta g(x) + o(\delta)$$

is the same statement as

$$\frac{df}{dx} = g(x).$$

- What considerations might help you choose which of these notations to use?

Question 11.1. From Markov chain to ODE. Find the expected value of $N_{SI}(t + \delta) - N_{SI}(t)$ and $N_{IR}(t + \delta) - N_{IR}(t)$ given the current state, $S(t)$, $I(t)$ and $R(t)$. Take the limit as $\delta \rightarrow 0$ and show that this gives the ODE model.

- A **simple counting process** is one which cannot count more than one event at a time.
- Technically, the SIR Markov chain model we have written is simple. One may want to model the extra randomness resulting from multiple simultaneous events: someone sneezing in a bus; large gatherings at football matches; etc. This extra randomness may even be critical to match the variability in data.
- Later in the course, we may see situations where this extra

Euler's method for ordinary differential equations (ODEs)

- Euler (1707-1783) wanted a numeric solution of an ODE $dx/dt = h(x)$ with an initial condition $x(0)$.
- He supposed this ODE has some true solution $x(t)$ which could not be worked out analytically. He therefore wished to approximate $x(t)$ numerically.
- He initialized the numerical solution at the known starting value,

$$\tilde{x}(0) = x(0).$$

- For $k = 1, 2, \dots$, he supposed that the gradient dx/dt is approximately constant over the small time interval $k\delta \leq t \leq (k+1)\delta$.
- Therefore, he defined

$$\tilde{x}((k+1)\delta) = \tilde{x}(k\delta) + \delta h(\tilde{x}(k\delta)).$$

This only defines $\tilde{x}(t)$ when t is a multiple of δ , but suppose $\tilde{x}(t)$ is constant between these discrete times.

- We now have a numerical scheme, stepping forwards in time increments of size δ , that can be readily evaluated by computer (or by hand, if you are Euler).

Euler's method versus other numerical methods

- Mathematical analysis of Euler's method says that, as long as the function $h(x)$ is not too exotic, then $x(t)$ is well approximated by $\tilde{x}(t)$ when the discretization time-step, δ , is sufficiently small.
- Euler's method is not the only numerical scheme to solve ODEs. More advanced schemes have better convergence properties, meaning that the numerical approximation is closer to $x(t)$. However, there are 3 reasons we choose to lean heavily on Euler's method:
 - ① Euler's method is the simplest (following the KISS principle).
 - ② Euler's method extends naturally to stochastic models, both continuous-time Markov chains models and stochastic differential equation (SDE) models.
 - ③ Close approximation of the numerical solutions to a continuous-time model is less important than it may at first appear, a topic to be discussed.

Some comments on using continuous-time models and discretized approximations

- In some physical and engineering situations, a system follows an ODE model closely. For example, Newton's laws provide a very good approximation to the motions of celestial bodies.
- In many biological situations, ODE models only become close mathematical approximations to reality at reasonably large scale. On small temporal scales, models cannot usually capture the full scope of biological variation and biological complexity.
- If we are going to expect substantial error in using $x(t)$ to model a biological system, maybe the numerical solution $\tilde{x}(t)$ represents the system being modeled as well as $x(t)$ does.
- If our model fitting, model investigation, and final conclusions are all based on our numerical solution $\tilde{x}(t)$ (i.e., we are sticking entirely to simulation-based methods) then we are most immediately concerned with how well $\tilde{x}(t)$ describes the system of interest. $\tilde{x}(t)$ becomes more important than the original model, $x(t)$.
- When following this perspective, it is important that the scientists fully describe the numerical model $\tilde{x}(t)$. The main advantages of the

Euler's method for a discrete SIR model

- Recall the simple continuous-time Markov chain interpretation of the SIR model without demography:

$$\begin{aligned}\mathbb{P}[N_{SI}(t + \delta) = N_{SI}(t) + 1] &= \mu_{SI}(t) S(t)\delta + o(\delta), \\ \mathbb{P}[N_{IR}(t + \delta) = N_{IR}(t) + 1] &= \mu_{IR} I(t)\delta + o(\delta).\end{aligned}$$

- We look for a numerical solution with state variables $\tilde{S}(k\delta)$, $\tilde{I}(k\delta)$, $\tilde{R}(k\delta)$.
- The counting processes for the flows between compartments are $\tilde{N}_{SI}(t)$ and $\tilde{N}_{IR}(t)$. The counting processes are related to the numbers of individuals in the compartments by the same flow equations we had before:

$$\begin{aligned}\tilde{S}(k\delta) &= S(0) - \tilde{N}_{SI}(k\delta) \\ \tilde{I}(k\delta) &= I(0) + \tilde{N}_{SI}(k\delta) - \tilde{N}_{IR}(k\delta) \\ \tilde{R}(k\delta) &= R(0) + \tilde{N}_{IR}(k\delta)\end{aligned}$$

- We focus on a numerical solution to $N_{SI}(t)$, since the same methods can also be applied to $N_{IR}(t)$.

- Here are three possibilities, defining a process at times $t = k\delta$ for all integer k and some fixed $\delta > 0$.

- 1 A Poisson approximation.

$$\tilde{N}_{SI}(t + \delta) = \tilde{N}_{SI}(t) + \text{Poisson}[\mu_{SI}(\tilde{I}(t)) \tilde{S}(t) \delta],$$

where $\text{Poisson}(\mu)$ is a Poisson random variable with mean μ and

$$\mu_{SI}(\tilde{I}(t)) = \beta \tilde{I}(t).$$

- 2 A binomial approximation with transition probabilities approximated by rate times time.

$$\tilde{N}_{SI}(t + \delta) = \tilde{N}_{SI}(t) + \text{Binomial}[\tilde{S}(t), \mu_{SI}(\tilde{I}(t)) \delta],$$

where $\text{Binomial}(n, p)$ is a binomial random variable with mean np and variance $np(1 - p)$.

- 3 A binomial approximation with exponential transition probabilities.

$$\tilde{N}_{SI}(t + \delta) = \tilde{N}_{SI}(t) + \text{Binomial}[\tilde{S}(t), 1 - \exp\{-\mu_{SI}(\tilde{I}(t))\delta\}].$$

- Conceptually, it is simplest to think of (1) or (2). Numerically, it is usually preferable to implement (3).

Theoretical exercise: Compartment models via stochastic differential equations (SDEs)

The Euler method extends naturally to stochastic differential equations. A natural way to add stochastic variation to an ODE $dx/dt = h(x)$ is

$$dX/dt = h(X) + \sigma dB/dt$$

where $\{B(t)\}$ is Brownian motion and so dB/dt is Brownian noise. Then, an Euler approximation $\tilde{X}(t)$ is generated by

$$\tilde{X}((k+1)\delta) = \tilde{X}(k\delta) + \delta h(\tilde{X}(k\delta)) + \sigma\sqrt{\delta} Z_k$$

where Z_1, Z_2, \dots is a sequence of independent standard normal random variables, i.e., $Z_k \sim N[0, 1]$. Although SDEs are often considered an advanced topic in probability, the Euler approximation doesn't demand much more than familiarity with the normal distribution.

- Write down a stochastic Euler method for an SDE representation of the SIR model. Consider some difficulties that might arise with non-negativity constraints, and propose some practical way one might deal with that issue.
- A useful method to deal with positivity constraints is to use Gamma

Conceptual exercise: Euler's method vs Gillespie's algorithm

- A widely used, exact simulation method for continuous time Markov chains is Gillespie's algorithm. We do not put much emphasis on Gillespie's algorithm here. Why? When would you prefer an implementation of Gillespie's algorithm to an Euler solution?
- Numerically, Gillespie's algorithm is often approximated using so-called tau-leaping methods. These are closely related to Euler's approach. Is it reasonable to call a suitable Euler approach a tau-leaping method?

Compartmental models in pomp

As an example that we can probe in some depth, we look at an isolated outbreak of influenza that occurred in a boarding school for boys in England [anonymous78]. Let's examine the data:

```
bsflu <- read.table("bsflu_data.txt")  
head(bsflu)
```

```
##           B  C day  
## 1978-01-22   1  0   1  
## 1978-01-23   6  0   2  
## 1978-01-24  26  0   3  
## 1978-01-25  73  1   4  
## 1978-01-26 222  8   5  
## 1978-01-27 293 16   6
```

The variable B refers to boys confined to bed and C to boys in convalescence. Let's restrict our attention for the moment to the B variable.

```
bsflu <- subset(bsflu, select=c(day, B))  
ggplot(data=bsflu, aes(x=day, y=B))+geom_line()+geom_point()
```


Viewing the SIR as a POMP model

- The unobserved state variables, in this case, are the numbers of individuals, S , I , R in the S, I, and R compartments, respectively.
- It is reasonable in this case to view the population size $N = S + I + R$, as fixed.
- The numbers that actually move from one compartment to another over any particular time interval are modeled as stochastic processes.
- In this case, we'll assume that the stochasticity is purely demographic, i.e., that each individual in a compartment at any given time faces the same risk of exiting the compartment.

Implementing the SIR POMP model in pomp

A starting point is to construct `rprocess`, a stochastic simulator for the unobserved state process. We saw several numerical approaches to do this. An attractive option here is to model the number moving from one compartment to the next over a very short time interval as a binomial random variable. In particular, we model the number, ΔN_{SI} , moving from S to I over interval Δt as

$$\Delta N_{SI} \sim \text{Binomial} \left(S, 1 - e^{-\lambda \Delta t} \right),$$

and the number moving from I to R as

$$\Delta N_{IR} \sim \text{Binomial} \left(I, 1 - e^{-\gamma \Delta t} \right).$$

A Csnippet that encodes such a simulator is as follows:

```
sir_step <- Csnippet("  
  double dN_SI = rbinom(S,1-exp(-Beta*I/N*dt));  
  double dN_IR = rbinom(I,1-exp(-gamma*dt));  
  S -= dN_SI;  
  I += dN_SI - dN_IR;  
  R += dN_IR;  
")
```

Toward a measurement model

Now let's assume that the case reports, B , result from a process by which new infections result in confinement with probability ρ , which we can think of as the probability that an infection is severe enough to be noticed by the school authorities. Since confined cases have, presumably, a much lower transmission rate, let's treat B as being a count of the number of boys who have moved from I to R over the course of the past day. We need a variable to track this. Let's modify our Csnippet above, adding a variable H to track the incidence. We'll then replace the rprocess with the new one.

```
sir_step <- Csnippet("  
  double dN_SI = rbinom(S,1-exp(-Beta*I/N*dt));  
  double dN_IR = rbinom(I,1-exp(-gamma*dt));  
  S -= dN_SI;  
  I += dN_SI - dN_IR;  
  R += dN_IR;  
  H += dN_IR;  
")
```

Now, we'll model the data, B , as a binomial process,

$$B_t \sim \text{Binomial}(H(t) - H(t-1), \rho).$$

But we have a problem, since at time t , the variable H we've defined will contain $H(t)$, not $H(t) - H(t-1)$. We can overcome this by telling `pomp` that we want H to be set to zero immediately following each observation. We do this by setting the `zernames` argument to `pomp`:

```
pomp(sir, zernames="H") -> sir
```

Now, to include the observations in the model, we must write both a `dmeasure` and an `rmeasure` component:

```
dmeas <- Csnippet("lik = dbinom(B,H,rho,give_log);")  
rmeas <- Csnippet("B = rbinom(H,rho);")
```

and put these into our `pomp` object:

```
sir <- pomp(sir, rmeasure=rmeas, dmeasure=dmeas, statenames="H", paramnames=)
```

A good first step is to perform some simulations to check whether things are working. To do so, we need to choose parameters. A little thought will get us some ballpark estimates. In the data, it looks like there were a total of 1540 infections, so the population size, N , must be somewhat in excess of this number. In fact, we can use the final-size equation

$$R_0 = -\frac{\log(1-f)}{f},$$

where $f = R(\infty)/N$ is the final size of the epidemic, together with the idea that R_0 must be, say, around 1.5, to estimate that $f \approx 0.6$, whence $N \approx 2600$. If the infectious period is roughly 1 day, then $1/\gamma \approx 1$ da and $\beta = \gamma R_0 \approx 1.5 \text{ da}^{-1}$.

```
sims <- simulate(sir, params=c(Beta=1.5, gamma=1, rho=0.9, N=2600),  
                 nsim=20, format="data.frame", include=TRUE)  
ggplot(sims, mapping=aes(x=day, y=B, group=.id, color=.id=="data")) +  
  geom_line() + guides(color=FALSE)
```



Question 11.2. Explore the SIR model by simulation. Fiddle with the parameters to see if you can't find parameters for which the data are a more plausible realization.

Extending the SIR model to include latency: The SEIR model

Below is a diagram of the so-called SEIR model. This differs from the SIR model in that infected individuals must pass a period of latency before becoming infectious.

Question 11.3. Modify the codes above to construct a pomp object containing the flu data and an SEIR model. Perform simulations as above

Rethinking the boarding-school flu data

In the preceding, we've been assuming that B_t represents the number of boys **sent** to bed on day t . Actually, this isn't correct at all. As described in the report (?), B_t represents the total number of boys **in** bed on day t . Since boys were potentially confined for more than one day, the data count each infection multiple times. On the other hand, we have information about the total number of boys at risk and the total number who were infected. In fact, we know from @anonymous78 that $N = 763$ boys were at risk and 512 boys in total spent between 3 and 7 days away from class (either in bed or convalescent). Moreover, there is information in the data on the number of boys, C_t , convalescent at day t . Since $1540 \text{ boy-da} / 512 \text{ boy} \approx 3 \text{ da}$, we know that the average duration spent in bed was 3 days and, since $\sum_t C_t = 0$, we can infer that the average time spent convalescing was $0 \text{ boy-da} / 512 \text{ boy} \approx 0 \text{ da}$.

```
require(reshape2)
ggplot(data=melt(bsflu, id="day"), mapping=aes(x=day, y=value, color=var))
  geom_line()+geom_point()
```

300



Question 11.4. Formulate a model with both confinement and convalescent stages. Implement it in `pomp` using a compartmental model like that diagrammed below.

You will have to give some thought to just how to model the relationship between the data (B and C) and the state variables. How many parameters can reasonably be fixed? How many must be estimated? Obtain some ballpark estimates of the parameters and simulate to see if you can plausibly explain the data as a realization of this model.

Acknowledgments and License

- Produced with R version 3.6.2 and pomp version 2.3.
- These notes build on previous versions at ionides.github.io/531w16 and ionides.github.io/531w18.
- Those notes draw on material developed for a short course on Simulation-based Inference for Epidemiological Dynamics (<http://kingaa.github.io/sbied/>) by Aaron King and Edward Ionides, taught at the University of Washington Summer Institute in Statistics and Modeling in Infectious Diseases, from 2015 through 2018.
- Licensed under the Creative Commons attribution-noncommercial license, <http://creativecommons.org/licenses/by-nc/3.0/>. Please share and remix noncommercially, mentioning its origin.



References I