



Tropical Spectrahedra

Xavier Allamigeon¹ · Stéphane Gaubert¹ · Mateusz Skomra² 

Received: 15 September 2017 / Revised: 14 December 2019 / Accepted: 8 January 2020 /

Published online: 18 February 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

We introduce tropical spectrahedra, defined as the images by the nonarchimedean valuation of spectrahedra over the field of real Puiseux series. We provide an explicit polyhedral characterization of generic tropical spectrahedra, involving principal tropical minors of size at most 2. One of the key ingredients is Denef–Pas quantifier elimination result over valued fields. We obtain from this that the nonarchimedean valuation maps semialgebraic sets to semilinear sets that are closed. We also prove that, under a regularity assumption, the image by the valuation of a basic semialgebraic set is obtained by tropicalizing the inequalities which define it.

Keywords Spectrahedra · Tropical geometry · Valued fields · Nonarchimedean fields

Mathematics Subject Classification 14T05 · 90C22 · 14P10 · 12J25

1 Introduction

Spectrahedra are one of the main generalizations of polyhedra. They are real convex semialgebraic sets, defined by a single matrix inequality of the form

Editor in Charge: János Pach

Xavier Allamigeon, Stéphane Gaubert, and Mateusz Skomra were partially supported by the ANR projects CAFEIN (ANR-12-INSE-0007) and MALTHY (ANR-13-INSE-0003), by the PGMO program of EDF and Fondation Mathématique Jacques Hadamard, and by the “Investissement d’avenir”, référence ANR-11-LABX-0056-LMH, LabEx LMH. M. Skomra was supported by a grant from Région Île-de-France.

Xavier Allamigeon
Xavier.Allamigeon@inria.fr

Stéphane Gaubert
Stephane.Gaubert@inria.fr

Mateusz Skomra
mateusz.skomra@ens-lyon.fr

¹ INRIA and CMAP, École Polytechnique, IP Paris, CNRS, 91128 Palaiseau Cedex, France

² Université de Lyon, EnsL, UCBL, CNRS, LIP, 69342 Lyon Cedex 07, France

$$Q^{(0)} + x_1 Q^{(1)} + \cdots + x_n Q^{(n)} \succcurlyeq 0,$$

where $Q^{(0)}, \dots, Q^{(n)}$ are real symmetric matrices, and \succcurlyeq denotes the Loewner order. (Recall that, for real symmetric matrices A, B , the *Loewner order* is such that $A \succcurlyeq B$ if all the eigenvalues of $A - B$ are nonnegative.) Spectrahedra arise in a number of applications from engineering sciences and combinatorial optimization [14,24]. Several theoretical questions concerning spectrahedra have been raised, such as the Helton–Nie conjecture [25], recently disproved by Scheiderer [35]. Other, like the generalized Lax conjecture [39] or the complexity of checking the emptiness [33], are still unsettled.

Spectrahedra can be considered more generally over any real closed field. In tropical geometry, one often works with the usual field of real Puiseux series (with rational exponents), or with a larger field of generalized Puiseux series with real exponents, as we do here.

Main results. In this paper, we introduce tropical spectrahedra. These are defined as the images by the nonarchimedean valuation of spectrahedra over the field of real generalized Puiseux series. Our main result (Theorem 5.17) provides an explicit polyhedral characterization of tropical spectrahedra, when the valuation of the defining matrices satisfies a genericity condition. We suppose first that the defining matrices have a Metzler type sign pattern. Metzler matrices are fundamental in mathematical modelling, since they yield dynamical systems whose flow is order preserving. However, we focus on these matrices because the tropicalization of the (generic) spectrahedra that they define is remarkably transparent (Theorem 5.13). Then, we show how to recover the tropicalization of a spectrahedron whose defining matrices have a general sign pattern by “gluing” tropicalizations of spectrahedra satisfying the Metzler condition. Our results are synthesized by Theorem 5.18, which shows, still *under a genericity condition*, that *the tropicalization of a spectrahedron is obtained by requiring only the positivity of principal tropical minors of order at most 2*. The genericity condition is expressed explicitly as a flow condition in a directed hypergraph (Lemma 5.23 and Theorem 5.24).

To this end, we study, more generally, tropical semialgebraic sets, defined as the images by the nonarchimedean valuation of semialgebraic sets over the field of real generalized Puiseux series. We exploit quantifier elimination methods in valued fields by Denef [19] and Pas [30], which imply that such sets are semilinear. Moreover, we show that tropical semialgebraic sets are always closed. It follows that, under a regularity assumption, the image by the valuation of a basic semialgebraic set is obtained by “tropicalizing” each polynomial inequality arising in the definition of this set. This shows in particular that this image is a polyhedral complex with an explicit description in terms of piecewise linear inequalities (Corollary 4.8).

Related work. A general question, in tropical geometry, consists in providing combinatorial characterizations of nonarchimedean amoebas, i.e., images by the nonarchimedean valuation of algebraic sets over nonarchimedean fields. Kapranov’s theorem on amoebas of hypersurfaces, or Viro’s patchworking method for real algebraic curves [40] and its extensions [13,37], address this question in different settings. In parallel, general results have been developed in model theory of valued fields, in

particular by Weispfenning [42], Denef [18,19], and Pas [30,31]. The fact that nonarchimedean amoebas have a polyhedral structure follows from these works.

Excepting tropical polyhedra, there are few works dealing with tropical semialgebraic sets. The most closely related works are those of Yu [43] and Alessandrini [2].

Yu characterized the image by the nonarchimedean valuation of the cone of positive semidefinite matrices over real Puiseux series, showing that it is determined by 2×2 principal tropical minors. We show that 2×2 , together with 1×1 , tropical minors still determine generic tropical spectrahedra.

Alessandrini studied the log-limits of real semialgebraic sets. His approach, based on o-minimal models, shows in particular that the image by the nonarchimedean valuation of a semialgebraic set over the field of absolutely convergent real generalized Puiseux series is a polyhedral complex. We avoid the recourse to o-minimal techniques by using Denef–Pas quantifier elimination instead. A by-product is that the property holds for any real closed valued field.

Applications and consequences of the main results. A general issue in computational optimization is to develop combinatorial algorithms for semidefinite programming. The present work, providing an explicit characterization of tropical spectrahedra, leads to combinatorial algorithms to solve a class of generic semidefinite feasibility problems over nonarchimedean fields. This is developed in the companion work [6], where it is shown that feasibility problems for generic tropical spectrahedra are equivalent to solving mean payoff stochastic games with perfect information, also known as “turn-based” mean payoff stochastic games. This allows one to apply game algorithms to solve nonarchimedean semidefinite feasibility problems. The reference [6] focuses on algorithmic aspects, relying on the present work for structural results. Conversely, tropical spectrahedra offer new methods to study turn-based mean payoff stochastic games. Recall that the complexity of these games is one of the fundamental unsettled problems in algorithmic game theory: these games are in $\text{NP} \cap \text{coNP}$, but they are not known to be in P (see [8] and the references therein). In the recent joint work of the authors with Katz [5], we associate “primal” and “dual” tropical spectrahedra to a turn-based mean payoff stochastic game. We show that metric characteristics of these spectrahedra define a condition number, allowing one to parametrize the complexity of these games.

The present work has also more theoretical consequences. In [7], we use the characterization of tropical spectrahedra to show that an analogue of the Helton–Nie conjecture over nonarchimedean fields is true “up to taking the valuations”, i.e., the images by the valuation of convex semialgebraic sets over a nonarchimedean field coincide with the images by the valuation of projected spectrahedra, over the same field.

Organization of the paper. In Sect. 2, we recall basic notions and results from tropical geometry and from the theory of valued fields. In Sect. 3, we apply the quantifier elimination results of Denef and Pas to show that tropical semialgebraic sets have a polyhedral structure. This allows us to show, in Sect. 4, that tropical semialgebraic sets are finite unions of closed polyhedra. In Sect. 5, we introduce tropical spectrahedra. We first provide an explicit combinatorial description in the simpler situation in which

the input matrices have a Metzler sign pattern (Sect. 5.2), and subsequently remove this assumption (Sect. 5.3). These results hold under a condition that is shown to be satisfied generically in Sect. 5.4.

2 Preliminaries

2.1 Puiseux Series

The main example of nonarchimedean structure which we use in this paper is the field \mathbb{K} of (formal generalized real) *Puiseux series*. This field is defined as follows. It consists of formal series of the form

$$x = \sum_{i=1}^{\infty} c_{\lambda_i} t^{\lambda_i}, \quad (1)$$

where t is a formal parameter, $(\lambda_i)_{i \geq 1}$ is a strictly decreasing sequence of real numbers that is either finite or unbounded, and $c_{\lambda_i} \in \mathbb{R} \setminus \{0\}$ for all λ_i . There is also a special, empty series, which is denoted by 0. We denote by $\text{lc}(x)$ the coefficient c_{λ_1} of the leading term in the series x , with the convention that $\text{lc}(0) = 0$. The addition and multiplication in \mathbb{K} are defined in a natural way. Moreover, \mathbb{K} can be endowed with a linear order \geq , which is defined as $x \geq y$ if $\text{lc}(x - y) \geq 0$. We denote as $\mathbb{K}_{\geq 0}$ the set of nonnegative series x , i.e., satisfying $x \geq 0$. The *valuation* of an element $x \in \mathbb{K}$ as in (1) is defined as the greatest exponent λ_1 occurring in the series. It is known that \mathbb{K} is a real closed field (see [29] for instance). We use the specific field \mathbb{K} to keep the exposition concrete. Our main results, however, work for any real closed field sent surjectively to \mathbb{R} by a nonarchimedean valuation. Other convenient choices of such fields include the field of Hahn series (series with well-ordered support) with real coefficients and real exponents [34], or its subfield consisting of those series that are absolutely convergent for t large enough, or the subfield of \mathbb{K} consisting of series subject to the same convergence requirement (see [21] for more information on the two latter fields).

2.2 Tropical Algebra

In this section, we briefly introduce the basic concepts of tropical algebra and its connection with the nonarchimedean field of Puiseux series.

We denote by $\text{val}: \mathbb{K} \rightarrow \mathbb{R} \cup \{-\infty\}$ the function which maps a Puiseux series $x \in \mathbb{K}$ to its valuation. We use the convention $\text{val}(0) = -\infty$. It is immediate to see that the map val satisfies the following properties:

$$\text{val}(x + y) \leq \max(\text{val}(x), \text{val}(y)), \quad (2)$$

$$\text{val}(xy) = \text{val}(x) + \text{val}(y) \quad (3)$$

meaning that val is a *nonarchimedean valuation*. Moreover, the equality holds in (2) if the leading terms of x and y do not cancel, which is the case if $\text{val}(x) \neq \text{val}(y)$ or if $x, y \geq 0$.

Loosely speaking, the *tropical semifield* \mathbb{T} can be thought of as the image of \mathbb{K} by the nonarchimedean valuation. We use the “max-plus” convention, so the base set of \mathbb{T} is defined to be $\mathbb{R} \cup \{-\infty\}$. It is endowed with the addition $x \oplus y := \max(x, y)$ and the multiplication $x \odot y := x + y$. The term “semifield” refers to the fact that the addition does not have an opposite law. We use the notation $\bigoplus_{i=1}^n a_i = a_1 \oplus \cdots \oplus a_n$ and $a^{\odot n} = a \odot \cdots \odot a$ (n times). We also endow \mathbb{T} with the standard order \geq . The map val yields an order-preserving morphism of semifields from $\mathbb{K}_{\geq 0}$ to \mathbb{T} . This follows from (3) and from the equality case in (2). We refer the reader to [15, 27] for more information on the tropical semifield.

We note that in the valued fields literature, the valuation map is usually defined using the min-plus notation. More precisely, in the definition of Puiseux series, one usually requires the sequence of exponents $(\lambda_i)_{i \geq 1}$ to be strictly increasing, and then the valuation of a series is defined as the smallest exponent occurring in this series. This leads to a map $\text{val}: \mathbb{K} \rightarrow \mathbb{R} \cup \{+\infty\}$ that satisfies $\text{val}(x_1 + x_2) \geq \min(\text{val}(x_1), \text{val}(x_2))$. This definition is equivalent to the one above—it is enough to replace $\text{val}(x)$ by $-\text{val}(x)$ in order to alternate between the two definitions. In this paper, we prefer to use the max-plus notation, since it yields an order preserving map from $\mathbb{K}_{\geq 0}$ to \mathbb{T} , allowing more direct arguments. Our convention also corresponds to an analytic interpretation of t as a “large positive parameter”, and it is consistent with earlier work of us on related problems, see [4, 6].

It is convenient to keep track not only of the valuation of a series but also of its sign. To this end, we define the *sign* of a series $\mathbf{x} \in \mathbb{K}$ as $+1$ if $\mathbf{x} > 0$, -1 if $\mathbf{x} < 0$, and 0 otherwise. We denote it by $\text{sign}(\mathbf{x})$. Besides, we introduce the *signed valuation*, denoted by sval , which associates the couple $(\text{sign}(\mathbf{x}), \text{val}(\mathbf{x}))$ with a series $\mathbf{x} \in \mathbb{K}$. We denote by \mathbb{T}_{\pm} the image of \mathbb{K} by sval . We refer to it as the set of *signed tropical numbers*. For brevity, we denote an element of the form (ε, a) by a if $\varepsilon = 1$, $\ominus a$ if $\varepsilon = -1$, and $-\infty$ if $\varepsilon = 0$. Here, \ominus is a formal symbol. We call the elements of the first and second kind the *positive* and *negative* tropical numbers, respectively. We denote by \mathbb{T}_+ and \mathbb{T}_- the corresponding sets. In this way, (-2) is tropically positive, but $\ominus(-2)$ is tropically negative. Also, \mathbb{T} is embedded in \mathbb{T}_{\pm} , i.e., $\mathbb{T} = \mathbb{T}_+ \cup \{-\infty\}$. We shall extend the valuation maps val and sval to vectors and matrices in a coordinate-wise manner.

In \mathbb{T}_{\pm} , we define a *modulus* function, $|\cdot|: \mathbb{T}_{\pm} \rightarrow \mathbb{T}$, as $|\infty| = -\infty$ and $|a| = |\ominus a| = a$ for all $a \in \mathbb{T}_+$. We point out that \odot straightforwardly extends to \mathbb{T}_{\pm} using the standard rules for the sign, for instance $2 \odot (\ominus 3) = \ominus 5$. In contrast, we only partially define the tropical addition \oplus to elements of \mathbb{T}_{\pm} of identical sign, e.g., $2 \oplus 3 = 3$ and $(\ominus 2) \oplus (\ominus 3) = \ominus 3$. Such a partially defined operation will be enough for the present purposes. We note, however, that we may extend the addition operation so that it is everywhere defined, by embedding \mathbb{T}_{\pm} in the *symmetrized tropical semiring* [1], or by allowing the addition to be multivalued in the setting of hyperfields [10, 17, 41].

Moreover, we use the notion of tropical polynomials. A (*signed*) *tropical polynomial* over the variables X_1, \dots, X_n is a formal expression of the form

$$P(X) = \bigoplus_{\alpha \in \Lambda} a_{\alpha} \odot X_1^{\odot \alpha_1} \odot \cdots \odot X_n^{\odot \alpha_n}, \quad (4)$$

where Λ is a finite subset of $\{0, 1, 2, \dots\}^n$, and $a_\alpha \in \mathbb{T}_\pm \setminus \{-\infty\}$ for all $\alpha \in \Lambda$. We set $\Lambda^+ := \{\alpha \in \Lambda : a_\alpha \in \mathbb{T}_+\}$ and $\Lambda^- := \{\alpha \in \Lambda : a_\alpha \in \mathbb{T}_-\}$. We shall occasionally write $\Lambda(P)$ or $\Lambda^\pm(P)$ to emphasize the dependence in P . We say that the tropical polynomial P *vanishes* on the point $x \in \mathbb{T}_\pm^n$ if two of the terms $a_\alpha \odot x_1^{\odot \alpha_1} \odot \dots \odot x_n^{\odot \alpha_n}$ which have the greatest modulus do not have the same sign. If P does not vanish on x , we define $P(x)$ as the tropical sum of the terms which have the greatest modulus. As an example, if $P(X) = 2 \odot X_1^{\odot 3} \odot X_2^{\odot 4} \oplus (\ominus 0 \odot X_2)$, then $P(1, \ominus 5) = 25$, $P(1, -5) = \ominus(-5)$, whereas P vanishes on $(1, -5/3)$.

Furthermore, if Λ is empty, then we define $P(x) = -\infty$ for all $x \in \mathbb{T}_\pm^n$. The next definition and lemma relate the structure laws of \mathbb{T}_\pm and the ones of \mathbb{K} .

Definition 2.1 If

$$P(X) = \sum_{\alpha \in \Lambda} a_\alpha X_1^{\alpha_1} \dots X_n^{\alpha_n} \in \mathbb{K}[X_1, \dots, X_n] \quad (5)$$

is a polynomial over Puiseux series, then we define its *formal tropicalization*, denoted $\text{trop}(P)$, as the tropical polynomial

$$\text{trop}(P) := \bigoplus_{\alpha \in \Lambda} \text{sval}(a_\alpha) \odot X_1^{\odot \alpha_1} \odot \dots \odot X_n^{\odot \alpha_n}.$$

Lemma 2.2 Let $P(X) \in \mathbb{K}[X_1, \dots, X_n]$ and let $P := \text{trop}(P)$. Then, for all $x \in \mathbb{K}^n$, $\text{sval}(P(x)) = P(\text{sval}(x))$ provided that P does not vanish on $\text{sval}(x)$.

Proof Let P be as in (5). If P is a zero polynomial, then the claim is trivial. Otherwise, let $x := \text{sval}(x)$ and $a_\alpha := \text{sval}(a_\alpha)$ for all $\alpha \in \Lambda$. Note that for every $\alpha \in \Lambda$ we have the equalities

$$\begin{aligned} \text{sval}(a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n}) &= a_\alpha \odot x_1^{\odot \alpha_1} \odot \dots \odot x_n^{\odot \alpha_n}, \\ \text{val}(a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n}) &= |a_\alpha \odot x_1^{\odot \alpha_1} \odot \dots \odot x_n^{\odot \alpha_n}|. \end{aligned}$$

Let $\bar{\Lambda} \subset \Lambda$ denote terms that maximize the modulus of $a_\alpha \odot x_1^{\odot \alpha_1} \odot \dots \odot x_n^{\odot \alpha_n}$. Write $P(x)$ as

$$P(x) = \left(\sum_{\alpha \in \bar{\Lambda}} a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n} \right) + \left(\sum_{\alpha \notin \bar{\Lambda}} a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n} \right).$$

Since P does not vanish on $\text{sval}(x)$, the terms $a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n}$ such that $\alpha \in \bar{\Lambda}$ have the same sign. Furthermore, note that $\text{sval}(x + y) = \text{sval}(x) \oplus \text{sval}(y)$ provided that x and y have the same sign. In particular, we have the equality

$$\text{sval}\left(\sum_{\alpha \in \bar{\Lambda}} a_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n}\right) = \bigoplus_{\alpha \in \bar{\Lambda}} a_\alpha \odot x_1^{\odot \alpha_1} \odot \dots \odot x_n^{\odot \alpha_n} = P(x).$$

Moreover, observe that $\text{sval}(\mathbf{x} + \mathbf{y}) = \text{sval}(\mathbf{x})$ whenever $\text{val}(\mathbf{y}) < \text{val}(\mathbf{x})$. Hence, we get

$$\text{sval}(\mathbf{P}(\mathbf{x})) = \text{sval}\left(\sum_{\alpha \in \bar{\Lambda}} a_{\alpha} x_1^{\alpha_1} \dots x_n^{\alpha_n}\right) = \mathbf{P}(\mathbf{x}). \quad \square$$

Given a polynomial \mathbf{P} as in (5), we denote by \mathbf{P}^+ the polynomial obtained by summing the terms $a_{\alpha} X_1^{\alpha_1} \dots X_n^{\alpha_n}$ such that $a_{\alpha} > 0$. Similarly, \mathbf{P}^- refers to the sum of the terms $-a_{\alpha} X_1^{\alpha_1} \dots X_n^{\alpha_n}$ verifying $a_{\alpha} < 0$. In this way, $\mathbf{P} = \mathbf{P}^+ - \mathbf{P}^-$. We also use the analogues of these polynomials in the tropical setting. If P is the tropical polynomial given in (4), we define P^+ (resp. P^-) as the tropical polynomial obtained by summing the terms $|a_{\alpha}| \odot X_1^{\odot \alpha_1} \odot \dots \odot X_n^{\odot \alpha_n}$ where $a_{\alpha} \in \mathbb{T}_+$ (resp. \mathbb{T}_-). Observe that the quantities $P^+(\mathbf{x})$ and $P^-(\mathbf{x})$ are well defined for all $\mathbf{x} \in \mathbb{T}^n$, since the tropical polynomials P^+ and P^- only involve tropically positive coefficients. In particular, if \mathbf{P} is as in (5), and if $P = \text{trop}(\mathbf{P})$, observe that $P^+ = \text{trop}(\mathbf{P}^+)$ and $P^- = \text{trop}(\mathbf{P}^-)$.

Throughout the paper, we denote the set $\{1, \dots, k\}$ by $[k]$.

2.3 Tropical Polynomial Inequalities and Polyhedral Complexes

Given a tropical polynomial P as in (4), we say that P is *nonzero* if the set Λ is nonempty. For every such tropical polynomial and every point $\mathbf{x} \in \mathbb{R}^n$ we define the *set of maximizing multi-indices at \mathbf{x}* as

$$\text{Argmax}(P, \mathbf{x}) := \{\alpha \in \Lambda : \forall \beta \in \Lambda, |a_{\alpha}| + \langle \alpha, \mathbf{x} \rangle \geq |a_{\beta}| + \langle \beta, \mathbf{x} \rangle\},$$

where $\langle \cdot, \cdot \rangle$ refers to the usual scalar product. If P is a nonzero tropical polynomial and we fix a multi-index $\alpha \in \Lambda$, then the set

$$\mathcal{W} = \text{cl}(\{\mathbf{x} \in \mathbb{R}^n : \text{Argmax}(P, \mathbf{x}) = \alpha\})$$

is a polyhedron that is either empty or full-dimensional. (Here and in the sequel, $\text{cl}(\cdot)$ refers to the closure of a subset of \mathbb{R}^n with respect to the Euclidean topology.) Moreover, the family of these polyhedra, together with their faces, forms a polyhedral complex whose support is equal to \mathbb{R}^n . More precisely, a polyhedron \mathcal{V} is a (possibly empty) cell of this complex if and only if there exists a subset $L \subset \Lambda$ such that

$$\mathcal{V} = \text{cl}(\{\mathbf{x} \in \mathbb{R}^n : \text{Argmax}(P, \mathbf{x}) = L\}). \quad (6)$$

We denote this complex by $\mathcal{C}(P)$. The union of all $(n - 1)$ -dimensional polyhedra belonging to $\mathcal{C}(P)$ is called a *tropical hypersurface*. In other words, a tropical hypersurface is the set of all points $\mathbf{x} \in \mathbb{R}^n$ such that $\text{Argmax}(P, \mathbf{x})$ has at least two elements.

For the purpose of this work, given a nonzero tropical polynomial P , it is also convenient to consider the set

$$\mathcal{S}^{\geq}(P) := \{x \in \mathbb{R}^n : P^+(x) \geq P^-(x)\}.$$

To describe this set, we consider the family $\mathcal{C}^{\geq}(P)$ of positive cells of $\mathcal{C}(P)$. We say that a cell $\mathcal{V} \in \mathcal{C}(P)$ as in (6) is *positive* if there exists at least one $\alpha \in L$ such that $a_\alpha \in \mathbb{T}_+$ or if \mathcal{V} is empty. The family $\mathcal{C}^{\geq}(P)$ is a polyhedral complex whose support is equal to $\mathcal{S}^{\geq}(P)$.

Given a system of nonzero tropical polynomials P_1, \dots, P_m , one can regard the refinements of complexes defined by P_1, \dots, P_m . More precisely, we define $\mathcal{C}(P_1, \dots, P_m)$ and $\mathcal{C}^{\geq}(P_1, \dots, P_m)$ as

$$\begin{aligned}\mathcal{C}(P_1, \dots, P_m) &= \{\bigcap_{i=1}^m \mathcal{W}_i : \forall i, \mathcal{W}_i \in \mathcal{C}(P_i)\}, \\ \mathcal{C}^{\geq}(P_1, \dots, P_m) &= \{\bigcap_{i=1}^m \mathcal{W}_i : \forall i, \mathcal{W}_i \in \mathcal{C}^{\geq}(P_i)\}.\end{aligned}$$

The families $\mathcal{C}(P_1, \dots, P_m)$ and $\mathcal{C}^{\geq}(P_1, \dots, P_m)$ are polyhedral complexes. The support of the former is equal to \mathbb{R}^n while the support of the latter coincides with

$$\mathcal{S}^{\geq}(P_1, \dots, P_m) := \{x \in \mathbb{R}^n : \forall i, P_i^+(x) \geq P_i^-(x)\}.$$

Finally, in this work we consider polyhedral complexes with regular supports. Recall that a closed set $S \subset \mathbb{R}^n$ is called *regular* if $S = \text{cl}(\text{int}(S))$ (here and in the sequel, $\text{int}(\cdot)$ denotes the interior of a subset of \mathbb{R}^n). If \mathcal{C} is a polyhedral complex, then its support is regular if and only if \mathcal{C} is pure and full-dimensional. A basic property of such complexes appears in the next lemma.

Lemma 2.3 *Suppose that the polyhedral complex $\mathcal{C}^{\geq}(P_1, \dots, P_m)$ has a regular support. Then this support, $\mathcal{S}^{\geq}(P_1, \dots, P_m)$, coincides with the closure of the set*

$$\mathcal{S}^>(P_1, \dots, P_m) := \{x \in \mathbb{R}^n : \forall i, P_i^+(x) > P_i^-(x)\}.$$

Proof The set $\mathcal{S}^{\geq}(P_1, \dots, P_m)$ is closed since the tropical polynomial functions P_i^{\pm} are continuous, and obviously, $\mathcal{S}^>(P_1, \dots, P_m) \subset \mathcal{S}^{\geq}(P_1, \dots, P_m)$. Therefore,

$$\text{cl}(\mathcal{S}^>(P_1, \dots, P_m)) \subset \mathcal{S}^{\geq}(P_1, \dots, P_m).$$

Consider now $y \in \mathcal{S}^{\geq}(P_1, \dots, P_m)$. Since this set is regular, y belongs to a full-dimensional cell \mathcal{W} of $\mathcal{C}^{\geq}(P_1, \dots, P_m)$. We have $\mathcal{W} = \bigcap_{1 \leq i \leq m} \mathcal{W}_i$, where \mathcal{W}_i is a full-dimensional cell of $\mathcal{C}^{\geq}(P_i)$. This implies that $\mathcal{W}_i = \text{cl}(\{x \in \mathbb{R}^n : \text{Argmax}(P_i, x) = L_i\})$ where L_i is a one element subset of $\Lambda^+(P_i)$. We conclude that $P_i^+(x) > P_i^-(x)$ holds for all $x \in \text{int}(\mathcal{W}_i)$, and so, $\text{int}(\mathcal{W}) \subset \mathcal{S}^>(P_1, \dots, P_m)$. Taking any \bar{y} in the interior of \mathcal{W} , we see that the half-open segment $[\bar{y}, y[$ is contained in the interior of \mathcal{W} , and thus in $\mathcal{S}^>(P_1, \dots, P_m)$. It follows that $y \in \text{cl}(\mathcal{S}^>(P_1, \dots, P_m))$. \square

2.4 Valued Fields

In this section, we recall some basic information about valued fields. We refer to [23, Chap. 2] for a complete account. If \mathcal{K} is a field and Γ is an ordered abelian group, then a surjective function $\text{val}: \mathcal{K} \rightarrow \Gamma \cup \{-\infty\}$ is called a *valuation* if it fulfills the following three conditions:

$$\begin{aligned} \text{val}(x) = -\infty &\iff x = 0, \\ \forall x_1, x_2 \in \mathcal{K}, \text{val}(x_1 x_2) &= \text{val}(x_1) + \text{val}(x_2), \\ \forall x_1, x_2 \in \mathcal{K}, \text{val}(x_1 + x_2) &\leq \max(\text{val}(x_1), \text{val}(x_2)). \end{aligned} \quad (7)$$

A tuple $(\mathcal{K}, \Gamma, \text{val})$ is called a *valued field*. Under these conditions, $\mathcal{O} := \{x \in \mathcal{K} : \text{val}(x) \leq 0\}$ is a subring of \mathcal{K} and $\mathcal{M} := \{x \in \mathcal{K} : \text{val}(x) < 0\}$ is its maximal ideal. The quotient field $\mathcal{K} := \mathcal{O}/\mathcal{M}$ is called the *residue field*. We denote by res the canonical projection from \mathcal{O} to \mathcal{K} . The valuation is called *trivial* if $\Gamma = \{0\}$. Otherwise, it is called *nontrivial*. Recall that, as explained in Sect. 2.2, we use the “max-plus” sign-convention to define the valuation, i.e., the map $-\text{val}$ is a valuation in the sense of [23].

A map $\text{csec}: \Gamma \rightarrow \mathcal{K}^*$ is called a *cross-section* if it is a multiplicative morphism such that $\text{val} \circ \text{csec}$ is the identity map. A map $\text{ac}: \mathcal{K} \rightarrow \mathcal{K}$ is called an *angular component* if it fulfills the following conditions:

- $\text{ac}(0) = 0$;
- ac is a multiplicative morphism from \mathcal{K}^* to \mathcal{K}^* ;
- the function from \mathcal{O} to \mathcal{K} , mapping x to $\text{ac}(x)$ if $\text{val}(x) = 0$, and to 0 otherwise, is a surjective morphism of rings whose kernel is equal to \mathcal{M} .

Not every valued field admits an angular component [32]. Nevertheless, if it admits a cross-section csec , then $\text{ac}(x) := \text{res}(\text{csec}(-\text{val}(x))x)$ for $x \neq 0$ defines an angular component. For example, if \mathbb{K} is a field of Puiseux series defined in Sect. 2.1, then $\text{csec}(y) = t^y$ is a cross-section, and lc is an angular component. In fact, every real closed valued field has a cross-section, as shown by the following lemma.

Lemma 2.4 *Suppose that \mathcal{K} is real closed. Then $(\mathcal{K}, \Gamma, \text{val})$ admits a cross-section. (In particular, it has an angular component.)*

Proof The case where the valuation is trivial is obtained by taking a cross-section equal to 1. Therefore, we assume that the valuation is nontrivial. First, observe that in this case Γ is a divisible group, as any positive element of \mathcal{K} admits an n th root for every nonzero natural number n . Moreover, since Γ is ordered, it is also torsion free. Therefore, given a nonzero natural number n and $y \in \Gamma$, the equation $nz = y$ has a unique solution z in Γ . It follows that we can regard Γ as a vector space over \mathbb{Q} . Let $\{y_i\}_{i \in I}$ be a basis of this space. For every i take $x_i \in \mathcal{K}$ such that $x_i > 0$ and $\text{val}(x_i) = y_i$. For every finite subset $J \subset I$ and every $(\alpha_j) \in \mathbb{Q}^J$ define

$$\text{csec}\left(\sum_{j \in J} \alpha_j y_j\right) = \prod_{j \in J} x_j^{\alpha_j}.$$

It is obvious that csec is a cross-section. \square

Finally, we recall the notion of a convex valuation. Suppose that \mathcal{K} is an ordered field with a total order \geq . We say that the valuation val is *convex with respect to* \geq if it satisfies the following property: for every $x_1 \in \mathcal{O}$ and every $x_2 \in \mathcal{K}$ we have the implication

$$0 \leq x_2 \leq x_1 \implies x_2 \in \mathcal{O}.$$

If \mathcal{K} is a real closed field, it has a unique total order. In this case, the convexity property is understood in the sense of this order. It can be shown that if \mathcal{K} is real closed and val is convex, then \mathcal{K} is also real closed (see [23, Thm. 4.3.7]). The field of Puiseux series is an example of a real closed field with convex valuation.

3 Semilinearity of Tropical Semialgebraic Sets

The goal of this section is to prove the following theorem.

Theorem 3.1 *Let \mathcal{K} be a real closed field equipped with a nontrivial and convex valuation val . Furthermore, suppose that the set $\mathcal{S} \subset \mathcal{K}^n$ is semialgebraic. Then every stratum of $\text{val}(\mathcal{S})$ is semilinear.*

Let us detail the notions used in this statement. If \mathcal{K} is a real closed field, then we say that a subset $\mathcal{S} \subset \mathcal{K}^n$ is *basic semialgebraic* if it is of the form

$$\mathcal{S} = \{x \in \mathcal{K}^n : \forall i = 1, \dots, p, P_i(x) > 0 \wedge \forall i = p+1, \dots, q, P_i(x) = 0\},$$

where $P_i \in \mathcal{K}[X_1, \dots, X_n]$ are polynomials. We say that \mathcal{S} is *semialgebraic* if it is a finite union of basic semialgebraic sets. Similarly, if Γ is a divisible ordered abelian group, then we say that a set $\mathcal{S} \subset \Gamma^n$ is *basic semilinear* if it is of the form

$$\mathcal{S} = \{g \in \Gamma^n : \forall i = 1, \dots, p, f_i(g) > h^{(i)}, \forall i = p+1, \dots, q, f_i(g) = h^{(i)}\},$$

where $f_i \in \mathbb{Z}[X_1, \dots, X_n]$ are homogeneous linear polynomials with integer coefficients and $h^{(i)} \in \Gamma$. We say that \mathcal{S} is *semilinear* if it is a finite union of basic semilinear sets.

Finally, since we are interested in valuations of semialgebraic sets defined in valued fields, we work with $\Gamma \cup \{-\infty\}$ rather than Γ . Any set $S \subset (\Gamma \cup \{-\infty\})^n$ is naturally stratified as follows: the *support* of a point $x \in (\Gamma \cup \{-\infty\})^n$ is defined as the set of indices $k \in [n]$ such that $x_k \neq -\infty$. Given a nonempty subset $K \subset [n]$, and a set $S \subset (\Gamma \cup \{-\infty\})^n$, we define the *stratum of S associated with K* as the subset of $\Gamma^{|K|}$ formed by the projection $(x_k)_{k \in K}$ of the points $x \in S$ with support K .

The rest of Sect. 3 is devoted to the presentation of the proof of Theorem 3.1, which relies on model theoretic results in valued fields. After a preliminary section on model theory (Sect. 3.1), we explain how Theorem 3.1 is obtained from a quantifier elimination technique in valued fields of Denef and Pas (Sect. 3.2).

3.1 Languages and Structures

In this section we recall some basic notions from model theory. We refer to [28, Chap. 1] and [38, Chap. 1] for more information. In model theory, a *language* \mathcal{L} is a collection of *symbols* that are divided into three sets: a set of *constant symbols*, a set of *function symbols*, and a set of *relation symbols*. For example, $\mathcal{L}_{\text{og}} := (0, +, \leq)$ is the language of ordered groups, while $\mathcal{L}_{\text{or}} := (0, 1, +, -, \cdot, \leq)$ is the language of ordered rings.

An \mathcal{L} -*structure* is a tuple $\mathcal{M} := (M, \mathcal{L})$, where M is a nonempty set (called a *domain*) and every symbol of \mathcal{L} can be interpreted in M . For instance, if $\Gamma = (\Gamma, 0, +, \leq)$ is an ordered abelian group, then we can interpret the symbol 0 as zero in Γ , the symbol $+$ as addition, and the symbol \leq as order in Γ . Thus, every ordered abelian group is an \mathcal{L}_{og} -structure.

The formalism introduced above enables us to study the first-order formulas over \mathcal{L} (or \mathcal{L} -*formulas*). The atoms of these formulas are constructed by applying relation symbols to terms built out of variables, and functions and constants from \mathcal{L} .

Given an \mathcal{L} -formula ψ and a variable x , an occurrence of x is said to be *bound* if it is located within the scope of a subformula of the form $\forall x \dots$ or $\exists x \dots$. Other occurrences of the variable x are said to be *free*. By extension, the variable x is said to be *free* when it occurs freely in the formula ψ . Up to renaming some of the variables, we can suppose that free variables do not have bound occurrences.

We often denote an \mathcal{L} -formula ψ as $\psi(X)$, where $X = (x_1, \dots, x_n)$ is a string of free variables that occur in ψ . If \mathcal{M} is an \mathcal{L} -structure with domain M and we fix a vector $\bar{X} \in M^n$, then $\psi(\bar{X})$ can be interpreted as a meaningful statement about \mathcal{M} . This statement can be either true or false. For example, if we fix an ordered abelian group Γ , then the \mathcal{L}_{og} -formula $\forall x_1(x_1 \geq 0 \rightarrow \exists x_2(x_2 \geq 0 \wedge x_1 = x_2 + x_2))$ has no free variables. It is interpreted in Γ as “for every nonnegative element $x_1 \in \Gamma$, there exists a nonnegative element $x_2 \in \Gamma$ such that x_1 is equal to x_2 added to x_2 .” Note that this is true if we take $\Gamma = (\mathbb{Q}, +, \leq)$, but false if we take $\Gamma = (\mathbb{Z}, +, \leq)$. Similarly, the \mathcal{L}_{og} -formula $\exists x_2(x_1 = x_2 + x_2)$ has one free variable x_1 . If we take $\Gamma = (\mathbb{Z}, +, \leq)$, then $\psi(2)$ is true, but $\psi(1)$ is false. We denote “ $\psi(\bar{X})$ is true in \mathcal{M} ” as $\mathcal{M} \models \psi(\bar{X})$. A formula without free variables is called a *sentence*. A set $\mathcal{S} \subset M^n$ is called *definable* (in \mathcal{L}) if there exists a number $m \geq 0$, a vector $\bar{b} \in M^m$, and an \mathcal{L} -formula $\psi(x_1, \dots, x_{n+m})$ such that

$$\mathcal{S} = \{x \in M^n : \mathcal{M} \models \psi(x_1, \dots, x_n, \bar{b})\}.$$

Example 3.2 Take an \mathcal{L}_{og} -structure $\mathcal{M} = (\Gamma, 0, +, \leq)$, where Γ is a divisible ordered abelian group. Suppose that $\psi(x_1, \dots, x_{n+m})$ is a quantifier-free \mathcal{L}_{og} -formula (i.e., a formula that does not contain quantifier symbols). Then $\mathcal{S} = \{x \in \Gamma^n : \mathcal{M} \models \psi(x_1, \dots, x_n, \bar{b})\}$ is a semilinear set. Conversely, every semilinear set can be written in such form.

If \mathcal{L} is a language, then any set of \mathcal{L} -sentences is called a *theory*. In our context, one can think that a theory is a set of axioms. If Th is a fixed theory in \mathcal{L} , then we say that an \mathcal{L} -structure \mathcal{M} is a *model* of Th when we have $\mathcal{M} \models \psi$ for every $\psi \in \text{Th}$. Furthermore,

if ψ is an \mathcal{L} -sentence that does not necessarily belong to Th , then we say that ψ is a *logical consequence* of Th , if ψ is true in every model of Th . We say that \mathcal{L} -formulas $\psi(X)$, $\phi(X)$ are *equivalent* in Th if the sentence $\forall x_1 \dots \forall x_n \psi(X) \leftrightarrow \phi(X)$ is a logical consequence of Th . We say that the theory Th admits *quantifier elimination* if every \mathcal{L} -formula is equivalent in Th to a quantifier-free formula. Finally, we say that a theory Th is *complete* if for every \mathcal{L} -sentence ψ , either ψ or $\neg\psi$ is a logical consequence of Th .

Example 3.3 The theory of real closed fields, denoted Th_{rcf} , is a theory in the language of ordered rings \mathcal{L}_{or} . It consists of the usual axioms of ordered fields, the axiom $\forall x_1 (x_1 \geq 0 \rightarrow \exists x_2 (x_1 = x_2 \cdot x_2))$ which governs the existence of square roots, and an infinite set of axioms that state the fact that every polynomial of an odd degree has a root. In other words, for every $n \geq 1$, Th_{rcf} contains the axiom $\forall x_0 \dots \forall x_{2n} \exists x (x^{2n+1} + x_{2n}x^{2n} + \dots + x_1x + x_0 = 0)$. A classical result due to Tarski states that this theory admits quantifier elimination and is complete (see [28, Thm. 3.3.15 and Cor. 3.3.16]). As an immediate corollary one sees that if \mathcal{K} is a real closed field, then a set $\mathcal{S} \subset \mathcal{K}^n$ is definable in \mathcal{L}_{or} if and only if it is semialgebraic.

In the next section, we use divisible ordered abelian groups which arise as value groups of nonarchimedean real closed fields. Since the valuation map may evaluate to $-\infty$, we need to deal with divisible ordered abelian groups with bottom element. In more detail, we denote by $\mathcal{L}_{\text{ogb}} := (0, -\infty, +, \leq)$ the language of ordered groups with bottom element. The theory of nontrivial divisible ordered abelian groups with bottom element, denoted Th_{dogb} , consists of the axioms of divisible ordered abelian groups, the nontriviality axiom $\exists y (y \neq 0 \wedge y \neq -\infty)$, and the axioms that extend the addition and order to $-\infty$, namely $\forall y (-\infty + y = -\infty)$ and $\forall y (y \geq -\infty)$. As stated in the next proposition, this theory admits quantifier elimination and is complete. It follows from the fact that the same result holds in the case of groups without bottom element [28, Cor. 3.1.17].

Proposition 3.4 *The theory Th_{dogb} admits quantifier elimination and is complete. Moreover, any \mathcal{L}_{ogb} -formula $\theta(Y)$ with $Y = (y_1, \dots, y_m)$ and $m \geq 1$ is equivalent to a quantifier-free formula of the form*

$$\bigvee_{\Sigma \subseteq [m]} ((\forall \sigma \in \Sigma, y_\sigma \neq -\infty) \wedge (\forall \sigma \notin \Sigma, y_\sigma = -\infty) \wedge \psi_\Sigma),$$

where every ψ_Σ is a quantifier-free \mathcal{L}_{og} -formula over a subset of variables in $\{y_\sigma\}_{\sigma \in \Sigma}$.

This proposition can be easily proven from [28, Cor. 3.1.17] using double induction over m and the length of θ . We omit the proof for brevity. We emphasize that every ψ_Σ is a \mathcal{L}_{og} -formula, i.e., a formula that does not contain the symbol $-\infty$. As a consequence of Proposition 3.4 and the discussion in Example 3.2, we get the following characterization of definable sets.

Corollary 3.5 *Suppose that Γ is a nontrivial divisible abelian group. Then $\mathcal{S} \subset (\Gamma \cup \{-\infty\})^n$ is definable in \mathcal{L}_{ogb} if and only if every stratum of \mathcal{S} is semilinear.*

3.2 Quantifier Elimination in Real Closed Valued Fields

In this section, we want to show quantifier elimination over real closed fields equipped with a nontrivial and convex valuation. We suppose that \mathcal{K} is a real closed field and $\text{val}: \mathcal{K} \rightarrow \Gamma \cup \{-\infty\}$ is a valuation that is nontrivial and convex. We denote by \mathcal{K} the residue field of $(\mathcal{K}, \Gamma, \text{val})$, and by ac we denote any angular component of this field. Under these conditions, Γ is divisible and \mathcal{K} is real closed, as noted in Sect. 2.4. In order to describe such structures, we consider the following *three-sorted language*

$$\mathcal{L}_{\text{rcvf}} := (\mathcal{L}_{\mathcal{K}}, \mathcal{L}_{\Gamma}, \mathcal{L}_{\mathcal{K}}, \text{val}, \text{ac}).$$

Here, $\mathcal{L}_{\mathcal{K}}$ and $\mathcal{L}_{\mathcal{K}}$ denote the language of ordered rings (respectively associated with \mathcal{K} and \mathcal{K}), \mathcal{L}_{Γ} denotes the language of ordered groups with bottom element, val is a symbol for valuation map, and ac is a symbol for angular component. In the language $\mathcal{L}_{\text{rcvf}}$, any formula has three kinds of variables, one kind for every sort. Let x_1, x_2, \dots denote the variables associated with \mathcal{K} , y_1, y_2, \dots denote the variables associated with $\Gamma \cup \{-\infty\}$, and z_1, z_2, \dots denote the variables associated with \mathcal{K} . If θ is an $\mathcal{L}_{\text{rcvf}}$ -formula, then we denote it as $\theta(X, Y, Z)$, where X, Y, Z are sequences of free variables associated with $\mathcal{K}, \Gamma \cup \{-\infty\}, \mathcal{K}$ respectively. The constant, function, and relation symbols of the language $\mathcal{L}_{\text{rcvf}}$ are implicitly typed. For instance, the addition symbol of $\mathcal{L}_{\mathcal{K}}$ takes two elements of the sort \mathcal{K} , and returns an element of the same sort. The symbol val yields an element of the sort Γ from an element of the sort \mathcal{K} . Then, $\mathcal{L}_{\text{rcvf}}$ -formulas are built from the symbols of the language $\mathcal{L}_{\text{rcvf}}$ and variables in such a way that every term and atom is well typed. We refer to [38, Chap. 1] for a formal treatment of multisorted languages.

Let us denote by Th_{rcvf} the theory of valued fields with angular component which are real closed and have a nontrivial and convex valuation. More precisely, the theory consists of the axioms of the theory of real closed fields for \mathcal{K} , the axioms of the theory of ordered abelian groups with bottom element for $\Gamma \cup \{-\infty\}$, the axioms of ordered fields for \mathcal{K} , the axioms specifying that val is a nontrivial and convex valuation, and the axioms specifying that ac is an angular component. Note that the latter axioms imply that \mathcal{K} is indeed the residue field of $(\mathcal{K}, \Gamma, \text{val})$. In the next theorem, we show that this theory admits quantifier elimination. The cornerstone of the proof is a result due to Pas [30,31], which establishes that the theory of henselian valued fields with angular component admits elimination of quantifiers over the \mathcal{K} -variables. We refer to [16] for more recent generalizations of Pas's result.

Theorem 3.6 *The theory Th_{rcvf} admits quantifier elimination and is complete. Moreover, any $\mathcal{L}_{\text{rcvf}}$ -formula $\theta(X, Y, Z)$ is equivalent in Th_{rcvf} to a formula of the form*

$$\bigvee_{i=1}^m \left(\phi_i(\text{val}(f_{i1}(X)), \dots, \text{val}(f_{ik_i}(X)), Y) \right. \\ \left. \wedge \psi_i(\text{ac}(f_{i(k_i+1)}(X)), \dots, \text{ac}(f_{il_i}(X)), Z) \right),$$

where, for every $i = 1, \dots, m$, $f_{i1}, \dots, f_{il_i} \in \mathbb{Z}[X]$ are polynomials with integer coefficients, ϕ_i is a quantifier-free \mathcal{L}_{Γ} -formula, and ψ_i is a quantifier-free $\mathcal{L}_{\mathcal{K}}$ -formula.

Proof Let $\theta(X, Y, Z)$ denote any $\mathcal{L}_{\text{rcvf}}$ -formula. Recall that the order \leq in any real closed field can be defined as $x_1 \leq x_2 \iff \exists x_3 (x_2 - x_1 = x_3^2)$. This enables us to inductively eliminate all occurrences of the symbols \leq of the languages $\mathcal{L}_{\mathcal{K}}$ and $\mathcal{L}_{\mathcal{K}}$. Therefore, $\theta(X, Y, Z)$ is equivalent in Th_{rcvf} to a formula $\hat{\theta}(X, Y, Z)$ without the symbol \leq . Moreover, by [23, Thm. 4.3.7], $(\mathcal{K}, \Gamma, \text{val})$ is henselian. This enables us to apply the quantifier elimination of Pas [30, Thm. 4.1]. (To be more precise, we use the formulation given in [16, Thm. 4.2].) As a result, $\hat{\theta}(X, Y, Z)$ is equivalent in Th_{rcvf} to a formula of the form

$$\bigvee_{i=1}^m \left(\phi_i(\text{val}(f_{i1}(X)), \dots, \text{val}(f_{ik_i}(X)), Y) \right. \\ \left. \wedge \psi_i(\text{ac}(f_{i(k_i+1)}(X)), \dots, \text{ac}(f_{il_i}(X)), Z) \right), \quad (8)$$

where, for every $i = 1, \dots, m$, $f_{i1}, \dots, f_{il_i} \in \mathbb{Z}[X]$ are polynomials with integer coefficients, ϕ_i is an \mathcal{L}_{Γ} -formula, and ψ_i is an $\mathcal{L}_{\mathcal{K}}$ -formula. Then, we apply Proposition 3.4 and [28, Thm. 3.3.15] to eliminate the quantifiers in the formulas ϕ_i and ψ_i . This shows the last part of the statement.

In the case where θ is a sentence, the formulas ϕ_i and ψ_i are also sentences. The completeness results in Proposition 3.4 and [28, Cor. 3.3.16] applied to each subformula ϕ_i and ψ_i in (8) allow to prove that either θ or $\neg\theta$ is a logical consequence of Th_{rcvf} . \square

As a corollary, we obtain Theorem 3.1.

Proof of Theorem 3.1 Let Γ denote the value group of \mathcal{K} and \mathcal{K} denote the residue field. The structure $\mathcal{M} = (\mathcal{K}, \Gamma \cup \{-\infty\}, \mathcal{K}, \mathcal{L}_{\text{rcvf}})$ is a model of Th_{rcvf} . Let $\phi(x_1, \dots, x_{n+m})$ be an $\mathcal{L}_{\mathcal{K}}$ -formula and $\bar{b} \in \mathcal{K}^m$ be a vector such that $\mathcal{S} = \{x \in \mathcal{K}^n : \mathcal{K} \models \phi(x, \bar{b})\}$. Take the formula $\theta(x_{n+1}, \dots, x_{n+m}, y_1, \dots, y_n)$ in $\mathcal{L}_{\text{rcvf}}$ defined as

$$\exists x_1 \dots \exists x_n \left(\phi(x_1, \dots, x_{n+m}) \wedge \text{val}(x_1) = y_1 \wedge \dots \wedge \text{val}(x_n) = y_n \right).$$

We obviously have

$$\text{val}(\mathcal{S}) = \{y \in (\Gamma \cup \{-\infty\})^n : \mathcal{M} \models \theta(\bar{b}, y)\}.$$

By Theorem 3.6, θ is equivalent to a formula of the form

$$\bigvee_{i=1}^m \left(\phi_i(\text{val}(f_{i1}(X)), \dots, \text{val}(f_{ik_i}(X)), Y) \right. \\ \left. \wedge \psi_i(\text{ac}(f_{i(k_i+1)}(X)), \dots, \text{ac}(f_{il_i}(X))) \right),$$

where we denote $X := (x_{n+1}, \dots, x_{n+m})$, $Y := (y_1, \dots, y_n)$, every ϕ_i is an \mathcal{L}_{Γ} -formula, every ψ_i is an $\mathcal{L}_{\mathcal{K}}$ -formula, and f_{i1}, \dots, f_{il_i} are polynomials with integer coefficients. If we fix X to be equal to \bar{b} , then this formula is equivalent to a formula of the form

$$\bigvee_{i \in I} \phi_i(\xi_{i1}, \dots, \xi_{ik_i}, Y),$$

where I is a subset of $[m]$ and we denote $\text{val}(f_{ik}(\bar{b})) = \xi_{ik} \in \Gamma$. Hence, $\text{val}(\mathcal{S})$ is definable in \mathcal{L}_Γ . By Corollary 3.5, $\text{val}(\mathcal{S})$ has semilinear strata. \square

4 Closedness of Tropical Semialgebraic Sets

In this section we strengthen Theorem 3.1 by showing that the strata of $\text{val}(\mathcal{S})$ are not only semilinear but also closed. More precisely, we show the following theorem.

Theorem 4.1 *Let \mathcal{K} be a real closed field equipped with a nontrivial and convex valuation val . Suppose that set $\mathcal{S} \subset \mathcal{K}^n$ is semialgebraic. Then every stratum of $\text{val}(\mathcal{S})$ is closed in the product topology of the order topology of the value group Γ .*

Furthermore, if \mathcal{S} is closed, then $\text{val}(\mathcal{S})$ is closed in the product topology of the order topology of $\Gamma \cup \{-\infty\}$.

Remark 4.2 Theorems 3.1 and 4.1 should be compared with Theorems 3.11 and 4.10 of Alessandrini [2], dealing with log-limits of sets that are definable in certain o-minimal polynomially bounded structures. The fact that we obtain the present results using quantifier elimination has two advantages. First, it gives a constructive proof, while the proofs in [2] rely on compactness arguments. Second, our methods apply to arbitrary real closed valued fields, while the analysis of [2] is restricted to fields whose value group is a subgroup of \mathbb{R} . On the other hand, the results of [2] allow to go beyond the semialgebraic setting; they apply for instance to the o-minimal structure of real numbers with restricted analytic functions.

Remark 4.3 We point out that every subset of $(\Gamma \cup \{-\infty\})^n$ that is closed in the product topology of the order topology of $\Gamma \cup \{-\infty\}$ has closed strata, but the converse is not true. For example, the set $\mathbb{R}^2 \cup \{-\infty\}$ has closed strata, but is not closed in the product topology of the order topology of \mathbb{T}^2 .

To prove Theorem 4.1, we first consider the case of Puiseux series, $\mathcal{K} = \mathbb{K}$. The proof needs a few auxiliary lemmas. Hereafter, $\mathbb{K}_{>0}^n := \{\mathbf{x} \in \mathbb{K} : \forall k, x_k > 0\}$ denotes the open positive orthant of \mathbb{K}^n . Let us fix a basic semialgebraic set $\mathcal{S} \subset \mathbb{K}_{>0}^n$ defined as

$$\mathcal{S} := \{\mathbf{x} \in \mathbb{K}_{>0}^n : \forall i = 1, \dots, p, P_i(\mathbf{x}) > 0 \wedge \forall i = p+1, \dots, q, Q_i(\mathbf{x}) = 0\} \quad (9)$$

for some polynomials $P_1, \dots, P_p, Q_{p+1}, \dots, Q_q \in \mathbb{K}[X_1, \dots, X_n]$. Equivalently, we put \mathcal{S} under the form

$$\mathcal{S} = \{\mathbf{x} \in \mathbb{K}_{>0}^n : \forall i = 1, \dots, p, P_i(\mathbf{x}) > 0 \wedge \forall i = p+1, \dots, q, P_i(\mathbf{x}) \geq 0\}, \quad (10)$$

where we set $P_i := -Q_i^2$ for all $i = p+1, \dots, q$. Denote $P_i := \text{trop}(P_i)$ for all $i = 1, \dots, q$. In the next lemma, we highlight a property of the full-dimensional cells of the complex $\mathcal{C}(P_1, \dots, P_q)$ whose interior is contained in $\text{val}(\mathcal{S})$.

Lemma 4.4 Suppose that \mathcal{W} is a full-dimensional cell of $\mathcal{C}(P_1, \dots, P_q)$ such that $\text{int}(\mathcal{W}) \cap \text{val}(\mathcal{S}) \neq \emptyset$. Let $w \in \text{int}(\mathcal{W})$, and $\mathbf{w} \in \text{val}^{-1}(w) \cap \mathbb{K}_{>0}^n$ be an arbitrary lift. Then $\mathbf{w} \in \mathcal{S}$.

Proof Take a point $z \in \mathcal{S}$ such that $z := \text{val}(z) \in \text{int}(\mathcal{W})$. For every $i = 1, \dots, q$ we have $P_i^-(z) \geq 0$. By Lemma 2.2 and the fact val is order preserving, we obtain $P_i^+(z) \geq P_i^-(z)$. Since \mathcal{W} is a full-dimensional cell of $\mathcal{C}(P_1, \dots, P_q)$, we have the equality $\text{int}(\mathcal{W}) = \bigcap_{i=1}^q \text{int}(\mathcal{W}_i)$, where, for every i , \mathcal{W}_i is a full-dimensional cell of $\mathcal{C}(P_i)$. In particular, $\text{Argmax}(P_i, z)$ has only one element and we have $P_i^+(z) > P_i^-(z)$. Furthermore, we have $\text{Argmax}(P_i, z) = \text{Argmax}(P_i, w)$ for any point $w \in \text{int}(\mathcal{W})$. This implies that $P_i^+(w) > P_i^-(w)$. Therefore, if $\mathbf{w} \in \text{val}^{-1}(w) \cap \mathbb{K}_{>0}^n$ is an arbitrary lift of w , then by Lemma 2.2 we have $\text{val}(P_i^+(\mathbf{w})) > \text{val}(P_i^-(\mathbf{w}))$ and hence $\mathbf{w} \in \mathcal{S}$. \square

Lemma 4.5 Let $A \in \mathbb{Q}^{m \times n}$ be any matrix. Define a function $f: \mathbb{K}_{>0}^n \rightarrow \mathbb{K}_{>0}^m$ as

$$f(\mathbf{x})_i := \mathbf{x}_1^{A_{i1}} \mathbf{x}_2^{A_{i2}} \dots \mathbf{x}_n^{A_{in}}.$$

Let $\mathcal{S} \subset \mathbb{K}_{>0}^n$ be any semialgebraic set. Then $f(\mathcal{S}) \subset \mathbb{K}_{>0}^m$ is semialgebraic and we have $\text{val}(f(\mathcal{S})) = A(\text{val}(\mathcal{S}))$.

Proof The first claim follows from the fact that the class of semialgebraic sets is closed under semialgebraic transformations [12, Prop. 2.83]. The second claim follows from the identity $\text{val}(f(\mathbf{x})_i) = A_i(\text{val}(\mathbf{x}))$. \square

Lemma 4.6 Suppose that $\mathcal{S} \subset \mathbb{K}_{>0}^n$ is a semialgebraic set. Then $\text{val}(\mathcal{S}) \subset \mathbb{R}^n$ is a union of finitely many closed polyhedra.

Proof We proceed by induction over the dimension n . First, suppose that $n = 1$. Since \mathbb{K} is a real closed field, every semialgebraic set in \mathbb{K} is a finite union of points and open intervals. Observe that the image by the valuation of an open interval in $\mathbb{K}_{>0}$ is an interval that is closed in \mathbb{R} . Therefore, the claim is true for $n = 1$.

Second, suppose that the claim holds in dimension $n - 1$. Observe that it is enough to prove the claim for basic semialgebraic sets. Fix a basic semialgebraic set $\mathcal{S} \subset \mathbb{K}_{>0}^n$ as in (10) and take the polyhedral complex $\mathcal{C} := \mathcal{C}(P_1, \dots, P_q)$. Let $\tilde{\mathcal{W}}_1, \dots, \tilde{\mathcal{W}}_r$ denote the cells of \mathcal{C} . By Lemma 3.1, $\text{val}(\mathcal{S})$ is a finite union of relatively open polyhedra. Denote these polyhedra by $\text{ri}(\tilde{\mathcal{V}}_1), \dots, \text{ri}(\tilde{\mathcal{V}}_s)$, where each $\tilde{\mathcal{V}}_j$ is a closed polyhedron and ri denotes the relative interior. For every (i, j) , let \mathcal{W}_{ij} be a polyhedron such that

$$\text{ri}(\mathcal{W}_{ij}) = \text{ri}(\tilde{\mathcal{W}}_i) \cap \text{ri}(\tilde{\mathcal{V}}_j).$$

Observe that $\text{val}(\mathcal{S})$ is a union of $\text{ri}(\mathcal{W}_{ij})$. We consider an element w^* of $\text{cl}(\text{val}(\mathcal{S}))$. Let us look at two cases.

Case I: There is a full-dimensional polyhedron \mathcal{W}_{ij} such that $w^* \in \mathcal{W}_{ij}$. In this case, let $\mathcal{H} = \{w \in \mathbb{R}^n : \langle a, w \rangle = \langle a, w^* \rangle\}$ be any hyperplane intersecting the

interior of \mathcal{W}_{ij} , and such that $a \in \mathbb{Q}^n$. Consider $w^{(1)}, w^{(2)}, \dots$ a sequence such that $w^{(h)} \in \mathcal{H} \cap \text{int}(\mathcal{W}_{ij})$ for all h and $w^{(h)} \rightarrow w^*$. Take the set $Y \subset \mathbb{K}_{>0}^n$ defined as

$$Y = \mathcal{S} \cap \left\{ x \in \mathbb{K}_{>0}^n : \prod_{k \in [n]} x_k^{a_k} = t^{\langle a, w^* \rangle} \right\}.$$

For every h define $w^{(h)} \in \text{val}^{-1}(w^{(h)}) \cap \mathbb{K}_{>0}^n$ as $w_k^{(h)} = t^{w_k^{(h)}}$ for all $k \in [n]$. Note that every $w^{(h)}$ belongs to the interior of the full-dimensional polyhedron $\tilde{\mathcal{W}}_i$. Consequently, $w^{(h)}$ belongs to Y by Lemma 4.4. Take $l \in [n]$ such that $a_l \neq 0$ and let $\pi: \mathbb{K}_{>0}^n \rightarrow \mathbb{K}_{>0}^{n-1}$ denote the projection that forgets the l -th coordinate. Similarly, let $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ denote the projection that forgets the l -th coordinate. By the induction hypothesis and Lemma 4.5, $\text{val}(\pi(Y))$ is a closed subset of \mathbb{R}^{n-1} and we have $\text{val}(\pi(Y)) = \pi(\text{val}(Y))$. The sequence $\pi(w^{(h)})$ converges to $\pi(w^*)$. Therefore, we have $\pi(w^*) \in \pi(\text{val}(Y))$. In other words, there exists a point $w^* \in Y$ such that $\pi(\text{val}(w^*)) = \pi(w^*)$. Moreover, we have $\text{val}(w^*) \in \mathcal{H}$ and $w^* \in \mathcal{H}$. Since $a_l \neq 0$, this implies that $\text{val}(w^*) = w^*$. Therefore $w^* \in \text{val}(\mathcal{S})$.

Case II: If w^* does not belong to any full-dimensional polyhedron \mathcal{W}_{ij} , then we denote by I the set of all indices (i, j) such that \mathcal{W}_{ij} contains w^* . We can take $\rho > 0$ so small that the closed Chebyshev ball $B(w^*, \rho)$ does not intersect any polyhedron \mathcal{W}_{ij} with $(i, j) \notin I$. Let $w^{(1)}, w^{(2)}, \dots$ be a convergent sequence of elements of \mathbb{R}^n , $w^{(h)} \rightarrow w^*$ such that $w^{(h)} \in \text{val}(\mathcal{S})$ for all h . Every polyhedron \mathcal{W}_{ij} such that $(i, j) \in I$ is not full-dimensional. Therefore, it is included in an affine hyperplane \mathcal{H}_{ij} . Let $X = \bigcup_{(i,j) \in I} \mathcal{H}_{ij}$ be a union of these hyperplanes. Observe that we have $w^* \in X$ and that $\text{val}(\mathcal{S}) \cap B(w^*, \rho) \subset X$. Let $v \in \mathbb{Q}^n$ be any rational vector such that $v \notin (X - w^*)$. (Here, by $X - w^*$ we mean the translation of X by vector $-w^*$.) Note that the affine line $w^* + \text{span}(v)$ intersects X only in w^* .

Let $A \in \mathbb{Q}^{(n-1) \times n}$ be a rational matrix such that $\ker(A) = \text{span}(v)$. Take the function $f: \mathbb{K}_{>0}^n \rightarrow \mathbb{K}_{>0}^{n-1}$ defined as

$$(f(x))_k := \prod_{l=1}^n x_l^{A_{kl}}, \quad k = 1, 2, \dots, n-1.$$

Let $U := \{x \in \mathbb{K}_{>0}^n : \forall l, x_l \in [t^{w_l^* - \rho}, t^{w_l^* + \rho}]\}$. By Lemma 4.5, the set $f(\mathcal{S} \cap U) \subset \mathbb{K}_{>0}^{n-1}$ is $\text{val}(f(\mathcal{S} \cap U)) = A(\text{val}(\mathcal{S} \cap U))$. Therefore, by the induction hypothesis, the set $A(\text{val}(\mathcal{S} \cap U))$ is closed. For every $w^{(h)}$, let $w^{(h)} \in \mathcal{S}$ denote any element of \mathcal{S} such that $\text{val}(w^{(h)}) = w^{(h)}$. For h large enough we have $w^{(h)} \in B(w^*, \rho/2)$ and hence $w^{(h)} \in \mathcal{S} \cap U$. Moreover, the sequence $Aw^{(h)}$ converges to Aw^* . Since $A(\text{val}(\mathcal{S} \cap U))$ is closed, there is $w^* \in \mathcal{S} \cap U$ such that $Aw^* = A\text{val}(w^*)$. As $w^* \in U$, we have $\text{val}(w^*) \in B(w^*, \rho)$. Therefore

$$\text{val}(w^*) \in \left((w^* + \text{span}(v)) \cap B(w^*, \rho) \cap \text{val}(\mathcal{S}) \right).$$

On the other hand, we have $\text{val}(\mathcal{S}) \cap B(w^*, \rho) \subset X$ and $(w^* + \text{span}(v)) \cap X = w^*$. Hence $\text{val}(w^*) = w^*$ and $w^* \in \text{val}(\mathcal{S})$. \square

Lemma 4.7 Suppose that $\mathcal{S} \subset \mathbb{K}^n$ is a nonempty bounded closed semialgebraic set. Let $K \subset [n]$ be a set of indices such that for every $\mathbf{a} \in \mathbb{K}_{>0}$ the set $\{\mathbf{x} \in \mathcal{S} : \forall k \in K, x_k \in [0, \mathbf{a}]\}$ is nonempty. Then there exists a point $\mathbf{y} \in \mathcal{S}$ such that $y_k = 0$ for all $k \in K$.

Proof We prove that the statement holds for any real closed field \mathcal{K} . Fix an \mathcal{L}_{or} -formula $\psi(x_1, \dots, x_{n+m})$. For every vector $\bar{b} \in \mathcal{K}^m$ we define the semialgebraic set $\mathcal{S}_{\bar{b}}$ by

$$\mathcal{S}_{\bar{b}} := \{x \in \mathcal{K}^n : \mathcal{K} \models \psi(x_1, \dots, x_n, \bar{b})\}.$$

The statement “for all $(x_{n+1}, \dots, x_{n+m})$, if the set $\mathcal{S}_{(x_{n+1}, \dots, x_{n+m})}$ is nonempty, bounded, closed, and the set $\{x \in \mathcal{S}_{(x_{n+1}, \dots, x_{n+m})} : \forall k \in K, x_k \in [0, a]\}$ is nonempty for every $a > 0$, then there exists a point $y \in \mathcal{S}_{(x_{n+1}, \dots, x_{n+m})}$ such that $y_k = 0$ for all $k \in K$ ” is a sentence in the language of ordered rings \mathcal{L}_{or} . It is true in \mathbb{R} , hence it is true in \mathcal{K} by the model completeness of real closed fields [28, Cor. 3.3.16]. \square

The lemmas above lead to the main theorem of this section.

Proof of Theorem 4.1 We first prove the result for a semialgebraic set \mathcal{S} included in the closed positive orthant $\mathbb{K}_{\geq 0}^n$. Let $K \subset [n]$ be any nonempty subset and let $X_K \subset \mathbb{K}^n$ be the set defined as

$$X_K := \{\mathbf{x} \in \mathbb{K}^n : \mathbf{x}_k \neq 0 \iff k \in K\}.$$

The sets X_K and subsequently $\mathcal{S} \cap X_K$ are semialgebraic. Let $\pi : \mathbb{K}^n \rightarrow \mathbb{K}^K$ denote the projection on the coordinates from K . Similarly, let $\pi : \mathbb{T}^n \rightarrow \mathbb{T}^K$ denote the projection on the coordinates from K . Observe that the stratum of $\text{val}(\mathcal{S})$ associated with K is equal to $\pi(\text{val}(\mathcal{S} \cap X_K)) = \text{val}(\pi(\mathcal{S} \cap X_K))$. Moreover, the set $\pi(\mathcal{S} \cap X_K)$ is included in $\mathbb{K}_{>0}^K$. Therefore the first claim follows from Lemma 4.6.

To prove the second claim, suppose that \mathcal{S} is closed. Let $x \in \mathbb{T}^n$ be any point that does not belong to $\text{val}(\mathcal{S})$. Let $K \subset [n]$ be the support of x . For any $M, N > 0$ we denote $I_k(M, N) = [-\infty, -M]$ if $k \notin K$ and $I_k(M, N) =]x_k - \frac{1}{N}, x_k + \frac{1}{N}[$ otherwise. Similarly, we denote $\mathbf{I}_k(M, N) = [0, t^{-M+1}] \subset \mathbb{K}_{\geq 0}$ for $k \notin K$ and $\mathbf{I}_k(M, N) = [t^{x_k - \frac{2}{N}}, t^{x_k + \frac{2}{N}}] \subset \mathbb{K}_{>0}$ otherwise. We want to show that there is an open neighborhood of x that does not belong to $\text{val}(\mathcal{S})$. Suppose that this is not the case. Then, for any $M, N > 0$, the set $\prod_{k=1}^n I_k(M, N)$ contains a point from $\text{val}(\mathcal{S})$. Therefore, the set $\mathcal{S}^{(M, N)} := \mathcal{S} \cap \prod_{k=1}^n \mathbf{I}_k(M, N)$ is nonempty, on top of being closed and bounded. If we fix $N > 0$, then, by Lemma 4.7, there is a point $\mathbf{y}^{(N)} \in \mathcal{S}^{(1, N)}$ such that $y_k^{(N)} = 0$ for all $k \notin K$. In other words, the set $\pi(\mathcal{S} \cap X_K)$ contains a point that belongs to $\prod_{k \in K} [t^{x_k - \frac{2}{N}}, t^{x_k + \frac{2}{N}}]$. Hence, the stratum of $\text{val}(\mathcal{S})$ associated with K contains a point that belongs to $\prod_{k \in K} [x_k - \frac{2}{N}, x_k + \frac{2}{N}]$. Since this is true for all $N > 0$, and the strata of $\text{val}(\mathcal{S})$ are closed, we have $x \in \text{val}(\mathcal{S})$, which gives a contradiction.

Second, suppose that $\mathcal{S} \subset \mathbb{K}^n$ is any semialgebraic set. Given a vector $\delta \in \{+1, -1\}^n$, we denote by f_δ the involution which maps $\mathbf{x} \in \mathbb{K}^n$ to the vector with

entries $\delta_k \mathbf{x}_k$. With this notation, \mathcal{S} is the union of the sets of the form $\mathcal{S} \cap f_\delta(\mathbb{K}_{\geq 0}^n)$. Moreover, the set $f_\delta(\mathcal{S} \cap f_\delta(\mathbb{K}_{\geq 0}^n))$ is a semialgebraic set included in $\mathbb{K}_{\geq 0}^n$, and its image under the valuation map coincides with that of $\mathcal{S} \cap f_\delta(\mathbb{K}_{\geq 0}^n)$. The claim follows by applying the result of the previous paragraph to each of the sets $f_\delta(\mathcal{S} \cap f_\delta(\mathbb{K}_{\geq 0}^n))$.

To prove the claim for an arbitrary field \mathcal{K} we use Theorem 3.6. We fix an \mathcal{L}_{or} -formula $\psi(x_1, \dots, x_{n+m})$. For every vector $\bar{b} \in \mathcal{K}^m$ we can look at the semialgebraic set

$$\mathcal{S}_{\bar{b}} := \{x \in \mathcal{K}^n : \mathcal{K} \models \psi(x_1, \dots, x_n, \bar{b})\}.$$

The statement “for all $(x_{n+1}, \dots, x_{n+m})$, the image by valuation of the set $\mathcal{S}_{(x_{n+1}, \dots, x_{n+m})}$ has closed strata” is a sentence in $\mathcal{L}_{\text{rcvf}}$. It is true in \mathbb{K} and hence, by the completeness result of Theorem 3.6, it is also true in \mathcal{K} . The same is true for the statement “for all $(x_{n+1}, \dots, x_{n+m})$, if $\mathcal{S}_{(x_{n+1}, \dots, x_{n+m})}$ is closed, then its image by valuation is closed.” \square

As a by-product, we get the following result, which generalizes the proposition of Develin and Yu [20, Prop. 2.9] on polyhedra to basic semialgebraic sets.

Corollary 4.8 *Suppose that $\mathcal{S} \subset \mathbb{K}_{>0}^n$ is a semialgebraic set defined as*

$$\mathcal{S} := \{\mathbf{x} \in \mathbb{K}_{>0}^n : P_1(\mathbf{x}) \square_1 0, \dots, P_m(\mathbf{x}) \square_m 0\},$$

where $P_i \in \mathbb{K}[X_1, \dots, X_n]$ are nonzero polynomials and $\square \in \{\geq, >\}^m$. Let $P_i := \text{trop}(P_i)$ for all i and suppose that $\mathcal{C}^{\geq}(P_1, \dots, P_m)$ has regular support. Then

$$\text{val}(\mathcal{S}) = \{x \in \mathbb{R}^n : \forall i, P_i^+(x) \geq P_i^-(x)\}.$$

Proof Denote $\mathcal{S}^{\geq} := \{x \in \mathbb{R}^n : \forall i, P_i^+(x) \geq P_i^-(x)\}$ and suppose that $\mathbf{x} \in \mathcal{S}$. Since val is order preserving, we have $\text{val}(\mathbf{x}) \in \mathcal{S}^{\geq}$ by Lemma 2.2. Therefore $\text{val}(\mathcal{S}) \subset \mathcal{S}^{\geq}$. On the other hand, if we take any point x such that $P_i^+(x) > P_i^-(x)$ for all i , then any lift $\mathbf{x} \in \text{val}^{-1}(x) \cap \mathbb{K}_{>0}^n$ belongs to \mathcal{S} . Hence, we have the inclusion

$$\{x \in \mathbb{R}^n : \forall i, P_i^+(x) > P_i^-(x)\} \subset \text{val}(\mathcal{S}) \subset \{x \in \mathbb{R}^n : \forall i, P_i^+(x) \geq P_i^-(x)\}$$

and the claim follows from Lemma 2.3 and Theorem 4.1. \square

Example 4.9 Take $P = 0 \oplus (X_1^{\odot 2} \odot X_2^{\odot 2}) \oplus (2 \odot X_1 \odot X_2) \oplus (\ominus 2 \odot X_1^{\odot 2}) \oplus (\ominus 2 \odot X_2^{\odot 2})$. Then $\mathcal{C}^{\geq}(P)$ is depicted in Fig. 1. This support of this complex is not regular and Corollary 4.8 does not apply. Indeed, take $P(\mathbf{x}_1, \mathbf{x}_2) = 1 + \mathbf{x}_1^2 \mathbf{x}_2^2 + t^2 \mathbf{x}_1 \mathbf{x}_2 - t^2 \mathbf{x}_1^2 - t^2 \mathbf{x}_2^2$. We have $\text{trop}(P) = P$, but the set $\text{val}(\{(\mathbf{x}_1, \mathbf{x}_2) \in \mathbb{K}_{>0}^2 : P(\mathbf{x}_1, \mathbf{x}_2) \geq 0\})$ does not contain the open segment $](-1, -1), (1, 1)[$.

Remark 4.10 The result of Corollary 4.8 may be thought of as a semialgebraic analogue of Kapranov’s theorem [22, Thm. 2.1.1]. This theorem characterizes an image by valuation of a complex hypersurface. A difference is that Kapranov’s theorem

orthants, defined in the following way. If $\mu \in \{-1, +1\}^n$, then the μ -orthant of \mathbb{T}_{\pm}^n is defined as

$$\mathbb{T}_{\pm, \mu}^n := \{x \in \mathbb{T}_{\pm}^n : \forall k, \text{sign}(x_k) \in \{0, \mu_k\}\}.$$

We note that the orthants are closed in the topology of \mathbb{T}_{\pm}^n . Moreover, they are homeomorphic to \mathbb{T}^n , with the homeomorphism given by the absolute value map

$$\mathbb{T}_{\pm, \mu} \ni (x_1, \dots, x_n) \mapsto (|x_1|, \dots, |x_n|) \in \mathbb{T}^n. \quad (11)$$

We can also divide \mathbb{T}_{\pm}^n into 3^n strata, in the following way. If $\sigma \in \{-1, 0, +1\}^n$ is a vector of signs, then we define the σ -stratum of \mathbb{T}_{\pm}^n as

$$\mathbb{T}_{\pm}^{n, (\sigma)} := \{x \in \mathbb{T}_{\pm}^n : \forall k \in [n], \text{sign}(x_k) = \sigma_k\}. \quad (12)$$

We point out that in Sect. 3 we defined “stratum” as a set of points with only finite coordinates. However, in the present context it is more convenient to consider σ -strata as sets of points with some coordinates equal to $-\infty$, as in (12). We note that the strata of \mathbb{T}_{\pm}^n are generally *not* closed in the topology of \mathbb{T}_{\pm}^n . More precisely, if $\sigma \neq 0$ and $\sum_k |\sigma_k| = p$, then $\mathbb{T}_{\pm}^{n, (\sigma)}$ is homeomorphic to \mathbb{T}_+^p and the homeomorphism is given by the *sign-forgetting projection* $\pi^{(\sigma)} : \mathbb{T}_{\pm}^{n, (\sigma)} \rightarrow \mathbb{T}_+^p$ defined as $\pi^{(\sigma)}(x_1, \dots, x_n) := (|x_{k_1}|, \dots, |x_{k_p}|)$, where $\{k_1, \dots, k_p\}$ are the indices such that $\sigma_{k_i} \neq 0$.

The following result about the topology of $\text{sval}(\mathcal{S})$ follows from Theorem 4.1, compare with [26, Thm. 6.5].

Corollary 4.11 *If $\mathcal{S} \subset \mathbb{K}^n$ is a semialgebraic set, then $\text{sval}(\mathcal{S}) \cap \mathbb{T}_{\pm}^{n, (\sigma)}$ is closed in the induced topology of $\mathbb{T}_{\pm}^{n, (\sigma)}$ for every $\sigma \in \{-1, 0, +1\}^n$. Moreover, if \mathcal{S} is closed, then $\text{sval}(\mathcal{S})$ is closed in the topology of \mathbb{T}_{\pm}^n .*

Proof To prove the first part, fix $\mathcal{S} \subset \mathbb{K}^n$ and $\sigma \in \{-1, 0, +1\}^n$. If $\sigma = 0$, then the claim is trivial. Otherwise, consider the semialgebraic set $\mathcal{S}_{\sigma} := \{x \in \mathcal{S} : \forall k, \text{sign}(x_k) = \sigma_k\}$ and note that $\text{sval}(\mathcal{S}) \cap \mathbb{T}_{\pm}^{n, (\sigma)} = \text{sval}(\mathcal{S}_{\sigma})$. Denote $p := \sum_k |\sigma_k|$ and $K := \{k \in [n] : \sigma_k \neq 0\}$. Let $\mathcal{S}' := \text{val}(\mathcal{S}_{\sigma}) \subset \mathbb{T}^n$ and let $\mathcal{S}'_K \subset \mathbb{T}_+^p$ denote the stratum of \mathcal{S}' associated with K . By Theorem 4.1, the set \mathcal{S}'_K is closed in the topology of \mathbb{T}_+^p . Therefore, the claim follows from the fact that $\text{sval}(\mathcal{S}_{\sigma}) = (\pi^{(\sigma)})^{-1}(\mathcal{S}'_K)$, where $\pi^{(\sigma)} : \mathbb{T}_{\pm}^{n, (\sigma)} \rightarrow \mathbb{T}_+^p$ is the sign-forgetting projection. To prove the second part, fix a vector $\mu \in \{-1, +1\}^n$, let

$$\mathbb{K}_{\mu}^n := \{x \in \mathbb{K}^n : \forall k, \text{sign}(x_k) \in \{0, \mu_k\}\}$$

denote the corresponding orthant of the Puiseux series, and let $\mathcal{S}_{\mu} := \mathcal{S} \cap \mathbb{K}_{\mu}^n$. By Theorem 4.1, the set $\text{val}(\mathcal{S}_{\mu})$ is closed in the topology of \mathbb{T}^n . Moreover, $\text{sval}(\mathcal{S}_{\mu}) \subset \mathbb{T}_{\pm, \mu}^n$ is the preimage of $\text{val}(\mathcal{S}_{\mu})$ under the absolute value map given in (11). Hence, $\text{sval}(\mathcal{S}_{\mu})$ is a closed subset of $\mathbb{T}_{\pm, \mu}^n$ and therefore it is also a closed subset of \mathbb{T}_{\pm}^n . The claim follows from the fact that $\text{sval}(\mathcal{S}) = \bigcup_{\mu \in \{-1, +1\}^n} \text{sval}(\mathcal{S}_{\mu})$. \square

5 Tropical Spectrahedra

5.1 Tropicalization of Nonarchimedean Spectrahedra

We now introduce the notion of tropical spectrahedra.

Definition 5.1 A set $\mathcal{S} \subset \mathbb{T}^n$ is said to be a *tropical spectrahedron* if there exists a spectrahedron $\mathcal{S} \subset \mathbb{K}_{\geq 0}^n$ such that $\mathcal{S} = \text{val}(\mathcal{S})$.

If $\mathcal{S} = \text{val}(\mathcal{S})$, then we refer to \mathcal{S} as the *tropicalization* of the spectrahedron \mathcal{S} , and \mathcal{S} is said to be a *lift (over the field \mathbb{K})* of \mathcal{S} .

Recall that we have the following characterization of positive semidefinite matrices:

Proposition 5.2 A symmetric matrix $A \in \mathbb{K}^{m \times m}$ is positive semidefinite if and only if every principal minor of A is nonnegative.

Given symmetric matrices $Q^{(0)}, \dots, Q^{(n)} \in \mathbb{K}^{m \times m}$ and $x \in \mathbb{K}^n$, we denote by $Q(x)$ the matrix pencil $Q^{(0)} + x_1 Q^{(1)} + \dots + x_n Q^{(n)}$. Proposition 5.2 provides a description of the spectrahedron $\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^n : Q(x) \succeq 0\}$ by a system of polynomial inequalities of the form $\det Q_{I \times I}(x) \geq 0$, where I is a nonempty subset of $[m]$, and $\det Q_{I \times I}(x)$ corresponds to the $(I \times I)$ -minor of the matrix $Q(x)$. Following this, we obtain that the tropical spectrahedron \mathcal{S} is included in the intersection of the sets $\{x \in \mathbb{T}^n : \text{trop}(P)^+(x) \geq \text{trop}(P)^-(x)\}$ where P is a polynomial of the form $\det Q_{I \times I}(x)$. In general, this inclusion may be strict. We refer to [3, Exam. 15] for an example in which \mathcal{S} is a polyhedron. Nevertheless, under the regularity assumption stated in Corollary 4.8, both sets coincide. In fact, we prove that, under similar assumptions, tropical spectrahedra have a description that is much simpler than the one provided by Corollary 4.8. This description only involves principal tropical minors of order 2.

Our results are divided into three parts. In Sect. 5.2 we deal with spectrahedra defined by Metzler matrices $Q^{(0)}, \dots, Q^{(n)}$ (i.e., matrices in which the off-diagonal entries are nonpositive). This enables us to use a lemma that is similar to Corollary 4.8 in order to give a description of tropical spectrahedra under a regularity assumption.

In Sect. 5.3 we switch to non-Metzler matrices. In this case, tropical spectrahedra may not be regular, even under strong genericity assumptions. Nevertheless, we are able to extend our previous analysis to this case and give a description, involving only principal minors of size at most 2, of non-Metzler spectrahedra, under a regularity assumption over some associated sets.

Finally, the purpose of Sect. 5.4 is to show that the regularity assumptions used in Sect. 5.2 and 5.3 hold generically.

Let us start with some introductory remarks. First, observe that in order to characterize the class of tropical spectrahedra, it is enough to restrict ourselves to tropical spectrahedral cones, as the image of a spectrahedron can be deduced from the image of its homogenized version. This is formally stated in the next lemma.

Lemma 5.3 Let $Q^{(0)}, \dots, Q^{(n)} \in \mathbb{K}^{m \times m}$ be a sequence of symmetric matrices. Define

$$\mathcal{S} := \left\{ x \in \mathbb{K}_{\geq 0}^n : Q^{(0)} + x_1 Q^{(1)} + \dots + x_n Q^{(n)} \succeq 0 \right\}$$

and

$$\mathcal{S}^h := \left\{ (x_0, \mathbf{x}) \in \mathbb{K}_{\geq 0}^{n+1} : x_0 \mathbf{Q}^{(0)} + x_1 \mathbf{Q}^{(1)} + \cdots + x_n \mathbf{Q}^{(n)} \succcurlyeq 0 \right\}.$$

Then

$$\text{val}(\mathcal{S}) = \pi(\{x \in \text{val}(\mathcal{S}^h) : x_0 = 0\}),$$

where $\pi : \mathbb{T}^{n+1} \rightarrow \mathbb{T}^n$ denotes the projection that forgets the first coordinate.

Proof We start by proving the inclusion \subset . Take any $x \in \text{val}(\mathcal{S})$ and its lift $\mathbf{x} \in \mathcal{S} \cap \text{val}^{-1}(x)$. Observe that the point $(1, \mathbf{x})$ belongs to \mathcal{S}^h . Therefore, the point $(0, x)$ belongs to $\text{val}(\mathcal{S}^h)$ and x belongs to $\pi(\{x \in \text{val}(\mathcal{S}^h) : x_0 = 0\})$. Conversely, let x belong to $\pi(\{x \in \text{val}(\mathcal{S}^h) : x_0 = 0\})$. Then $(0, x)$ belongs to $\text{val}(\mathcal{S}^h)$. In other words, there exists a lift $(z, \mathbf{x}) \in \mathcal{S}^h$ such that $\text{val}(z) = 0$ and $\text{val}(\mathbf{x}) = x$. Take the point $(1, \mathbf{x}/z)$. This point also belongs to \mathcal{S}^h . Moreover, \mathbf{x}/z belongs to \mathcal{S} . Hence, the point $x = \text{val}(\mathbf{x}/z)$ belongs to $\text{val}(\mathcal{S})$. \square

Second, let us explain our approach to the tropicalization of spectrahedra. It relies on the next elementary lemma. The latter should be compared with a result of Yu [43] which shows that the set of images by the valuation of the set of positive semidefinite matrices \mathbf{A} over the field of Puiseux series is determined by the inequalities $\text{val}(\mathbf{A}_{ii}) + \text{val}(\mathbf{A}_{jj}) \geq 2\text{val}(\mathbf{A}_{ij})$ for $i \neq j$.

Lemma 5.4 *Let $\mathbf{A} \in \mathbb{K}^{m \times m}$ be a symmetric matrix. Suppose that \mathbf{A} has nonnegative entries on its diagonal and that the inequality $\mathbf{A}_{ii}\mathbf{A}_{jj} \geq (m-1)^2\mathbf{A}_{ij}^2$ holds for all pairs (i, j) such that $i \neq j$. Then \mathbf{A} is positive semidefinite.*

Proof If \mathbf{A} is a zero matrix, then there is nothing to show. From now on we suppose that \mathbf{A} has at least one nonzero entry. First, let us suppose that \mathbf{A} has positive entries on its diagonal. In this case, let $\mathbf{B} \in \mathbb{K}^{m \times m}$ be the diagonal matrix defined by $\mathbf{B}_{ii} := \mathbf{A}_{ii}^{-1/2}$ for all i . Observe that \mathbf{A} is positive semidefinite if and only if the matrix $\mathbf{D} := \mathbf{B}\mathbf{A}\mathbf{B}$ is positive semidefinite. Moreover, \mathbf{D} has ones on its diagonal and $\mathbf{D}_{ij} = \mathbf{A}_{ii}^{-1/2}\mathbf{A}_{ij}\mathbf{A}_{jj}^{-1/2}$ for all $i \neq j$. Hence $|\mathbf{D}_{ij}| \leq 1/(m-1)$ for all $i \neq j$. Therefore \mathbf{D} is diagonally dominant and hence positive semidefinite.

Second, if \mathbf{A} has some zeros on its diagonal, let $I = \{i \in [m] : \mathbf{A}_{ii} \neq 0\}$. Since the inequality $\mathbf{A}_{ii}\mathbf{A}_{jj} \geq (m-1)^2\mathbf{A}_{ij}^2$ holds, we have $\mathbf{A}_{ij} = 0$ if either $i \notin I$ or $j \notin I$. Let \mathbf{A}_I denote the submatrix formed by the rows and columns with indices from I . Then \mathbf{A} is positive semidefinite if and only if \mathbf{A}_I is positive semidefinite. Finally, \mathbf{A}_I is positive semidefinite by the considerations from the previous paragraph. \square

Given a spectrahedron $\mathcal{S} = \{\mathbf{x} \in \mathbb{K}_{\geq 0}^n : \mathbf{Q}(\mathbf{x}) \succcurlyeq 0\}$ we define two sets $\mathcal{S}^{\text{out}}, \mathcal{S}^{\text{in}} \subset \mathbb{K}_{\geq 0}^n$ as

$$\begin{aligned} \mathcal{S}^{\text{out}} &:= \left\{ \mathbf{x} \in \mathbb{K}_{\geq 0}^n : \forall i, \mathbf{Q}_{ii}(\mathbf{x}) \geq 0, \forall i \neq j, \mathbf{Q}_{ii}(\mathbf{x})\mathbf{Q}_{jj}(\mathbf{x}) \geq (\mathbf{Q}_{ij}(\mathbf{x}))^2 \right\}, \\ \mathcal{S}^{\text{in}} &:= \left\{ \mathbf{x} \in \mathbb{K}_{\geq 0}^n : \forall i, \mathbf{Q}_{ii}(\mathbf{x}) \geq 0, \forall i \neq j, \mathbf{Q}_{ii}(\mathbf{x})\mathbf{Q}_{jj}(\mathbf{x}) \geq (m-1)^2(\mathbf{Q}_{ij}(\mathbf{x}))^2 \right\}. \end{aligned}$$

Proposition 5.2 shows that $\mathcal{S} \subset \mathcal{S}^{\text{out}}$, while Lemma 5.4 shows that $\mathcal{S}^{\text{in}} \subset \mathcal{S}$. In order to describe the set $\text{val}(\mathcal{S})$, we will exhibit conditions that ensure that the tropicalizations of \mathcal{S}^{in} and \mathcal{S}^{out} coincide, i.e., $\text{val}(\mathcal{S}^{\text{out}}) = \text{val}(\mathcal{S}) = \text{val}(\mathcal{S}^{\text{in}})$.

5.2 Tropical Metzler Spectrahedra

In this section, we study the spectrahedra defined by Metzler matrices. Recall that a square matrix $A \in \mathbb{K}^{m \times m}$ is a (*negated*) *Metzler matrix* if its off-diagonal coefficients are nonpositive. Similarly, we say that a matrix $M \in \mathbb{T}_{\pm}^{m \times m}$ is a *tropical Metzler matrix* if $M_{ij} \in \mathbb{T}_- \cup \{-\infty\}$ for all $i \neq j$. Let $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$ be symmetric tropical Metzler matrices. Given $i, j \in [m]$, we refer to $Q_{ij}(X)$ as the tropical polynomial:

$$Q_{ij}(X) := Q_{ij}^{(1)} \odot X_1 \oplus \dots \oplus Q_{ij}^{(n)} \odot X_n.$$

Definition 5.5 If $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$ are symmetric tropical Metzler matrices, we define the *tropical Metzler spectrahedron* $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$ described by $Q^{(1)}, \dots, Q^{(n)}$ as the set of points $x \in \mathbb{T}^n$ that fulfill the following two conditions:

- for all $i \in [m]$, $Q_{ii}^+(x) \geq Q_{ii}^-(x)$;
- for all $i, j \in [m]$, $i < j$, $Q_{ii}^+(x) \odot Q_{jj}^+(x) \geq (Q_{ij}(x))^{\odot 2}$.

Observe that the term $Q_{ij}(x)$ ($i \neq j$) is well defined for any $x \in \mathbb{T}^n$ thanks to the Metzler property of the matrices $Q^{(k)}$.

Where there is no ambiguity, we denote $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$ by \mathcal{S} . With standard notation, the constraints defining this set respectively read: for all $i \in [m]$,

$$\max_{Q_{ii}^{(k)} \in \mathbb{T}_+} (Q_{ii}^{(k)} + x_k) \geq \max_{Q_{ii}^{(l)} \in \mathbb{T}_-} (|Q_{ii}^{(l)}| + x_l), \quad (13)$$

and for all $i, j \in [m]$ such that $i < j$,

$$\max_{Q_{ii}^{(k)} \in \mathbb{T}_+} (Q_{ii}^{(k)} + x_k) + \max_{Q_{jj}^{(k')} \in \mathbb{T}_+} (Q_{jj}^{(k')} + x_{k'}) \geq 2 \max_{l \in [n]} (|Q_{ij}^{(l)}| + x_l). \quad (14)$$

Remark 5.6 The system of constraints (13) and (14) has an interpretation in terms of games. Indeed, it is shown in [6, 7] to be equivalent to an equation of the form $x \leq T(x)$, where T is the dynamic programming operator of a turn-based stochastic mean payoff game. The existence of a solution $x \in \mathbb{T}^n$, not identically $-\infty$, is equivalent to the existence of an initial state that is winning. See also [5, 36] for more information on the relation between tropical spectrahedra and turn-based games.

The next proposition justifies the terminology introduced in Definition 5.5, and ensures that the set \mathcal{S} is indeed a tropical spectrahedron. To this end, we explicitly construct a spectrahedron $\mathcal{S} \subset \mathbb{K}_{\geq 0}^n$ verifying $\text{val}(\mathcal{S}) = \mathcal{S}$. (This result was already announced in the conference version of the paper [6]. We provide here the full proof for the sake of completeness.)

Proposition 5.7 *The set $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$ is a tropical spectrahedron.*

Proof Let us define the matrices $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{K}^{m \times m}$ as follows:

- if $Q_{ij}^{(k)} \in \mathbb{T}_-$, then we set $Q_{ij}^{(k)} := -t^{|Q_{ij}^{(k)}|}$;
- if $Q_{ij}^{(k)} \in \mathbb{T}_+$ (which, under our assumptions, can happen only if $i = j$), then $Q_{ij}^{(k)} := mnt^{Q_{ij}^{(k)}}$;
- if $Q_{ij}^{(k)} = -\infty$, then $Q_{ij}^{(k)} := 0$.

Consider the spectrahedron $\mathcal{S} := \{x \in \mathbb{K}_{\geq 0}^n : Q(x) \succ 0\}$. We claim that $\text{val}(\mathcal{S}) = \mathcal{S}$.

We start with the inclusion $\text{val}(\mathcal{S}^{\text{out}}) \subset \mathcal{S}$. Let $x \in \mathcal{S}^{\text{out}}$. Observe that for all $i \neq j$, the inequality $Q_{ii}^+(x)Q_{jj}^+(x) \geq (Q_{ij}^+(x))^2$ holds thanks to the fact that $Q_{ii}(x) \geq 0$. Moreover, we have $\text{val}(Q_{ii}^+(x)) = Q_{ii}^+(x)$, where $x = \text{val}(x)$. Similarly, $\text{val}(Q_{ii}^-(x)) = Q_{ii}^-(x)$. As the $Q^{(k)}$ are tropical Metzler matrices, we have $\text{val}(Q_{ij}(x)) = |Q_{ij}(x)|$ for $i \neq j$. Since the map val is order preserving over $\mathbb{K}_{\geq 0}$, we deduce that $x \in \mathcal{S}$.

Now, let us prove the inclusion $\mathcal{S} \subset \text{val}(\mathcal{S}^{\text{in}})$. Take any $x \in \mathcal{S}$ and its lift $x_k = t^{x_k}$, with the convention that $t^{-\infty} = 0$. First, as noted in the previous paragraph, we have $\text{val}(Q_{ii}^+(x)) = Q_{ii}^+(x)$ and $\text{val}(Q_{ii}^-(x)) = Q_{ii}^-(x)$. We have chosen the matrices $Q^{(k)}$ and the point x in such a way that

$$Q_{ii}^+(x) = \sum_{Q_{ii}^{(k)} \in \mathbb{T}_+} mnt^{Q_{ii}^{(k)} + x_k} \geq mnt^{Q_{ii}^+(x)}. \quad (15)$$

Similarly, we have $Q_{ii}^-(x) = \sum_{Q_{ii}^{(k)} \in \mathbb{T}_-} t^{|Q_{ii}^{(k)}| + x_k} \leq nt^{Q_{ii}^-(x)}$. Since $Q_{ii}^+(x) \geq Q_{ii}^-(x)$, we deduce that $Q_{ii}^-(x) \leq \frac{1}{m} Q_{ii}^+(x)$, and so $Q_{ii}(x) \geq (1 - \frac{1}{m}) Q_{ii}^+(x) \geq 0$. Second, for all $i \neq j$ we have $0 \geq Q_{ij}(x) \geq -nt^{|Q_{ij}(x)|}$. Using (15) and the fact that $Q_{ii}^+(x) \odot Q_{jj}^+(x) \geq (Q_{ij}(x))^{\odot 2}$, we obtain

$$Q_{ij}(x)^2 \leq \frac{1}{m^2} Q_{ii}^+(x) Q_{jj}^+(x).$$

Therefore, by the previous inequalities,

$$\begin{aligned} & Q_{ii}(x) Q_{jj}(x) - (m-1)^2 Q_{ij}(x)^2 \\ & \geq \left(1 - \frac{1}{m}\right)^2 Q_{ii}^+(x) Q_{jj}^+(x) - (m-1)^2 Q_{ij}(x)^2 \geq 0. \end{aligned}$$

Hence $x \in \mathcal{S}^{\text{in}}$. Therefore, by Proposition 5.2 and Lemma 5.4 we have $\text{val}(\mathcal{S}) \subset \text{val}(\mathcal{S}^{\text{out}}) \subset \mathcal{S} \subset \text{val}(\mathcal{S}^{\text{in}}) \subset \text{val}(\mathcal{S})$, which implies that $\text{val}(\mathcal{S}) = \mathcal{S}$. \square

Example 5.8 If $A^{(1)}, \dots, A^{(p)}$ are matrices, then $\text{tdiag}(A^{(1)}, \dots, A^{(p)})$ refers to the block diagonal matrix with blocks $A^{(s)}$ on the diagonal and all other entries equal to




Fig. 2 A tropical Metzler spectrahedron

$-\infty$. Let $Q^{(0)}, Q^{(1)}, Q^{(2)} \in \mathbb{T}_{\pm}^{9 \times 9}$ be symmetric tropical Metzler matrices defined as follows:

$$\begin{aligned} Q^{(0)} &:= \text{tdiag} \left(8, \ominus 1, \ominus 1, \begin{bmatrix} -\infty & \ominus 3 \\ \ominus 3 & -\infty \end{bmatrix}, \begin{bmatrix} 2 & -\infty \\ -\infty & 8 \end{bmatrix}, \begin{bmatrix} 3 & -\infty \\ -\infty & 9 \end{bmatrix} \right), \\ Q^{(1)} &:= \text{tdiag} \left(\ominus 0, 0, -\infty, \begin{bmatrix} 0 & -\infty \\ -\infty & -2 \end{bmatrix}, \begin{bmatrix} -\infty & \ominus 0 \\ \ominus 0 & -\infty \end{bmatrix}, \begin{bmatrix} 0 & -\infty \\ -\infty & 4 \end{bmatrix} \right), \\ Q^{(2)} &:= \text{tdiag} \left(\ominus 0, -\infty, 0, \begin{bmatrix} -1 & -\infty \\ -\infty & -1 \end{bmatrix}, \begin{bmatrix} 0 & -\infty \\ -\infty & 4 \end{bmatrix}, \begin{bmatrix} -\infty & \ominus 0 \\ \ominus 0 & -\infty \end{bmatrix} \right). \end{aligned}$$

The intersection of the tropical Metzler spectrahedron $\mathcal{S}(Q^{(0)}, Q^{(1)}, Q^{(2)})$ with the hyperplane $\{x_0 = 0\}$ is depicted in Fig. 2.

We now focus on the main problem of characterizing the image by the valuation of a spectrahedron defined by Metzler matrices. Our goal is to show that any spectrahedron $\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^n : Q(x) \succcurlyeq 0\}$ verifying $\text{sval}(Q^{(k)}) = Q^{(k)}$ is mapped to the tropical Metzler spectrahedron \mathcal{S} , provided that some assumptions related to the genericity of the matrices $Q^{(k)}$ and the regularity of the set \mathcal{S} hold. To do so, we prove a weaker result, Theorem 5.12, on the tropicalization of the spectrahedron restricted to the open positive orthant $\mathbb{K}_{>0}^n$.

Lemma 5.9 *Let $A \in \mathbb{T}^{m \times m}$ be a symmetric matrix such that $A_{ii} \in \mathbb{T}_+ \cup \{-\infty\}$ for all i and $A_{ii} \odot A_{jj} > A_{ij}^{\odot 2}$ for all $i < j$ such that $A_{ij} \neq -\infty$. Let $\mathbf{A} \in \mathbb{K}^{m \times m}$ be any symmetric matrix such that $\text{sval}(\mathbf{A}) = A$. Then \mathbf{A} fulfills the conditions of Lemma 5.4. (In particular, it is positive semidefinite.)*

Proof Since $A_{ii} \in \mathbb{T}_+ \cup \{-\infty\}$ for all i , we have $A_{ii} \geq 0$ for all i . Moreover, if $A_{ij} = -\infty$, then $A_{ii} A_{jj} \geq 0$ and if $A_{ij} \neq -\infty$, then $\text{val}(A_{ii} A_{jj}) = A_{ii} \odot A_{jj} > A_{ij}^{\odot 2} = \text{val}((m-1)^2 A_{ij}^2)$. Therefore \mathbf{A} fulfills the conditions of Lemma 5.4. \square

Lemma 5.10 *Let \mathcal{T} be the set of points $x \in \mathbb{R}^n$ that fulfill the following two conditions:*

- for all $i \in [m]$ such that Q_{ii}^- is nonzero we have $Q_{ii}^+(x) > Q_{ii}^-(x)$;
- for all $i, j \in [m]$, $i < j$, such that Q_{ij} is nonzero we have $Q_{ii}^+(x) \odot Q_{jj}^+(x) > (Q_{ij}(x))^{\odot 2}$.

Then $\text{cl}(\mathcal{T}) \subset \text{val}(\mathcal{S}^{\text{in}} \cap \mathbb{K}_{>0}^n)$ for every spectrahedron $\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^n \mid Q(x) \succcurlyeq 0\}$ such that $\text{sval}(Q^{(k)}) = Q^{(k)}$.

Proof By Theorem 4.1, the set $\text{val}(\mathcal{S}^{\text{in}} \cap \mathbb{K}_{>0}^n)$ is closed. Therefore, it is enough to prove that $\mathcal{T} \subset \text{val}(\mathcal{S}^{\text{in}})$. Fix any $x \in \mathcal{T}$ and take any lift $\mathbf{x} \in \text{val}^{-1}(x) \cap \mathbb{K}_{\geq 0}^n$. Let $A := \text{sval}(Q(x))$. For any i such that Q_{ii}^- is nonzero we have $\text{val}(Q_{ii}^+(\mathbf{x})) = Q_{ii}^+(x) > Q_{ii}^-(x) = \text{val}(Q_{ii}^-(\mathbf{x}))$. Therefore $A_{ii} = \text{sval}(Q_{ii}(\mathbf{x})) = Q_{ii}^+(x) \in \mathbb{T}_+ \cup \{-\infty\}$ for all i (even if Q_{ii}^- is zero). Furthermore, we have $A_{ij} = Q_{ij}(x)$ for any $i < j$. Therefore, for any $i < j$ such that $A_{ij} \neq -\infty$, we have $A_{ii} \odot A_{jj} > A_{ij}^{\odot 2}$. Hence, by Lemma 5.9, $Q(x)$ fulfills the conditions of Lemma 5.4. In other words, $x \in \mathcal{S}^{\text{in}}$ and $x \in \text{val}(\mathcal{S}^{\text{in}} \cap \mathbb{K}_{>0}^n)$. \square

Assumption 5.11 We suppose that for every matrix $Q^{(k)}$ and every pair $i \neq j$ such that $Q_{ii}^{(k)}$ and $Q_{jj}^{(k)}$ belong to \mathbb{T}_+ the inequality $Q_{ii}^{(k)} + Q_{jj}^{(k)} \neq 2|Q_{ij}^{(k)}|$ holds.

We point out that Assumption 5.11 can be interpreted in terms of the nonsingularity of some (tropical) minors of order 2 of the matrices $Q^{(k)}$.

Theorem 5.12 Let $\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^n \mid Q(x) \succcurlyeq 0\}$ be a spectrahedron described by Metzler matrices $Q^{(1)}, \dots, Q^{(n)}$ such that $\text{sval}(Q^{(k)}) = Q^{(k)}$. Suppose that Assumption 5.11 holds and that the set $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)}) \cap \mathbb{R}^n$ is regular. Then

$$\text{val}(\mathcal{S} \cap \mathbb{K}_{>0}^n) = \mathcal{S}(Q^{(1)}, \dots, Q^{(n)}) \cap \mathbb{R}^n.$$

Proof Let \mathcal{T} be defined as in Lemma 5.10. Using the same arguments as in the proof of Proposition 5.7, we can show that $\text{val}(\mathcal{S}^{\text{out}}) \subset \mathcal{S}$, and subsequently, $\text{val}(\mathcal{S}^{\text{out}} \cap \mathbb{K}_{>0}^n) \subset \mathcal{S} \cap \mathbb{R}^n$. Then, by Lemma 5.10, it is enough to show that $\text{cl}(\mathcal{T}) = \mathcal{S} \cap \mathbb{R}^n$. Observe that the inequalities defining \mathcal{S} so that Q_{ij}^- ($i \leq j$) is the zero tropical polynomial are trivially satisfied. Therefore, \mathcal{S} can be expressed as the set of points $x \in \mathbb{T}^n$ verifying:

- for all $i \in [m]$ such that Q_{ii}^- is nonzero, $Q_{ii}^+(x) \geq Q_{ii}^-(x)$;
- for all $i, j \in [m]$, $i < j$, such that Q_{ij} (or, equivalently, Q_{ij}^-) is nonzero, $Q_{ii}^+(x) \odot Q_{jj}^+(x) \geq (Q_{ij}(x))^{\odot 2}$.

We denote by \mathcal{E} the set of $(i, j) \in [m] \times [m]$ such that $i \leq j$ and Q_{ij}^- is nonzero. Since $\mathcal{S} \cap \mathbb{R}^n$ is supposed to be regular, we propose to use Lemma 2.3, and thus, to exhibit nonzero tropical polynomials P_{ij} such that

$$\mathcal{S} \cap \mathbb{R}^n = \{x \in \mathbb{R}^n : \forall (i, j) \in \mathcal{E}, P_{ij}^+(x) \geq P_{ij}^-(x)\} \quad (16)$$

and

$$\mathcal{T} = \{x \in \mathbb{R}^n : \forall (i, j) \in \mathcal{E}, P_{ij}^+(x) > P_{ij}^-(x)\}. \quad (17)$$

In other words, we want to express the inequalities of the form $Q_{ii}^+(x) > Q_{ii}^-(x)$ and $Q_{ii}^+(x) \odot Q_{jj}^+(x) > (Q_{ij}(x))^{\odot 2}$ as tropical polynomial inequalities in which no term appears both on the left- and on the right-hand side. The inequalities of the first kind already satisfy this condition, and it suffices to set $P_{ii} := Q_{ii}$ for all $i \in [m]$ such that $(i, i) \in \mathcal{E}$. In contrast, we have to transform the inequalities of the second kind into equivalent constraints of the form $P_{ij}^+(x) > P_{ij}^-(x)$, where $(i, j) \in \mathcal{E}$ and $i < j$. To this end, we use Assumption 5.11. First, observe that the functions $(Q_{ij}(x))^{\odot 2}$ and $\bigoplus_k (Q_{ij}^{(k)} \odot x_k)^{\odot 2}$ are equal. Therefore, we can replace the inequalities $Q_{ii}^+(x) \odot Q_{jj}^+(x) > (Q_{ij}(x))^{\odot 2}$ by $Q_{ii}^+(x) \odot Q_{jj}^+(x) > \bigoplus_k (Q_{ij}^{(k)} \odot x_k)^{\odot 2}$. Now, we can define a formal subtraction of these tropical expressions. More precisely, for every $(i, j) \in \mathcal{E}$ such that $i < j$, we define

$$P_{ij} := \left(\bigoplus_{\substack{k \neq l \\ Q_{ii}^{(k)}, Q_{jj}^{(l)} \in \mathbb{T}_+}} (Q_{ii}^{(k)} \odot Q_{jj}^{(l)}) \odot (X_k \odot X_l) \right) \oplus \left(\bigoplus_{\substack{Q_{ii}^{(k)}, Q_{jj}^{(k)} \in \mathbb{T}_+ \\ \text{or } |Q_{ij}^{(k)}| \neq -\infty}} \alpha_k \odot X_k^{\odot 2} \right),$$

where α_k is given by

$$\alpha_k := \begin{cases} Q_{ii}^{(k)} \odot Q_{jj}^{(k)} & \text{if } Q_{ii}^{(k)}, Q_{jj}^{(k)} \in \mathbb{T}_+ \text{ and } Q_{ii}^{(k)} + Q_{jj}^{(k)} > 2|Q_{ij}^{(k)}|, \\ \ominus(Q_{ij}^{(k)})^{\odot 2} & \text{otherwise.} \end{cases}$$

Recall that any inequality of the form $\max(x, \alpha + y) > \max(x', \beta + y)$ is equivalent to $\max(x, \alpha + y) > x'$ if $\alpha > \beta$, and to $x > \max(x', \beta + y)$ if $\beta > \alpha$. Therefore, Assumption 5.11 ensures that $P_{ij}^+(x) > P_{ij}^-(x)$ is equivalent to $Q_{ii}^+(x) \odot Q_{jj}^+(x) > \bigoplus_k (Q_{ij}^{(k)} \odot x_k)^{\odot 2}$. The same applies to the nonstrict counterparts of these inequalities. We conclude that (16) and (17) are satisfied. \square

Theorem 5.13 *Let $\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^n : Q(x) \succcurlyeq 0\}$ be a spectrahedron described by Metzler matrices $Q^{(1)}, \dots, Q^{(n)}$ such that $\text{sval}(Q^{(k)}) = Q^{(k)}$. Suppose that Assumption 5.11 is fulfilled and that every stratum of the set $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$ is regular. Then*

$$\text{val}(\mathcal{S}) = \mathcal{S}(Q^{(1)}, \dots, Q^{(n)}).$$

Proof Fix a nonempty subset $K \subset [n]$. Observe that the stratum of $\text{val}(\mathcal{S})$ associated with K is equal to $\text{val}(\mathcal{S}^{(K)} \cap \mathbb{K}_{>0}^K)$, where $\mathcal{S}^{(K)}$ is the spectrahedron described by $(Q^{(k)})_{k \in K}$. Similarly, the stratum of \mathcal{S} associated with K is equal to $\mathcal{S}^{(K)} \cap \mathbb{R}^K$, where $\mathcal{S}^{(K)}$ denotes the tropical Metzler spectrahedron described by $(Q^{(k)})_{k \in K}$. Therefore, we obtain the claim by applying Theorem 5.12 to every stratum. \square

5.3 Non-Metzler Spectrahedra

In this section, we abandon the Metzler assumption that was imposed in the previous section. Let $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$ be symmetric tropical matrices. We introduce the

set $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$ (or simply \mathcal{S}) of points $x \in \mathbb{T}^n$ that fulfill the following two conditions:

- for all $i \in [m]$, $Q_{ii}^+(x) \geq Q_{ii}^-(x)$;
- for all $i, j \in [m]$, $i < j$, we have $Q_{ii}^+(x) \odot Q_{jj}^+(x) \geq (Q_{ij}^+(x) \oplus Q_{ij}^-(x))^{\odot 2}$ or $Q_{ij}^+(x) = Q_{ij}^-(x)$.

We point out that this generalizes Definition 5.5 to the case of non-Metzler matrices. More precisely, if the matrices $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$ are Metzler, then $Q_{ij}^+(x) = -\infty$ and $(Q_{ij}^+(x) \oplus Q_{ij}^-(x))^{\odot 2} = (Q_{ij}(x))^{\odot 2}$ for all $i \neq j$ and $x \in \mathbb{T}^n$. However, if the matrices $Q^{(k)}$ are not Metzler, then we have two possibilities. If Q_{ij} does not vanish on x , then $(Q_{ij}^+(x) \oplus Q_{ij}^-(x))^{\odot 2} = (Q_{ij}(x))^{\odot 2}$ and we recover the same kind of inequality as in the Metzler case. On the other hand, if Q_{ij} vanishes on x , then $Q_{ij}^+(x) = Q_{ij}^-(x)$ and we ignore the corresponding inequality.

We do *not* claim that the set \mathcal{S} defined above is a tropical spectrahedron. In this work we only show that this is true under some additional assumptions (which are generically fulfilled as shown in Sect. 5.4). First, we need some notation. For every subset

$$\Sigma \subset \{(i, j) \in [m]^2 : i < j\}$$

we denote

$$\Sigma^{\complement} := \{(i, j) \in [m]^2 : i < j, (i, j) \notin \Sigma\}.$$

For every Σ and every $\diamond \in \{\leq, \geq\}^{\Sigma^{\complement}}$ we define $\mathcal{S}_{\Sigma, \diamond}(Q^{(1)}, \dots, Q^{(n)})$ (or $\mathcal{S}_{\Sigma, \diamond}$ for short) as the set of all $x \in \mathbb{T}^n$ such that

- for all $i \in [m]$, $Q_{ii}^+(x) \geq Q_{ii}^-(x)$;
- for all $i, j \in [m]$, $i < j$, $(i, j) \in \Sigma$, $Q_{ii}^+(x) \odot Q_{jj}^+(x) \geq (Q_{ij}^+(x) \oplus Q_{ij}^-(x))^{\odot 2}$;
- for all $i, j \in [m]$, $i < j$, $(i, j) \in \Sigma^{\complement}$, $Q_{ij}^+(x) \diamond_{(i,j)} Q_{ij}^-(x)$.

Observe that we have the equality

$$\mathcal{S} = \bigcup_{\Sigma} \bigcap_{\diamond} \mathcal{S}_{\Sigma, \diamond}, \quad (18)$$

where the intersection goes over every $\diamond \in \{\leq, \geq\}^{\Sigma^{\complement}}$ and the union goes over every $\Sigma \subset \{(i, j) \in [m]^2 : i < j\}$. Moreover, we note that every set $\mathcal{S}_{\Sigma, \diamond}$ is a tropical Metzler spectrahedron (we give a detailed proof of this fact in the proof of Theorem 5.17).

Let $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{K}^{m \times m}$ be any symmetric matrices such that $\text{sval}(Q^{(k)}) = Q^{(k)}$, and $\mathcal{S} := \{x \in \mathbb{K}_{\geq 0}^n : Q(x) \succ 0\}$ be the associated spectrahedron. We will use the following observation, which already appeared in the proof of [3, Cor. 3.6] on the tropicalization of polyhedra. We denote by $\text{conv}(S)$ the convex hull of the set $S \subset \mathbb{K}^n$.

Lemma 5.14 Let $\mathbf{a}^{(1)}, \dots, \mathbf{a}^{(p)} \in \mathbb{K}^n$ and $\mathbf{b} \in \mathbb{K}^p$. Suppose that for every sign pattern $\delta \in \{+1, -1\}^p$ there is a point $\mathbf{x}^\delta \in \mathbb{K}^n$ such that for all $s \in [p]$ we have $\delta_s(\langle \mathbf{a}^{(s)}, \mathbf{x}^\delta \rangle - \mathbf{b}_s) \geq 0$. Then, there exists a point $\mathbf{y} \in \text{conv}_\delta \{\mathbf{x}^\delta\}$ such that for all s , $\langle \mathbf{a}^{(s)}, \mathbf{y} \rangle = \mathbf{b}_s$.

Proof If $p = 1$ then we have two points $\mathbf{x}^{(1)}$ and $\mathbf{x}^{(2)}$ such that $\langle \mathbf{a}^{(1)}, \mathbf{x}^{(1)} \rangle \geq \mathbf{b}_1$ and $\langle \mathbf{a}^{(1)}, \mathbf{x}^{(2)} \rangle \leq \mathbf{b}_1$. Therefore, there exists λ such that $0 \leq \lambda \leq 1$ and $\langle \mathbf{a}^{(1)}, \lambda \mathbf{x}^{(1)} + (1 - \lambda) \mathbf{x}^{(2)} \rangle = \mathbf{b}_1$. This completes the proof for $p = 1$.

Suppose that the claim is true for p . We will prove it for $p + 1$. Take

$$\Delta^+ := \{\delta \in \{+1, -1\}^{p+1} : \text{last entry of } \delta \text{ is equal to } +1\}$$

and

$$\Delta^- := \{\delta \in \{+1, -1\}^{p+1} : \text{last entry of } \delta \text{ is equal to } -1\}.$$

By the induction hypothesis, there exists a point $\mathbf{x}^{(1)} \in \text{conv}_{\delta \in \Delta^+} \{\mathbf{x}^\delta\}$ such that $\langle \mathbf{a}^{(s)}, \mathbf{x}^{(1)} \rangle = \mathbf{b}_s$ for all $s \leq p$. Moreover, we have $\langle \mathbf{a}^{(p+1)}, \mathbf{x}^\delta \rangle \geq \mathbf{b}_{p+1}$ for all $\delta \in \Delta^+$ and therefore $\langle \mathbf{a}^{(p+1)}, \mathbf{x}^{(1)} \rangle \geq \mathbf{b}_{p+1}$. Analogously, there exists a point $\mathbf{x}^{(2)} \in \text{conv}_{\delta \in \Delta^-} \{\mathbf{x}^\delta\}$ such that $\langle \mathbf{a}^{(s)}, \mathbf{x}^{(2)} \rangle = \mathbf{b}_s$ for all $s \leq p$ and $\langle \mathbf{a}^{(p+1)}, \mathbf{x}^{(2)} \rangle \leq \mathbf{b}_{p+1}$. Therefore, there is a point $\mathbf{y} \in \text{conv}\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}\} \subset \text{conv}_\delta \{\mathbf{x}^\delta\}$ such that $\langle \mathbf{a}^{(p+1)}, \mathbf{y} \rangle = \mathbf{b}_{p+1}$. Furthermore, since $\langle \mathbf{a}^{(s)}, \mathbf{x}^{(1)} \rangle = \langle \mathbf{a}^{(s)}, \mathbf{x}^{(2)} \rangle = \mathbf{b}_s$ for all $s \leq p$, we have $\langle \mathbf{a}^{(s)}, \mathbf{y} \rangle = \mathbf{b}_s$ for all $s \leq p$. \square

Lemma 5.15 We have the inclusion $\text{val}(\mathcal{S}^{\text{out}} \cap \mathbb{K}_{>0}^n) \subset \mathcal{S} \cap \mathbb{R}^n$.

Proof Take a point $\mathbf{x} \in \mathcal{S}^{\text{out}} \cap \mathbb{K}_{>0}^n$ and denote $x := \text{val}(\mathbf{x})$. For every $i \in [m]$ we have $Q_{ii}(\mathbf{x}) \geq 0$ and hence $Q_{ii}^+(\mathbf{x}) \geq Q_{ii}^-(\mathbf{x})$. Furthermore, for every $i < j$ such that $\text{val}(Q_{ij}^+(\mathbf{x})) \neq \text{val}(Q_{ij}^-(\mathbf{x}))$, we have $\text{val}(Q_{ij}(\mathbf{x})) = Q_{ij}^+(\mathbf{x}) \oplus Q_{ij}^-(\mathbf{x})$ and hence $Q_{ii}^+(\mathbf{x}) \odot Q_{jj}^+(\mathbf{x}) \geq (Q_{ij}^+(\mathbf{x}) \oplus Q_{ij}^-(\mathbf{x}))^{\odot 2}$. On the other hand, for every $i < j$ such that $\text{val}(Q_{ij}^+(\mathbf{x})) = \text{val}(Q_{ij}^-(\mathbf{x}))$ we have $Q_{ii}^+(\mathbf{x}) = Q_{jj}^-(\mathbf{x})$. In particular, $x \in \mathcal{S} \cap \mathbb{R}^n$. \square

In Lemma 5.10 we introduced the symbol \mathcal{T} to denote the set of all real points that fulfill the strict version of nontrivial inequalities defining a tropical Metzler spectrahedron \mathcal{S} . Likewise, we denote by $\mathcal{T}_{\Sigma, \diamond}$ the set of all points $x \in \mathbb{R}^n$ which fulfill the strict versions of (nontrivial) inequalities defining $\mathcal{S}_{\Sigma, \diamond}$.

Lemma 5.16 We have

$$\bigcup_{\Sigma} \bigcap_{\diamond} \text{cl}(\mathcal{T}_{\Sigma, \diamond}) \subset \text{val}(\mathcal{S}^{\text{in}} \cap \mathbb{K}_{>0}^n).$$

Proof Fix any Σ and take $x \in \bigcap_{\diamond} \text{cl}(\mathcal{T}_{\Sigma, \diamond})$. By Lemma 5.10, for every $\diamond \in \{\leq, \geq\}^{\mathbb{C}}$ there exists a lift $\mathbf{x}^\diamond \in \mathbb{K}_{>0}^n \cap \text{val}^{-1}(x)$ such that we have the inequalities

$$\begin{aligned} \forall i, \mathbf{Q}_{ii}(\mathbf{x}^\diamond) &\geq 0, \\ \forall (i, j) \in \Sigma, \mathbf{Q}_{ii}(\mathbf{x}^\diamond) \mathbf{Q}_{jj}(\mathbf{x}^\diamond) &\geq (m-1)^2 (\mathbf{Q}_{ij}(\mathbf{x}^\diamond))^2, \\ \forall (i, j) \in \Sigma^c, \mathbf{Q}_{ij}(\mathbf{x}^\diamond) &= 0. \end{aligned}$$

Observe that the set

$$\{\mathbf{y} \in \mathbb{K}_{\geq 0}^n : \forall i, \mathbf{Q}_{ii}(\mathbf{y}) \geq 0 \wedge \forall (i, j) \in \Sigma, \mathbf{Q}_{ii}(\mathbf{y}) \mathbf{Q}_{jj}(\mathbf{y}) \geq (m-1)^2 (\mathbf{Q}_{ij}(\mathbf{y}))^2\}$$

is convex. Indeed, it is a spectrahedron defined by some block diagonal matrices with blocks of size at most 2. Therefore, by Lemma 5.14, there exists a point $\mathbf{z} \in \text{conv}_\diamond \{\mathbf{x}^\diamond\}$ such that

$$\begin{aligned} \forall i, \mathbf{Q}_{ii}(\mathbf{z}) &\geq 0, \\ \forall (i, j) \in \Sigma, \mathbf{Q}_{ii}(\mathbf{z}) \mathbf{Q}_{jj}(\mathbf{z}) &\geq (m-1)^2 (\mathbf{Q}_{ij}(\mathbf{z}))^2, \\ \forall (i, j) \in \Sigma^c, \mathbf{Q}_{ij}(\mathbf{z}) &= 0. \end{aligned}$$

In particular, we have $\mathbf{Q}_{ii}(\mathbf{z}) \mathbf{Q}_{jj}(\mathbf{z}) \geq (m-1)^2 (\mathbf{Q}_{ij}(\mathbf{z}))^2$ for all (i, j) such that $i \neq j$. Therefore $\mathbf{z} \in \mathcal{S}^{\text{in}}$. Moreover, since $\mathbf{x}^\diamond \in \mathbb{K}_{>0}^n \cap \text{val}^{-1}(x)$ for all \diamond , we have $\mathbf{z} \in \mathbb{K}_{>0}^n \cap \text{val}^{-1}(x)$. \square

Theorem 5.17 *Let $\mathcal{S} = \{\mathbf{x} \in \mathbb{K}_{\geq 0}^n : \mathbf{Q}(\mathbf{x}) \succcurlyeq 0\}$ be a spectrahedron described by matrices $\mathbf{Q}^{(1)}, \dots, \mathbf{Q}^{(n)}$ such that $\text{sval}(\mathbf{Q}^{(k)}) = Q^{(k)}$. Suppose that Assumption 5.11 is fulfilled and that every stratum of the set $\mathcal{S}_{\Sigma, \diamond}(\mathbf{Q}^{(1)}, \dots, \mathbf{Q}^{(n)})$ is regular for every choice of (Σ, \diamond) . Then*

$$\text{val}(\mathcal{S}) = \mathcal{S}(Q^{(1)}, \dots, Q^{(n)}).$$

Proof We focus on the proof of the identity $\text{val}(\mathcal{S} \cap \mathbb{K}_{>0}^n) = \mathcal{S}(Q^{(1)}, \dots, Q^{(n)}) \cap \mathbb{R}^n$, as the generalization to all strata can be obtained analogously to the proof of Theorem 5.13. Let (Σ, \diamond) be fixed and observe that $\mathcal{S}_{\Sigma, \diamond}$ is a tropical Metzler spectrahedron. More precisely, it is described by the following tropical block diagonal matrices

$$\begin{bmatrix} \tilde{Q}^{(k)} & -\infty \\ -\infty & R^{(k)} \end{bmatrix}, \quad (19)$$

where $\tilde{Q}^{(k)} \in \mathbb{T}_{\pm}^{m \times m}$ is the symmetric matrix defined by

$$\tilde{Q}_{ij}^{(k)} := \begin{cases} Q_{ii}^{(k)} & \text{if } i = j, \\ \ominus |Q_{ij}^{(k)}| & \text{if } (i, j) \in \Sigma, \\ -\infty & \text{if } (i, j) \in \Sigma^c. \end{cases}$$

and $R^{(k)} \in \mathbb{T}_{\pm}^{\Sigma^{\mathbb{G}} \times \Sigma^{\mathbb{G}}}$ is the (tropical) diagonal matrix consisting of the coefficients $Q_{ij}^{(k)}$ if $\diamond_{(i,j)}$ is equal to \geq and $\ominus Q_{ij}^{(k)}$ otherwise, where (i, j) ranges over the set $\Sigma^{\mathbb{G}}$.

It can be verified that, as soon as the matrices $Q^{(k)}$ satisfy Assumption 5.11, this assumption is also satisfied by all block matrices (19). In consequence, as shown in the proof of Theorem 5.12, the sets $\text{cl}(\mathcal{T}_{\Sigma, \diamond})$ and $\mathcal{S}_{\Sigma, \diamond} \cap \mathbb{R}^n$ coincide. Then, the theorem follows from (18) and Lemmas 5.16 and 5.15. \square

We now show that, under the regularity conditions discussed above, tropical spectrahedra arise as tropicalizations of minors of size 1 and 2 of the linear matrix inequalities that define them. To do so, let $\mathcal{S} = \{\mathbf{x} \in \mathbb{K}_{\geq 0}^n : \mathbf{Q}(\mathbf{x}) \succcurlyeq 0\}$ be a spectrahedron described by matrices $Q^{(1)}, \dots, Q^{(n)}$ as above. For every $i, j \in [m]$ such that $i \neq j$, let

$$P_{ij} := Q_{ii}(\mathbf{x})Q_{jj}(\mathbf{x}) - Q_{ij}^2(\mathbf{x})$$

denote the corresponding 2×2 minor of the matrix pencil $\mathbf{Q}(\mathbf{x})$. Furthermore, for every $i \neq j$ let $P_{ij} := \text{trop}(P_{ij})$ and, for every $i \in [m]$, let $P_{ii} := \text{trop}(Q_{ii})$. In this way, P_{ij} are the tropicalizations of the 1×1 and 2×2 minors of $\mathbf{Q}(\mathbf{x})$. Finally, let

$$\mathcal{S}^{\text{form}} := \{x \in \mathbb{T}^n : \forall i, P_{ii}^+(x) \geq P_{ii}^-(x) \wedge \forall i \neq j, P_{ij}^+(x) \geq P_{ij}^-(x)\}$$

denote the set obtained by this formal tropicalization.

Theorem 5.18 *Suppose that the assumptions of Theorem 5.17 are satisfied. Then, the tropical spectrahedron $\text{val}(\mathcal{S})$ is determined only by the formal tropicalization of 1×1 and 2×2 minors, in other words*

$$\text{val}(\mathcal{S}) = \mathcal{S}^{\text{form}}.$$

We shall deduce this theorem from the following general “sandwich” lemma, which does not require any genericity assumption.

Lemma 5.19 *We have $\text{val}(\mathcal{S}) \subset \text{val}(\mathcal{S}^{\text{out}}) \subset \mathcal{S}^{\text{form}} \subset \mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$.*

Let us first explain why Theorem 5.18 follows in a straightforward way from Lemma 5.19.

Derivation of Theorem 5.18 from Lemma 5.19 By Lemma 5.19 we have $\text{val}(\mathcal{S}) \subset \mathcal{S}^{\text{form}} \subset \mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$. Furthermore, Theorem 5.17 implies that $\text{val}(\mathcal{S}) = \mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$. Hence, we have the equality $\text{val}(\mathcal{S}) = \mathcal{S}^{\text{form}}$. \square

The proof of Lemma 5.19 requires more technical efforts.

Proof of Lemma 5.19 The inclusion $\text{val}(\mathcal{S}) \subset \text{val}(\mathcal{S}^{\text{out}})$ is trivial. The inclusion $\text{val}(\mathcal{S}^{\text{out}}) \subset \mathcal{S}^{\text{form}}$ follows from Lemma 2.2, as in the proof of Corollary 4.8. It remains to show that $\mathcal{S}^{\text{form}}$ is included in $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)})$. To do so, let us choose an arbitrary point $\bar{x} \in \mathcal{S}^{\text{form}}$. We first observe that $P_{ii} = Q_{ii}$ for all i by the definition

of the formal tropicalization. In particular, $Q_{ii}^+(\bar{x}) \geq Q_{ii}^-(\bar{x})$ for all $i \in [m]$. We must show that

$$Q_{ii}^+(\bar{x}) \odot Q_{jj}^+(\bar{x}) \geq (Q_{ij}^+(\bar{x}) \oplus Q_{ij}^-(\bar{x}))^{\odot 2} \quad \text{or} \quad Q_{ij}^+(\bar{x}) = Q_{ij}^-(\bar{x}) \quad \text{for all } i \neq j.$$

It suffices to consider $i \neq j$ such that $Q_{ij}^+(\bar{x}) \neq Q_{ij}^-(\bar{x})$ and to show that $Q_{ii}^+(\bar{x}) \odot Q_{jj}^+(\bar{x}) \geq (Q_{ij}^+(\bar{x}) \oplus Q_{ij}^-(\bar{x}))^{\odot 2}$. Since $Q_{ij}^+(\bar{x}) \neq Q_{ij}^-(\bar{x})$, at least one of the polynomials Q_{ij}^+ , Q_{ij}^- is nonzero. By the definition of Q_{ij}^+ and Q_{ij}^- we have $Q_{ij}^+(x) \oplus Q_{ij}^-(x) = \max_{k \in [n]} \{\text{val}(Q_{ij}^{(k)}) + x_k\}$. Let $k' \in [n]$ be an index such that

$$Q_{ij}^+(\bar{x}) \oplus Q_{ij}^-(\bar{x}) = \text{val}(Q_{ij}^{(k')}) + \bar{x}_{k'}. \quad (20)$$

Let us point out that $\text{val}(Q_{ij}^{(k')}) \neq -\infty$ and $\bar{x}_{k'} \neq -\infty$. Indeed, $Q_{ij}^+(\bar{x}) \neq Q_{ij}^-(\bar{x})$ implies that $Q_{ij}^+(\bar{x}) \oplus Q_{ij}^-(\bar{x}) \neq -\infty$, so neither $\text{val}(Q_{ij}^{(k')})$ nor $\bar{x}_{k'}$ can be equal to $-\infty$. To prove the claim, it remains to show that

$$2\text{val}(Q_{ij}^{(k')}) + 2\bar{x}_{k'} \leq Q_{ii}^+(\bar{x}) \odot Q_{jj}^+(\bar{x}). \quad (21)$$

Consider two cases. First, suppose that $2\text{val}(Q_{ij}^{(k')}) \leq \text{val}(Q_{ii}^{(k')}) + \text{val}(Q_{jj}^{(k')})$. In this case, using the inequalities $Q_{ii}^+(\bar{x}) \geq Q_{ii}^-(\bar{x})$ and $Q_{jj}^+(\bar{x}) \geq Q_{jj}^-(\bar{x})$, we obtain

$$\begin{aligned} 2\text{val}(Q_{ij}^{(k')}) + 2\bar{x}_{k'} &\leq \text{val}(Q_{ii}^{(k')}) + \text{val}(Q_{jj}^{(k')}) + 2\bar{x}_{k'} \\ &\leq (Q_{ii}^+(\bar{x}) \oplus Q_{ii}^-(\bar{x})) \odot (Q_{jj}^+(\bar{x}) \oplus Q_{jj}^-(\bar{x})) = Q_{ii}^+(\bar{x}) \odot Q_{jj}^+(\bar{x}), \end{aligned}$$

and so, (21) is established in this case. Second, suppose that $2\text{val}(Q_{ij}^{(k')}) > \text{val}(Q_{ii}^{(k')}) + \text{val}(Q_{jj}^{(k')})$. This case requires a longer argument.

To start, note that $2\text{val}(Q_{ij}^{(k')}) > \text{val}(Q_{ii}^{(k')}) + \text{val}(Q_{jj}^{(k')})$ implies $(Q_{ij}^{(k')})^2 > Q_{ii}^{(k')} Q_{jj}^{(k')}$ and $\text{val}(Q_{ii}^{(k')} Q_{jj}^{(k')} - (Q_{ij}^{(k')})^2) = 2\text{val}(Q_{ij}^{(k')})$. Furthermore, the minor P_{ij} is given as

$$\begin{aligned} P_{ij}(\mathbf{x}) &= \sum_{k \in [n]} (Q_{ii}^{(k)} Q_{jj}^{(k)} - (Q_{ij}^{(k)})^2) \mathbf{x}_k^2 \\ &\quad + \sum_{k < l} (Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)} - 2Q_{ij}^{(k)} Q_{ij}^{(l)}) \mathbf{x}_k \mathbf{x}_l. \end{aligned} \quad (22)$$

In particular, the polynomial $P_{ij}(\mathbf{x})$ has at least one monomial with (strictly) negative coefficient, namely $(Q_{ii}^{(k')} Q_{jj}^{(k')} - (Q_{ij}^{(k')})^2) \mathbf{x}_{k'}^2$. Remembering that $P_{ij}^- = \text{trop}(P_{ij}^-)$ is the tropical polynomial whose monomials are obtained by tropicalizing the monomials of P_{ij} with negative coefficients (as in Sect. 2.2), we see that P_{ij}^- is a nonzero tropical polynomial and that it contains a monomial of the form $\text{val}(Q_{ij}^{(k')})^{\odot 2} \odot \mathbf{x}_{k'}^{\odot 2}$.

Furthermore, recall that we have $P_{ij}^+(\bar{x}) \geq P_{ij}^-(\bar{x})$ and that we already showed that $\bar{x}_{k'} \neq -\infty$. Hence, we also have $P_{ij}^-(\bar{x}) \neq -\infty$, which implies that $P_{ij}^+(\bar{x}) \neq -\infty$. Therefore, the polynomial P_{ij} has at least one monomial with (strictly) positive coefficient. In other words, there exist indices $k, l \in [n]$ (possibly equal) such that

$$Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)} - 2Q_{ij}^{(k)} Q_{ij}^{(l)} > 0.$$

By definition, $P_{ij}^+(\bar{x})$ is the maximum of the tropical monomials arising from the monomials with positive coefficients in (22), evaluated at point \bar{x} . Choosing a monomial attaining this maximum yields a pair of indices (k, l) such that $P_{ij}^+(\bar{x}) = \text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)} - 2Q_{ij}^{(k)} Q_{ij}^{(l)}) + \bar{x}_k + \bar{x}_l$. Since $P_{ij}^+(\bar{x}) \geq P_{ij}^-(\bar{x})$, we also have

$$P_{ij}^+(\bar{x}) \oplus P_{ij}^-(\bar{x}) = \text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)} - 2Q_{ij}^{(k)} Q_{ij}^{(l)}) + \bar{x}_k + \bar{x}_l.$$

Hence, we get

$$\begin{aligned} 2\text{val}(Q_{ij}^{(k')}) + 2\bar{x}_{k'} &= \text{val}(Q_{ii}^{(k')} Q_{jj}^{(k')} - (Q_{ij}^{(k')})^2) + 2\bar{x}_{k'} \\ &\leq P_{ij}^+(\bar{x}) \oplus P_{ij}^-(\bar{x}) \leq \max\{\text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)}), \text{val}(Q_{ij}^{(k)} Q_{ij}^{(l)})\} + \bar{x}_k + \bar{x}_l. \end{aligned} \quad (23)$$

We will show that

$$\max\{\text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)}), \text{val}(Q_{ij}^{(k)} Q_{ij}^{(l)})\} = \text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)}).$$

To do so, suppose that $\text{val}(Q_{ij}^{(k)} Q_{ij}^{(l)}) > \text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)})$. Then, we have $Q_{ij}^{(k)} Q_{ij}^{(l)} < 0$, because $Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)} - 2Q_{ij}^{(k)} Q_{ij}^{(l)} \geq 0$. In particular, since $Q_{ij}^+(\bar{x}) \neq Q_{ij}^-(\bar{x})$, we obtain

$$\begin{aligned} 2\text{val}(Q_{ij}^{(k')}) + 2\bar{x}_{k'} &\leq \text{val}(Q_{ij}^{(k)} Q_{ij}^{(l)}) + \bar{x}_k + \bar{x}_l \\ &\leq Q_{ij}^+(\bar{x}) \odot Q_{ij}^-(\bar{x}) < (Q_{ij}^+(\bar{x}) \oplus Q_{ij}^-(\bar{x}))^{\odot 2}, \end{aligned}$$

which gives a contradiction with (20). Hence, $\text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)}) \geq \text{val}(Q_{ij}^{(k)} Q_{ij}^{(l)})$ and (23) gives

$$\begin{aligned} 2\text{val}(Q_{ij}^{(k')}) + 2\bar{x}_{k'} &\leq \text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)} + Q_{ii}^{(l)} Q_{jj}^{(k)}) + \bar{x}_k + \bar{x}_l \\ &\leq \max\{\text{val}(Q_{ii}^{(k)} Q_{jj}^{(l)}) + \bar{x}_k + \bar{x}_l, \text{val}(Q_{ii}^{(l)} Q_{jj}^{(k)}) + \bar{x}_k + \bar{x}_l\} \\ &\leq (Q_{ii}^+(\bar{x}) \oplus Q_{ii}^-(\bar{x})) \odot (Q_{jj}^+(\bar{x}) \oplus Q_{jj}^-(\bar{x})) \\ &= Q_{ii}^+(\bar{x}) \odot Q_{jj}^+(\bar{x}), \end{aligned} \quad (24)$$




Fig. 3 A nonregular tropical spectrahedron that fulfills the regularity conditions of Theorem 5.17

which shows (21). This concludes the proof of Lemma 5.19, and so, of Theorem 5.18. \square

Example 5.20 Take the matrices

$$Q^{(0)} := \begin{bmatrix} a & -\infty \\ -\infty & b \end{bmatrix}, \quad Q^{(1)} := \begin{bmatrix} -\infty & c \\ c & -\infty \end{bmatrix}, \quad Q^{(2)} := \begin{bmatrix} -\infty & \ominus d \\ \ominus d & -\infty \end{bmatrix}.$$

The set $\mathcal{S}(Q^{(0)}, Q^{(1)}, Q^{(2)})$ is given by

$$\{x \in \mathbb{T}^3 : (a + x_0) + (b + x_0) \geq 2 \max\{c + x_1, d + x_2\} \text{ or } c + x_1 = d + x_2\}.$$

Moreover, $\mathcal{S}(Q^{(0)}, Q^{(1)}, Q^{(2)})$ fulfills the conditions of Theorem 5.17. Figure 3 depicts the intersection of this tropical spectrahedron with the hyperplane $\{x_0 = 0\}$ for $a = b = c = d = 0$. Note that this tropical spectrahedron is not regular for any choice of $a, b, c, d \in \mathbb{R}$. Furthermore, let

$$Q^{(0)} := \begin{bmatrix} t^a & 0 \\ 0 & t^b \end{bmatrix}, \quad Q^{(1)} := \begin{bmatrix} 0 & t^c \\ t^c & 0 \end{bmatrix}, \quad Q^{(2)} := \begin{bmatrix} 0 & -t^d \\ -t^d & 0 \end{bmatrix}$$

be a lift of the matrices $Q^{(k)}$. Then, the associated tropical spectrahedron \mathcal{S} is given by

$$\mathcal{S} = \{x \in \mathbb{K}_{\geq 0}^3 : t^{a+b} x_0^2 - t^{2c} x_1^2 - t^{2d} x_2^2 + 2t^{c+d} x_1 x_2 \geq 0\}.$$

The formal tropicalization $\mathcal{S}^{\text{form}}$ of this set is given by

$$\{x \in \mathbb{T}^3 : ((a+b) \odot x_0^{\odot 2}) \oplus ((c+d) \odot x_1 \odot x_2) \geq (2c \odot x_1^{\odot 2}) \oplus (2d \odot x_2^{\odot 2})\}.$$

It is easy to check that $\mathcal{S}^{\text{form}}$ coincides with $\mathcal{S}(Q^{(0)}, Q^{(1)}, Q^{(2)})$, as implied by Theorem 5.18.

5.4 Genericity Conditions

In this section we show that the requirements of Theorems 5.12 and 5.17 on the matrices $Q^{(k)}$ and the regularity of sets are fulfilled generically. In [3] it was shown that genericity conditions for tropical polyhedra can be described by the means of tangent digraphs. We extend this characterization to tropical spectrahedra. For this purpose, we work with hypergraphs instead of graphs. A (directed) hypergraph is a pair $\mathcal{G} := (V, E)$, where V is a finite set of vertices and E is a finite set of (hyper)edges. Every edge $e \in E$ is a pair (T_e, h_e) , where $h_e \in V$ is called the head of the edge, and T_e is a multiset with elements taken from V . We call T_e the multiset of tails of e . By $|T_e|$ we denote the cardinality of T_e (counting multiplicities). Note that we do not exclude the situation in which a head is also a tail, i.e., it is possible that $h_e \in T_e$.

Let us now define the notion of a circulation in a hypergraph. If $v \in V$ is a vertex, then by $\text{In}(v) \subset E$ we denote the set of incoming edges, i.e., the set of all edges e such that $h_e = v$. By $\text{Out}(v)$ we denote the multiset of outgoing edges, i.e., a multiset of edges e such that $v \in T_e$. We treat $\text{Out}(v)$ as a multiset, with the convention that $e \in E$ appears p times in $\text{Out}(v)$ if v appears p times in T_e . A circulation in a hypergraph is a nonzero vector $\gamma = (\gamma_e)_{e \in E}$ such that $\gamma_e \geq 0$ for all $e \in E$, and for all $v \in V$ we have the equality

$$\sum_{e \in \text{In}(v)} |T_e| \gamma_e = \sum_{e \in \text{Out}(v)} \gamma_e.$$

We always suppose that circulations are normalized, i.e., that $\sum_{e \in E} \gamma_e = 1$. Observe that if a hypergraph \mathcal{G} is fixed, then the set of all normalized circulations on \mathcal{G} forms a polytope. We say that a hypergraph *does not admit a circulation* if this polytope is empty.

In our framework, every edge has at most two tails (counting multiplicities). Hereafter, ε_k denotes the k th vector of the standard basis in \mathbb{R}^n . Given a sequence of tropical symmetric Metzler matrices $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$, and a point $x \in \mathbb{R}^n$, we construct a hypergraph associated with x , denoted \mathcal{G}_x , as follows:

- we put $V := [n]$;
- for every $i \in [m]$ verifying $Q_{ii}^+(x) = Q_{ii}^-(x) \neq -\infty$, and every pair $\varepsilon_k \in \text{Argmax}(Q_{ii}^+, x)$, $\varepsilon_l \in \text{Argmax}(Q_{ii}^-, x)$, \mathcal{G}_x contains an edge (k, l) ;
- for every $i < j$ such that $Q_{ii}^+(x) \odot Q_{jj}^+(x) = (Q_{ij}(x))^{\odot 2} \neq -\infty$ and every triple $\varepsilon_{k_1} \in \text{Argmax}(Q_{ii}^+, x)$, $\varepsilon_{k_2} \in \text{Argmax}(Q_{jj}^+, x)$, $\varepsilon_l \in \text{Argmax}(Q_{ij}, x)$, \mathcal{G}_x contains an edge $(\{k_1, k_2\}, l)$.

Lemma 5.21 *Suppose that for every $x \in \mathbb{R}^n$ the hypergraph \mathcal{G}_x does not admit a circulation. Then $Q^{(1)}, \dots, Q^{(n)}$ fulfill Assumption 5.11 and $\mathcal{S}(Q^{(1)}, \dots, Q^{(n)}) \cap \mathbb{R}^n$ is regular.*

Proof To prove the first part, suppose that we have $Q_{ii}^{(k)} + Q_{jj}^{(k)} = 2|Q_{ij}^{(k)}|$ for some $i \neq j$ and $Q_{ii}^{(k)}, Q_{jj}^{(k)} \in \mathbb{T}_+$. Take the point $x := N\varepsilon_k \in \mathbb{R}^n$. If N is large enough, then we have

$$Q_{ii}^+(x) \odot Q_{jj}^+(x) = Q_{ii}^{(k)} + Q_{jj}^{(k)} + 2N = 2|Q_{ij}^{(k)}| + 2N = (Q_{ij}(x))^{\odot 2}$$

and the hypergraph \mathcal{G}_x contains the edge $(\{k, k\}, k)$. This hypergraph admits a circulation (we put $\gamma_e := 1$ for $e = (\{k, k\}, k)$ and $\gamma_e := 0$ for other edges).

We now claim that the set $\mathcal{S} \cap \mathbb{R}^n$ is regular. Let \mathcal{T} be defined as in Lemma 5.10. Let us show that for every $x \in \mathcal{S} \cap \mathbb{R}^n$ there exists a vector $\eta \in \mathbb{R}^n$ such that $x + \rho\eta$ belongs to \mathcal{T} for $\rho > 0$ small enough. This is sufficient to prove the claim because \mathcal{T} is a subset of the interior of $\mathcal{S} \cap \mathbb{R}^n$. Fix a point $x \in \mathcal{S} \cap \mathbb{R}^n$. If x belongs to \mathcal{T} , then we can take $\eta := 0$. Otherwise, let \mathcal{G}_x denote the hypergraph associated with x . The polytope of normalized circulations of this hypergraph is empty. Therefore, by Farkas' lemma, there exists a vector $\eta \in \mathbb{R}^n$ such that for every edge $e \in E$ we have

$$\sum_{v \in T_e} \eta_v > |T_e| \eta_{h_e}.$$

Take the vector $x^{(\rho)} := x + \rho\eta$. Let us look at two cases.

First, suppose that there is $i \in [m]$ such that $Q_{ii}^+(x) = Q_{ii}^-(x) \neq -\infty$. Fix any k^* such that $\varepsilon_{k^*} \in \text{Argmax}(Q_{ii}^+, x)$ and take any l such that $\varepsilon_l \in \text{Argmax}(Q_{ii}^-, x)$. Then (k^*, l) is an edge in \mathcal{G}_x . Therefore $\eta_{k^*} > \eta_l$. Moreover, $Q_{ii}^{(k^*)} + x_{k^*} = |Q_{ii}^{(l)}| + x_l$ and hence $Q_{ii}^{(k^*)} + x_{k^*}^{(\rho)} > |Q_{ii}^{(l)}| + x_l^{(\rho)}$. Furthermore, for every $l' \notin \text{Argmax}(Q_{ii}^-, x)$ we have

$$Q_{ii}^{(k^*)} + x_{k^*} = |Q_{ii}^{(l)}| + x_l > |Q_{ii}^{(l')}| + x_{l'}.$$

Therefore $Q_{ii}^{(k^*)} + x_{k^*}^{(\rho)} > |Q_{ii}^{(l')}| + x_{l'}^{(\rho)}$ for ρ small enough. Since l, l' were arbitrary, for every sufficiently small ρ we have

$$Q_{ii}^+(x^{(\rho)}) \geq Q_{ii}^{(k^*)} + x_{k^*}^{(\rho)} > Q_{ii}^-(x^{(\rho)}).$$

The second case is analogous. If there is $i < j$ such that $Q_{ii}^+(x) \odot Q_{jj}^+(x) = (Q_{ij}(x))^{\odot 2} \neq -\infty$, then we fix (k_1^*, k_2^*) such that $\varepsilon_{k_1^*} \in \text{Argmax}(Q_{ii}^+, x)$, $\varepsilon_{k_2^*} \in \text{Argmax}(Q_{jj}^+, x)$. For every $\varepsilon_l \in \text{Argmax}(Q_{ij}, x)$, $(\{k_1^*, k_2^*\}, l)$ is an edge in \mathcal{G} . Hence $\eta_{k_1^*} + \eta_{k_2^*} > 2\eta_l$. Therefore $Q_{ii}^{(k_1^*)} + Q_{jj}^{(k_2^*)} + x_{k_1^*}^{(\rho)} + x_{k_2^*}^{(\rho)} > 2|Q_{ij}^{(l)}| + 2x_l^{(\rho)}$. As before, this implies that $Q_{ii}^+(x^{(\rho)}) \odot Q_{jj}^+(x^{(\rho)}) > (Q_{ij}(x^{(\rho)}))^{\odot 2}$ for $\rho > 0$ small enough. Since we supposed that $x \in \mathcal{S} \cap \mathbb{R}^n$, we have $x^{(\rho)} \in \mathcal{T}$ for ρ small enough. \square




Fig. 4 The hypergraph from Example 5.22, consisting of the edges $(0, 1)$, $(0, 2)$, and $(\{1, 2\}, 0)$

Example 5.22 Take the matrices

$$\begin{aligned} Q^{(0)} &:= \text{tdiag} \left(0, 0, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right), \\ Q^{(1)} &:= \text{tdiag}(\ominus 0, -\infty, 0, -\infty), \\ Q^{(2)} &:= \text{tdiag}(-\infty, \ominus 0, -\infty, 0). \end{aligned}$$

In this case, the set $\mathcal{S}(Q^{(0)}, Q^{(1)}, Q^{(2)}) \cap \mathbb{R}^3 = \{(\lambda, \lambda, \lambda) : \lambda \in \mathbb{R}\}$ is reduced to a line. In particular, it is not regular. The hypergraph associated with $(0, 0, 0)$ is depicted in Fig. 4. This hypergraph admits a circulation (we put $\gamma_e := 1/3$ for all edges).

We now want to show that the condition of Lemma 5.21 is fulfilled generically.

Lemma 5.23 *There exists a set $X \subset \mathbb{T}^d$ with $d = nm(m+1)/2$ such that every stratum of X is a finite union of hyperplanes and such that if the vector with entries $|Q_{ij}^{(k)}|$ (for $i \leq j$) does not belong to X , then the hypergraph \mathcal{G}_x does not admit a circulation for any $x \in \mathbb{R}^n$.*

Proof Fix a nonempty subset $D \subset [d]$, $|D| = d'$ and let $\mathbb{R}^{d'}$ be the stratum of \mathbb{T}^d associated with D . Suppose that $Q^{(1)}, \dots, Q^{(n)}$ are tropical Metzler matrices, that the support of the vector $|Q_{ij}^{(k)}|$ is equal to D , and that $x \in \mathbb{R}^n$ is such that \mathcal{G}_x admits a circulation. Fix any such circulation γ . For every edge $e = (k, l)$ of \mathcal{G} we can fix $i_e \in [m]$ such that $Q_{i_e i_e}^{(k)} + x_k = |Q_{i_e l}^{(l)}| + x_l \neq -\infty$. Similarly, for every edge $e = (\{k_1, k_2\}, l)$ of \mathcal{G} we can fix $i_e < j_e$ so that $Q_{i_e i_e}^{(k_1)} + Q_{j_e j_e}^{(k_2)} + x_{k_1} + x_{k_2} = 2|Q_{i_e j_e}^{(l)}| + 2x_l \neq -\infty$. We take the sum of these equalities weighted by γ . This gives the equality

$$\begin{aligned} & \sum_{k \in [n]} \sum_{e \in \text{Out}(k)} \gamma_e Q_{i_e i_e}^{(k)} + \sum_{k \in [n]} \sum_{e \in \text{Out}(k)} \gamma_e x_k \\ &= \sum_{l \in [n]} \sum_{e \in \text{In}_1(l)} \gamma_e |Q_{i_e i_e}^{(l)}| + \sum_{l \in [n]} \sum_{e \in \text{In}_2(l)} 2\gamma_e |Q_{i_e j_e}^{(l)}| + \sum_{l \in [n]} \sum_{e \in \text{In}(l)} |T_e| \gamma_e x_l, \end{aligned}$$

where $\text{In}_1(l)$ denotes the set of incoming edges with tails of cardinality 1 and $\text{In}_2(l)$ denotes the set of incoming edges with tails of cardinality 2. Since γ is a circulation,

this expression simplifies to

$$\sum_{k \in [n]} \sum_{e \in \text{Out}(k)} \gamma_e Q_{i_e i_e}^{(k)} = \sum_{l \in [n]} \sum_{e \in \text{In}_1(l)} \gamma_e |Q_{i_e i_e}^{(l)}| + \sum_{l \in [n]} \sum_{e \in \text{In}_2(l)} 2\gamma_e |Q_{i_e j_e}^{(l)}|.$$

Consider the set \mathcal{H} of all $z \in \mathbb{R}^{d'}$ such that

$$\sum_{k \in [n]} \sum_{e \in \text{Out}(k)} \gamma_e z_{i_e i_e}^{(k)} = \sum_{l \in [n]} \sum_{e \in \text{In}_1(l)} \gamma_e z_{i_e i_e}^{(l)} + \sum_{l \in [n]} \sum_{e \in \text{In}_2(l)} 2\gamma_e z_{i_e j_e}^{(l)}. \quad (25)$$

This set is a hyperplane. Indeed, suppose that the equality above is trivial (i.e., that it reduces to $0 = 0$). Take any edge e such that $\gamma_e \neq 0$ and any vertex $k \in T_e$. Then the coefficient $z_{i_e i_e}^{(k)}$ appears on the left-hand side. Moreover, we have $\text{sign}(Q_{i_e i_e}^{(k)}) = 1$. On the other hand, for every coefficient $z_{j_e j_e}^{(l)}$ that appears on the right-hand side we have $\text{sign}(Q_{j_e j_e}^{(l)}) = -1$. This gives a contradiction.

Therefore, we can construct the stratum of X associated with D (denoted X_D) as follows: we take all possible hypergraphs that can arise in our construction (since n is fixed, we have finitely many of them). Out of them, we choose those hypergraphs that admit a circulation. For every such hypergraph we pick exactly one circulation γ . After that, for every possible choice of functions $e \mapsto i_e$, $e \mapsto (i_e, j_e)$ ¹, we take a set \mathcal{H} defined as in (25). If \mathcal{H} is equal to $\mathbb{R}^{d'}$, then we ignore it. Otherwise, \mathcal{H} is a hyperplane. We take X_D to be the union of all hyperplanes obtained in this way. \square

The proof of Lemma 5.23 can be easily adapted to give a genericity condition both for Metzler and non-Metzler spectrahedra.

Theorem 5.24 *Let $Q^{(1)}, \dots, Q^{(n)} \in \mathbb{T}_{\pm}^{m \times m}$ be a sequence of symmetric tropical matrices. There exists a set $X \subset \mathbb{T}^d$ with $d = nm(m+1)/2$ such that every stratum of X is a finite union of hyperplanes and such that if the vector with entries $|Q_{ij}^{(k)}|$ (for $i \leq j$) does not belong to X , then the matrices $Q^{(1)}, \dots, Q^{(n)}$ fulfill Assumption 5.11 and for all (Σ, \diamond) , every stratum of $\mathcal{S}_{\Sigma, \diamond}(Q^{(1)}, \dots, Q^{(n)})$ is regular.*

Proof As previously, we fix a nonempty set $D \subset [d]$, $|D| = d'$, and we will present a construction of the stratum of X associated with D , denoted X_D . Take symmetric matrices $(Q^{(k)}) \in \mathbb{T}^{m \times m}$ such that the sequence $(|Q_{ij}^{(k)}|) \in \mathbb{T}^d$ has support equal to D . Take any nonempty subset $K \subset [n]$ and let $\mathcal{S}^{(K)}$ denote the set $\mathcal{S}((Q^{(k)})_{k \in K})$. Fix a pair (Σ, \diamond) and take the tropical Metzler spectrahedron $\mathcal{S}_{\Sigma, \diamond}^{(K)}$. Take any $x \in \mathbb{R}^K$ and a graph \mathcal{G}_x associated with $\mathcal{S}_{\Sigma, \diamond}^{(K)}$ (note that this graph has vertices enumerated by numbers from K). Suppose that this graph admits a circulation γ . As previously, for every edge $e = (\{k_1, k_2\}, l)$ of \mathcal{G} we can take $(i_e, j_e) \in \Sigma$ such that $Q_{i_e i_e}^{(k_1)} + Q_{j_e j_e}^{(k_2)} + x_{k_1} + x_{k_2} = 2|Q_{i_e j_e}^{(l)}| + 2x_l$. For every edge $e = (k, l)$ we have two possibilities: either there exists $i_e \in [m]$ such that $Q_{i_e i_e}^{(k)} + x_k = |Q_{i_e i_e}^{(l)}| + x_l$ or there exists $(i_e, j_e) \in \Sigma^{\mathcal{G}}$ such that

¹ Note that the dependence on D lies here, as the choice of D restricts the amount of possible functions $e \mapsto i_e$, $e \mapsto (i_e, j_e)$.

$|Q_{i_e j_e}^{(k)}| + x_k = |Q_{i_e j_e}^{(l)}| + x_l$. As before, we take the sum of these equalities weighted by γ . This gives the identity

$$\sum_{k \in [n]} \sum_{e \in \text{Out}(k)} \gamma_e |Q_{i_e j_e}^{(k)}| = \sum_{l \in [n]} \sum_{e \in \text{In}_1(l)} \gamma_e |Q_{i_e j_e}^{(l)}| + \sum_{l \in [n]} \sum_{e \in \text{In}_2(l)} 2\gamma_e |Q_{i_e j_e}^{(l)}|.$$

As previously, the set of all $z \in \mathbb{R}^{d'}$ that fulfill this equality is a hyperplane. Indeed, any coefficient $z_{i_e j_e}^{(k)}$ which appears on the left-hand side does not appear on the right-hand side (note that here we use the fact that $\Sigma \cap \Sigma^{\mathbb{G}} = \emptyset$). As before, we take all possible hypergraphs (where “all possible” takes into account the fact that K can vary), one circulation for each hypergraph, all possible functions $e \mapsto (i_e, j_e)$ (the number of such functions depends on D), and all hyperplanes that can arise in this way. The union of these hyperplanes constitutes X_D . We deduce the result from Lemma 5.21. \square

6 Concluding Remarks

We characterized the images by the valuation of nonarchimedean spectrahedra which satisfy a certain genericity condition. Our results imply that the images of nongeneric spectrahedra are still closed semilinear sets. It is an open question to characterize the semilinear sets which arise in this way. A special situation in which such a description is known in the nongeneric case concerns tropical polyhedra. It relies on the Minkowski–Weyl theorem and does not carry over to spectrahedra.

Acknowledgements We thank the referees for their comments which led to improvements of this article.

References

1. Akian, M., Gaubert, S., Guterman, A.: Linear independence over tropical semirings and beyond. In: Litvinov, G.L., Sergeev, S.N. (eds.) *Proceedings of the International Conference on Tropical and Idempotent Mathematics*. Contemporary Mathematics, vol. 495, pp. 1–38. American Mathematical Society, Providence (2009)
2. Alessandrini, D.: Logarithmic limit sets of real semi-algebraic sets. *Adv. Geom.* **13**(1), 155–190 (2013)
3. Allamigeon, X., Benchimol, P., Gaubert, S., Joswig, M.: Tropicalizing the simplex algorithm. *SIAM J. Discrete Math.* **29**(2), 751–795 (2015)
4. Allamigeon, X., Benchimol, P., Gaubert, S., Joswig, M.: Log-barrier interior point methods are not strongly polynomial. *SIAM J. Appl. Algebra Geom.* **2**(1), 140–178 (2018)
5. Allamigeon, X., Gaubert, S., Katz, R.D., Skomra, M.: Condition numbers of stochastic mean payoff games and what they say about nonarchimedean semidefinite programming. In: *Proceedings of the 23rd International Symposium on Mathematical Theory of Networks and Systems (MTNS)*, pp. 160–167. <http://mtns2018.ust.hk/media/files/0213.pdf> (2018)
6. Allamigeon, X., Gaubert, S., Skomra, M.: Solving generic nonarchimedean semidefinite programs using stochastic game algorithms. *J. Symbolic Comp.* **85**, 25–54 (2018). This is an extended version of an article which appeared in the *Proceedings of the 41st International Symposium on Symbolic and Algebraic Computation (ISSAC 2016)*
7. Allamigeon, X., Gaubert, S., Skomra, M.: The tropical analogue of the Helton–Nie conjecture is true. *J. Symbolic Comput.* **91**, 129–148 (2019). This is an extended version of an article presented at MEGA 2017, *Effective Methods in Algebraic Geometry*, Nice (France), June 12–16, 2017

8. Andersson, D., Miltersen, P.B.: The complexity of solving stochastic games on graphs. In: Dong, Y. et al. (eds.) *Proceedings of the 20th International Symposium on Algorithms and Computation (ISAAC)*, Lecture Notes in Computer Science, vol. 5878, pp. 112–121. Springer, Berlin (2009)
9. Aroca, F.: Krull-tropical hypersurfaces. *Ann. Fac. Sci. Toulouse Math.* **19**(3–4), 525–538 (2010)
10. Baker, M., Bowler, N.: Matroids over partial hyperstructures. *Adv. Math.* **343**, 821–863 (2019)
11. Banerjee, S.D.: Tropical geometry over higher dimensional local fields. *J. Reine Angew. Math.* **698**, 71–87 (2015)
12. Basu, S., Pollack, R.: *Algorithms in Real Algebraic Geometry*. Algorithms and Computation in Mathematics, vol. 10, 2nd edn. Springer, Berlin (2006)
13. Bihan, F.: Viro method for the construction of real complete intersections. *Adv. Math.* **169**(2), 177–186 (2002)
14. Blekherman, G., Parrilo, P.A., Thomas, R.R. (eds.): *Semidefinite Optimization and Convex Algebraic Geometry*. MOS-SIAM Series on Optimization, vol. 13. SIAM, Philadelphia (2013)
15. Butkovič, P.: *Max-linear Systems: Theory and Algorithms*. Springer Monographs in Mathematics. Springer, London (2010)
16. Cluckers, R., Lipshitz, L., Robinson, Z.: Analytic cell decomposition and analytic motivic integration. *Ann. Sci. Éc. Norm. Supér.* **39**(4), 535–568 (2006)
17. Connes, A., Consani, C.: The hyperring of adèle classes. *J. Number Theory* **131**(2), 159–194 (2011)
18. Denef, J.: The rationality of the Poincaré series associated to the p -adic points on a variety. *Invent. Math.* **77**(1), 1–23 (1984)
19. Denef, J.: p -adic semi-algebraic sets and cell decomposition. *J. Reine Angew. Math.* **369**, 154–166 (1986)
20. Develin, M., Yu, J.: Tropical polytopes and cellular resolutions. *Exp. Math.* **16**(3), 277–291 (2007)
21. van den Dries, L., Speissegger, P.: The real field with convergent generalized power series. *Trans. Am. Math. Soc.* **350**(11), 4377–4421 (1998)
22. Einsiedler, M., Kapranov, M., Lind, D.: Non-archimedean amoebas and tropical varieties. *J. Reine Angew. Math.* **601**, 139–157 (2006)
23. Engler, A.J., Prestel, A.: *Valued Fields*. Springer Monographs in Mathematics. Springer, Berlin (2005)
24. Gärtner, B., Matoušek, J.: *Approximation Algorithms and Semidefinite Programming*. Springer, Heidelberg (2012)
25. Helton, J.W., Nie, J.: Sufficient and necessary conditions for semidefinite representability of convex hulls and sets. *SIAM J. Optim.* **20**(2), 759–791 (2009)
26. Jell, P., Scheiderer, C., Yu, J.: Real tropicalization and analytification of semialgebraic sets. [arXiv:1810.05132](https://arxiv.org/abs/1810.05132) (2018)
27. Maclagan, D., Sturmfels, B.: *Introduction to Tropical Geometry*. Graduate Studies in Mathematics, vol. 161. American Mathematical Society, Providence (2015)
28. Marker, D.: *Model Theory: An Introduction*. Graduate Texts in Mathematics, vol. 217. Springer, New York (2002)
29. Markwig, T.: A field of generalized Puiseux series for tropical geometry. *Rend. Sem. Mat. Univ. Politec. Torino* **68**(1), 79–92 (2010)
30. Pas, J.: Uniform p -adic cell decomposition and local zeta functions. *J. Reine Angew. Math.* **399**, 137–172 (1989)
31. Pas, J.: Cell decomposition and local zeta functions in a tower of unramified extensions of a p -adic field. *Proc. Lond. Math. Soc.* **60**(3), 37–67 (1990)
32. Pas, J.: On the angular component map modulo P . *J. Symb. Log.* **55**(3), 1125–1129 (1990)
33. Ramana, M.V.: An exact duality theory for semidefinite programming and its complexity implications. *Math. Program.* **77**(2), 129–162 (1997)
34. Ribenboim, P.: Fields: algebraically closed and others. *Manusc. Math.* **75**(2), 115–150 (1992)
35. Scheiderer, C.: Spectrahedral shadows. *SIAM J. Appl. Algebra Geom.* **2**(1), 26–44 (2018)
36. Skomra, M.: *Tropical spectrahedra: Application to semidefinite programming and mean payoff games*. PhD thesis, Université Paris-Saclay (2018). <https://pastel.archives-ouvertes.fr/tel-01958741>
37. Sturmfels, B.: Viro’s theorem for complete intersections. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **21**(3), 377–386 (1994)
38. Tent, K., Ziegler, M.: *A Course in Model Theory*. Lecture Notes in Logic, vol. 40. Cambridge University Press, Cambridge (2012)
39. Vinnikov, V.: LMI representations of convex semialgebraic sets and determinantal representations of algebraic hypersurfaces: past, present, and future. In: Dym, H., de Oliveira, M.C., Putinar, M.

- (eds.) *Mathematical Methods in Systems, Optimization, and Control. Operator Theory: Advances and Applications*, vol. 222, pp. 325–349. Springer, Basel (2012)
40. Viro, O.Ya.: Real plane algebraic curves: constructions with controlled topology. *Leningrad Math. J.* **1**(5), 1059–1134 (1990)
 41. Viro, O.: Hyperfields for tropical geometry I. Hyperfields and dequantization. [arXiv:1006.3034](https://arxiv.org/abs/1006.3034) (2010)
 42. Weispfenning, V.: Quantifier elimination and decision procedures for valued fields. In: Müller, G.H., Richter, M.M. (eds.) *Models and Sets. Lecture Notes in Mathematics*, vol. 1103, pp. 419–472. Springer, Berlin (1984)
 43. Yu, J.: Tropicalizing the positive semidefinite cone. *Proc. Am. Math. Soc.* **143**(5), 1891–1895 (2015)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.