

# LIST OF TODOS

■	I made the inequality a corollary here . . . . .	7
■	but the theorem is marked Cor13a? . . . . .	8
■	It may be appropriate here to indicate that $\text{ar}(K) < \text{mr}(K)$ some- times (maybe in previous subsection), and talk about the 2-cable of the trefoil that does not have $\text{ar}(K, \mathbb{C}) = 4$ . . . . .	8
■	figure out this two tensor products nonsense . . . . .	8
■	how do I bring in equations to fit margins? . . . . .	8
■	make consistent throughout paper . . . . .	9
■	does this need justification? . . . . .	13
■	do I need to explain what I'm doing here? . . . . .	13
■	check this . . . . .	14
■	check . . . . .	14
■	check . . . . .	15
■	check . . . . .	15
■	is this clear/can it be shortened? . . . . .	16

# AUGMENTATION RANK OF SATELLITES WITH BRAID PATTERN

DAVID R. HEMMINGER AND CHRISTOPHER R. CORNWELL

ABSTRACT. Given a knot  $K$  in  $S^3$ , a question raised by Cappell and Shaneson asks if the meridional rank of  $K$  equals the bridge number of  $K$ . Using augmentations in knot contact homology we consider the persistence of equality between these two invariants under satellite operations on  $K$  with a braid pattern. In particular, we answer the question in the affirmative for a large class of iterated torus knots.

## 1. INTRODUCTION

Let  $K$  be an oriented knot in  $S^3$  and denote by  $\pi_K$  the fundamental group of the knot complement  $\overline{S^3 \setminus n(K)}$ . We call an element of  $\pi_K$  a *meridian* if it is represented by the oriented boundary of a disc  $D$  embedded in  $S^3$  such that  $D$  intersects  $K$  geometrically once on its interior. The group  $\pi_K$  is generated by meridians; the *meridional rank* of  $K$ , written  $\text{mr}(K)$ , is the minimal size of a generating set containing only meridians.

Choose a height function  $h : S^3 \rightarrow \mathbb{R}$ . The *bridge number* of  $K$ , denoted  $b(K)$ , is the minimal number of local maxima of  $h|_K$  among all realizations of  $K$ . Equivalently, call a 2-sphere  $S \subset S^3$  a *bridge sphere* if each ball in  $S^3 \setminus S$  intersects  $K$  in some number  $b(S, K)$  of trivial arcs. The minimum of  $b(S, K)$  over all bridge spheres and all realizations of  $K$  equals  $b(K)$ .

A basic argument shows that  $\text{mr}(K) \leq b(K)$  for any  $K \subset S^3$ . Whether this bound is, in fact, equality for all knots is an open question asked by Cappell and Shaneson [Kir95, Prob. 1.11]. Equality is known to hold for some families of knots due to work of various authors.

In this paper we study the behavior of augmentations, which arise in the study of knot contact homology, upon taking satellites of  $K$  with a braid pattern. As we explain below, augmentations provide a lower bound on  $\text{mr}(K)$  which is sometimes large enough to force  $\text{mr}(K) = b(K)$ . Our main result shows that for many braid patterns this equality will persist upon taking a satellite if augmentations forced the equality for  $K$ .

More precisely, suppose that  $B \in B_k$  and  $B' \in B_p$  are two braids (here  $B_n$  denotes Artin's group of  $n$ -braids) and that  $K$  is the braid closure of  $B$  (see Figure 2 below). Denote by  $B^{(p)}$  the braid in  $B_{kp}$  which consists of each strand of  $B$  being replaced by  $p$  parallel copies of that strand (in the blackboard framing). Also, include  $B'$  into  $B_{kp}$  by juxtaposing it with an

identity braid on  $(k - 1)p$  strands. The *braid satellite* of  $K$  associated to  $B, B'$  is the braid closure of  $B^{(p)}B'$ , denoted  $K(B, B')$ .

Augmentations of  $K$  have an associated *rank*, and we write  $\text{ar}(K)$  for the maximum rank among all augmentations of  $K$ . This number is well-defined and we have the following inequalities (see Section 2.2)

$$(1) \quad \text{ar}(K) \leq \text{mr}(K) \leq b(K)$$

To minimize notation, we will write  $\text{ar}(B)$  to denote the augmentation rank of the braid closure of a braid  $B$ . We show the following.

**Theorem 1.1.** *If  $B$  is a  $k$ -braid such that  $\text{ar}(B) = k$  and  $B'$  a  $p$ -braid such that  $\text{ar}(B') = p$  then  $\text{ar}(K(B, B')) = kp$ .*

**Remark 1.2.** Our theorem shows that if  $B, B'$  have largest possible augmentation rank (equal to their index) then the augmentation rank acts multiplicatively under braid satellites. It is natural to ask if this multiplicative property holds under weaker assumptions on  $B, B'$ . Braid stabilization provides a way to construct counterexamples to augmentation rank being multiplicative, even when the bridge number does multiply.

A corollary to Theorem 1.1 involves Cappell and Shaneson's question for iterated torus knots. Let  $\mathbf{p} = (p_1, \dots, p_n)$  and  $\mathbf{q} = (q_1, \dots, q_n)$  be integral vectors. We write  $T(\mathbf{p}, \mathbf{q})$  for the  $(\mathbf{p}, \mathbf{q})$  *iterated torus knot*, defined as follows. Define  $T(\mathbf{p}, \mathbf{q})$  inductively so that, if  $\hat{\mathbf{p}}, \hat{\mathbf{q}}$  are the truncated lists obtained from  $\mathbf{p}, \mathbf{q}$  by removing the last integer in each, then  $T(\mathbf{p}, \mathbf{q})$  is the  $(p_n, q_n)$ -cable of  $T(\hat{\mathbf{p}}, \hat{\mathbf{q}})$ . By convention  $T(\emptyset, \emptyset)$  is the unknot.

The knot (or link)  $T(\mathbf{p}, \mathbf{q})$  is only well-defined if a framing convention is chosen at each stage of cabling. The traditional choice is to take the Seifert framing at each stage. In the statement of the corollary below, we use a different convention which is suited to the braid satellite construction (see Section ??).

**Corollary 1.3.** *Given integral vectors  $\mathbf{p}$  and  $\mathbf{q}$ , suppose that  $|p_i| < |q_i|$  and  $\gcd(p_i, q_i) = 1$  for each  $1 \leq i \leq n$ . Then  $\text{ar}(T(\mathbf{p}, \mathbf{q})) = p_1 p_2 \dots p_n$ . Moreover,  $\text{mr}(T(\mathbf{p}, \mathbf{q})) = b(T(\mathbf{p}, \mathbf{q}))$  for all such  $\mathbf{p}$  and  $\mathbf{q}$ .*

The paper is organized as follows. In Section 2 we give the needed background in knot contact homology and discuss the rank of augmentations and the relationship to the meridional rank. In Section 2.3 we review some techniques that will be used in the proof of Theorem 1.1. We also discuss there the difficulty of generalizing Theorem 1.1 to braid satellites of knots with smaller augmentation rank. Section 3 is mainly devoted to the proof of Theorem 1.1.

## 2. BACKGROUND

We review in Section 2.1 the construction of  $HC_0(K)$  from the viewpoint of the combinatorial knot DGA, which was first defined in [Ng08]; our conventions are those given in [Ng12]. In Section 2.2 we discuss augmentations

in knot contact homology and their rank, which gives a lower bound on the meridional rank of the knot group useful for studying the relation between meridional rank and bridge number. Finally, in Section 2.3 is a discussion of techniques from [Cor13a] that we use to calculate the augmentation rank.

Throughout the paper we denote by  $B_n$  the group of  $n$ -braids. We orient  $n$ -braids from left to right, labeling the strands  $1, \dots, n$ , with 1 the topmost and  $n$  the bottommost strand. We work with Artin's generators  $\{\sigma_i^\pm, i = 1, \dots, n-1\}$  of  $B_n$ , where in  $\sigma_i$  only the  $i$  and  $i+1$  strands interact, and they cross once in the manner depicted in Figure 1. Given a braid  $B$  the




FIGURE 1. Generators of  $B_n$

braid closure  $\widehat{B}$  of  $B$  is the link obtained as shown in Figure 2. The *writhe* (or algebraic sum) of  $B$ , denoted  $\omega(B)$ , is the sum of the exponents in a factorization of  $B$  as a product of the  $\sigma_i, i = 1, \dots, n-1$ .




FIGURE 2. The braid closure of  $B$

**2.1. Knot contact homology.** We review the construction of the combinatorial knot DGA of Ng (in fact, we discuss only the degree zero part as this will suffice for our purposes). This DGA was defined in order to be a calculation of knot contact homology and was shown to be so in [EENS13] (see [Ng12] for more details). Let  $\mathcal{A}_n$  be the noncommutative unital algebra over  $\mathbb{Z}$  freely generated by  $a_{ij}, 1 \leq i \neq j \leq n$ . We define a homomorphism  $\phi : B_n \rightarrow \text{Aut } \mathcal{A}_n$  by defining it on the generators of  $B_n$ :

$$(2) \quad \phi_{\sigma_k} : \begin{cases} a_{ij} \mapsto a_{ij} & i, j \neq k, k+1 \\ a_{k+1,i} \mapsto a_{ki} & i \neq k, k+1 \\ a_{i,k+1} \mapsto a_{ik} & i \neq k, k+1 \\ a_{k,k+1} \mapsto -a_{k+1,k} \\ a_{k+1,k} \mapsto -a_{k,k+1} \\ a_{ki} \mapsto a_{k+1,i} - a_{k+1,k}a_{ki} & i \neq k, k+1 \\ a_{ik} \mapsto a_{i,k+1} - a_{ik}a_{k,k+1} & i \neq k, k+1 \end{cases}$$

Let  $\iota: B_n \rightarrow B_{n+1}$  be the inclusion  $\sigma_i \mapsto \sigma_i$  so that the  $(n+1)$  strand does not interact with those from  $B \in B_n$ , and define  $\phi_B^* \in \text{Aut } \mathcal{A}_{n+1}$  by  $\phi_B^* = \phi_B \circ \iota$ . We then define the  $n \times n$  matrices  $\Phi_B^L$  and  $\Phi_B^R$  with entries in  $\mathcal{A}_n$  by

$$\begin{aligned}\phi_B^*(a_{i,n+1}) &= \sum_{j=1}^n (\Phi_B^L)_{ij} a_{j,n+1} \\ \phi_B^*(a_{n+1,i}) &= \sum_{j=1}^n a_{n+1,j} (\Phi_B^R)_{ji}\end{aligned}$$

Finally, let  $R_0$  be the Laurent polynomial ring  $\mathbb{Z}[\lambda^{\pm 1}, \mu^{\pm 1}]$  and define matrices  $\mathbf{A}$  and  $\mathbf{\Lambda}$  over  $R_0$  by

$$(3) \quad \mathbf{A}_{ij} = \begin{cases} a_{ij} & i < j \\ -\mu a_{ij} & i > j \\ 1 - \mu & i = j \end{cases}$$

$$(4) \quad \mathbf{\Lambda} = \text{diag}[\lambda \mu^{\omega(B)}, 1, \dots, 1].$$


**Definition 2.1.** Suppose that  $K$  is the closure of  $B \in B_n$ . Define  $\mathcal{I} \subset \mathcal{A}_n \otimes R_0$  to be the ideal generated by the entries of  $\mathbf{A} - \mathbf{\Lambda} \cdot \Phi_B^L \cdot \mathbf{A}$  and  $\mathbf{A} - \mathbf{A} \cdot \Phi_B^R \cdot \mathbf{\Lambda}^{-1}$ . The *degree zero homology of the combinatorial knot DGA* is  $\text{HC}_0(K) = (\mathcal{A}_n \otimes R_0)/\mathcal{I}$ .

It was shown in [Ng08] that the isomorphism class of  $\text{HC}_0(K)$  is unchanged by the Markov moves on  $B$ , hence  $\text{HC}_0(K)$  is an invariant of the knot  $K$ . We only consider  $\text{HC}_0(K)$  here, but there is a larger invariant, the differential graded algebra discussed in [Ng12].

The proofs in Section 3 require a number of computations of  $\phi_B$  for particular braids  $B$ . Such computations are greatly benefited by an alternate description of the map  $\phi_B$ , which we now give and will use without comment in Section 3.

Let  $D$  be a flat disk, to the right of  $B$ , with  $n$  points (punctures) where it intersects  $K = \widehat{B}$  (see Figure 3). We assume the  $n$  punctures of  $D$  to be collinear, on a line that separates  $D$  into upper and lower half-disks. Denote by  $c_{ij}$  the isotopy class (fixing endpoints) of a path that is contained in the upper half-disk of  $D$ , with initial endpoint on the  $i^{\text{th}}$  strand and terminal endpoint on the  $j^{\text{th}}$  strand.

Consider  $B$  as a mapping class and let  $B \cdot c_{ij}$  denote the isotopy class of the path to which  $c_{ij}$  is sent. If  $D$ , as viewed from the left (as pictured), is oriented as the plane then  $\sigma_k$  acts by rotating the  $k$ - and  $(k+1)$ -punctures an angle of  $\pi$  about their midpoint in counter-clockwise fashion. Consider the *algebra of paths* over  $\mathbb{Z}$  generated by isotopy classes of paths in  $D$  with endpoints on punctures, modulo the relation in Figure 4 (paths depicted there are understood to agree outside the neighborhood of the puncture

FIGURE 3. Cord  $c_{ij}$  of  $K = \widehat{B}$ 

shown). It was shown in [Ng05] that the algebra map to  $\mathcal{A}_n$  defined by  $F(c_{ij}) = a_{ij}$  if  $i < j$ , and  $F(c_{ij}) = -a_{ij}$  if  $i > j$  satisfies  $F(B \cdot c_{ij}) = \phi_B(F(c_{ij}))$ .

$$[\text{curved arrow}] = [\text{curved arrow with dot}] - [\text{straight arrow with dot}] \cdot [\text{straight arrow}]$$

FIGURE 4. Relation in the algebra of paths

Let  $\text{perm} : B_n \rightarrow S_n$  denote the homomorphism from  $B_n$  to the symmetric group sending  $\sigma_k$  to the simple transposition interchanging  $k, k+1$ . We will make use of the following.

**Lemma 2.2.** *For some  $B \in B_n$  and  $1 \leq i \neq j \leq n$ , consider  $(\Phi_B^L)_{ij} \in \mathcal{A}_n$  as a polynomial expression in the (non-commuting) variables  $\{a_{kl}, 1 \leq k \neq l \leq n\}$ . Writing  $i' = \text{perm}(B)(i)$ , every monomial in  $(\Phi_B^L)_{ij}$  is a constant times  $a_{i'i_1} a_{i_1 i_2} \dots a_{i_{l-1} j}$  for some  $l \geq 0$ , the monomial being a constant if  $l = 0$  and only if  $i' = j$ .*

*Proof.* Include  $B$  in  $B_{n+1}$  and consider the isotopy class of the path  $B \cdot c_{i,n+1}$  which begins at  $i'$  and ends at  $n+1$  (as  $B$  does not interact with the  $(n+1)$  strand). Applying the relation in Figure 4 to the path equates it with a sum (or difference) of another path with the same endpoints and a product of two paths, the first beginning at  $i'$  and the other ending at  $n+1$ . A finite number of applications of this relation allows one to express the path as a polynomial in the  $c_{kl}, 1 \leq k \neq l \leq n$  where each monomial has the form  $c_{i'i_1} \dots c_{i_{l-1} j}$ . The result then follows from the fact that  $\phi_B(a_{i,n+1}) = \phi_B(F(c_{i,n+1})) = F(B \cdot c_{i,n+1})$ .

Alternatively, the statement follows from noting that (2) defining  $\phi_{\sigma_k}$  has the desired property and that  $\phi : B_n \rightarrow \text{Aut}(\mathcal{A}_n)$  is a homomorphism.  $\square$

**2.2. Augmentations and augmentation rank.** Let  $S$  be a ring with 1, and consider it a differential graded algebra supported in grading 0, with trivial differential. Augmentations of  $(\mathcal{A}, \partial)$  are DGA maps  $(\mathcal{A}, \partial) \rightarrow (S, 0)$ . For our setting, if  $B \in B_n$  is a braid representative of  $K$ , such a map corresponds precisely to a homomorphism  $\epsilon : \mathcal{A}_n \otimes R_0 \rightarrow \mathbb{C}$  such that  $\epsilon$  sends each generator of  $\mathcal{I}$  to zero (see Definition 2.1).

**Definition 2.3.** Suppose that  $K$  is the closure of  $B \in B_n$ . An *augmentation* of  $K$  is a homomorphism  $\epsilon : \mathcal{A}_n \otimes R_0 \rightarrow \mathbb{C}$  such that each element of  $\mathcal{I}$  is sent by  $\epsilon$  to zero.

A correspondence between augmentations and particular representations of the knot group  $\pi_K$  were studied in [Cor13a]. Recall that  $\pi_K$  is generated by meridians. We may fix a meridian  $m$  and generate  $\pi_K$  by conjugates of  $m$ .

**Definition 2.4.** For any integer  $r \geq 1$ , a homomorphism  $\rho : \pi_K \rightarrow \mathrm{GL}_r \mathbb{C}$  is a *KCH representation* if there is a meridian  $m$  of  $K$  such that  $\rho(m)$  is diagonalizable and has an eigenvalue of 1 with multiplicity  $r - 1$ . We call  $\rho$  a *KCH irrep* if it is irreducible.

In [Ng08], Ng describes an isomorphism between  $HC_0(K)$  and an algebra constructed from elements of  $\pi_K$ . As discussed in [Ng12] a KCH representation  $\rho : \pi_K \rightarrow \mathrm{GL}_r \mathbb{C}$  induces an augmentation  $\epsilon_\rho$  of  $K$ . Given an augmentation, the first author showed how to construct a KCH representation that induces it. In fact, we have the following rephrasing of results from [Cor13a].

**Theorem 2.5** ([Cor13a]). *Let  $\epsilon : \mathcal{A}_n \otimes R_0 \rightarrow \mathbb{C}$  be an augmentation with  $\epsilon(\mu) \neq 1$ . There is a KCH irrep  $\rho : \pi_K \rightarrow \mathrm{GL}_r \mathbb{C}$  such that  $\epsilon_\rho = \epsilon$ . Furthermore, for any KCH irrep  $\rho : \pi_K \rightarrow \mathrm{GL}_r \mathbb{C}$  such that  $\epsilon_\rho = \epsilon$ ,  $r$  equals the rank of  $\epsilon(\mathbf{A})$ .*

Considering Theorem 2.5 we make the following definition.

**Definition 2.6.** The *rank* of an augmentation  $\epsilon : \mathcal{A}_n \otimes R_0 \rightarrow \mathbb{C}$  with  $\epsilon(\mu) \neq 1$  is the rank of  $\epsilon(\mathbf{A})$ . Given a knot  $K$ , the *augmentation rank* of  $K$ , denoted  $\mathrm{ar}(K)$ , is the maximum rank among augmentations of  $K$ .

**Remark 2.7.** The augmentation rank can be defined for target rings other than  $\mathbb{C}$ , but this paper only considers augmentations as in Definition 2.3.

It is the case that  $\mathrm{ar}(K)$  is well-defined. That is, given  $K$  there is a bound on the maximal rank of an augmentation of  $K$ .

**Theorem 2.8** ([Cor13b]). *Given a knot  $K \subset S^3$ , if  $g_1, \dots, g_d$  are meridians that generate  $\pi_K$  and  $\rho : \pi_K \rightarrow \mathrm{GL}_r \mathbb{C}$  is a KCH irrep then  $r \leq d$ .*

As in the introduction, if we denote the meridional rank of  $\pi_K$  by  $\mathrm{mr}(K)$ , then Theorem 2.8 implies that  $\mathrm{ar}(K) \leq \mathrm{mr}(K)$ . In addition, the geometric quantity  $b(K)$  called the bridge index of  $K$  is never less than  $\mathrm{mr}(K)$ . Thus we have the following corollary:

**Corollary 2.9** ([Cor13b]). *Given a knot  $K \subset S^3$ ,*

$$\mathrm{ar}(K) \leq \mathrm{mr}(K) \leq b(K)$$

I made the inequality a corollary here

As a result, to verify for  $K$  that  $\mathrm{mr}(K) = b(K)$  it suffices to find an augmentation of  $K$  with rank equal to  $b(K)$ . As we discuss in the next section, we will concern ourselves in this paper with a setting where  $\mathrm{ar}(K) = n$  and there is a braid  $B \in B_n$  which closes to  $K$ . This is a special situation, since  $b(K)$  is strictly less than the braid index for many knots.

**2.3. Finding augmentations.** The following theorem concerns the behavior of the matrices  $\Phi_B^L$  and  $\Phi_B^R$  under the product in  $B_n$ . It is an essential tool for studying  $HC_0(K)$  and will be central to our arguments.

**Theorem 2.10** ([Ng05], Chain Rule). *Let  $B, B'$  be braids in  $B_n$ . Then  $\Phi_{BB'}^L = \phi_B(\Phi_{B'}^L) \cdot \Phi_B^L$  and  $\Phi_{BB'}^R = \Phi_B^R \cdot \phi_B(\Phi_{B'}^R)$ .*

The main result of this paper concerns augmentations with rank equal to the braid index of the knot  $K$ . Define the diagonal matrix  $\Delta(B) = \text{diag}[(-1)^{w(B)}, 1, \dots, 1]$ . The following statement follows from results in [Cor13b, Section 5].

but the theorem is marked Cor13a?

**Theorem 2.11** ([Cor13a]). *If  $K$  is the closure of  $B \in B_n$  and has a rank  $n$  augmentation  $\epsilon : \mathcal{A}_n \otimes R_0 \rightarrow \mathbb{C}$ , then*

$$(5) \quad \epsilon(\Phi_B^L) = \Delta(B) = \epsilon(\Phi_B^R).$$

*Furthermore, any homomorphism  $\epsilon : \mathcal{A}_n \rightarrow \mathbb{C}$  which satisfies (5) can be extended to  $\mathcal{A}_n \otimes R_0$  to produce a rank  $n$  augmentation of  $K$ .*

The proof of Theorem 1.1 relies on this characterization of rank  $n$  augmentations. That is, given a braid  $B \in B_k$  with a rank  $k$  augmentation  $\epsilon : \mathcal{A}_k \rightarrow \mathbb{C}$ , and  $B' \in B_p$  with a rank  $p$  augmentation  $\epsilon' : \mathcal{A}_p \rightarrow \mathbb{C}$  we show that  $K(B, B')$  has a rank  $kp$  augmentation by defining a homomorphism  $\psi : \mathcal{A}_{kp} \rightarrow \mathcal{A}_k \otimes \mathcal{A}_p$  so that  $(\epsilon \otimes \epsilon') \circ \psi$  satisfies (5) for the braid  $B^{(p)}B'$ .

There is a symmetry on the matrices  $\Phi_B^L$  and  $\Phi_B^R$  that is relevant to the study of augmentations in this setting. Define an involution  $x \mapsto \bar{x}$  on  $\mathcal{A}_n$  (termed *conjugation*) as follows: first set  $\bar{a}_{ij} = a_{ji}$ ; then, for any  $x, y \in \mathcal{A}_n$ , define  $\overline{xy} = \bar{y}\bar{x}$  and extend the operation linearly to  $\mathcal{A}_n$ . We have the following symmetry.

**Theorem 2.12** ([Ng05], Prop. 6.2). *For a matrix of elements in  $\mathcal{A}_n$ , let  $\overline{M}$  be the matrix such that  $(\overline{M})_{ij} = \overline{M_{ij}}$ . Then for  $B \in B_n$ ,  $\Phi_B^R$  is the transpose of  $\overline{\Phi_B^L}$ .*

It may be appropriate here to indicate that  $\text{ar}(K) < \text{mr}(K)$  sometimes (maybe in previous subsection), and talk about the 2-cable of the trefoil that does not have  $\text{ar}(K, \mathbb{C}) = 4$

### 3. MAIN RESULT

figure out this two tensor products nonsense

how do I bring in equations to fit margins?

We begin in Section 3.1 with our main theorem. The proof relies upon an intermediate result, Proposition 3.1, which is shown in the subsequent Section 3.2 along with some supporting lemmas.

**3.1. Proof of main result.** In this section we prove our main result, which we now recall. The notation of Theorem 1.1 will be used throughout Section 3.



**Theorem 1.1.** Let  $B \in B_k$  have augmentation rank  $k$ , and let  $B'' \in B_p$  have augmentation rank  $p$ . If  $B'$  is the braid  $B''$  included into  $B_{pk}$ , then  $B^{(p)}B'$  has augmentation rank  $pk$ .

Theorem 1.1 is proved by defining an algebra map  $\psi: \mathcal{A}_{pk} \rightarrow \mathcal{A}_k \otimes \mathcal{A}_p$  such that  $\psi(\Phi_{B^{(p)}B'}^L)$  factors suitably across the tensor product. This is the content of Proposition 3.1, which follows from Lemmas 3.2 and 3.4, the former following from a calculation (Lemma 3.5) and the latter depending on the former.

For each  $1 \leq i \leq pk$  let  $q_i \geq 0$  and  $0 < r_i \leq p$  be the integers such that  $i = q_i p + r_i$ . We define the map  $\psi: \mathcal{A}_{pk} \rightarrow \mathcal{A}_k \otimes \mathcal{A}_p$  on generators as follows

$$\psi(a_{ij}) = \begin{cases} 1 \otimes a_{r_i r_j} & : q_i = q_j \\ a_{q_i+1, q_j+1} \otimes 1 & : r_i = r_j \\ 0 & : (q_i - q_j)(r_i - r_j) < 0 \\ a_{q_i+1, q_j+1} \otimes a_{r_i r_j} & : (q_i - q_j)(r_i - r_j) > 0 \end{cases}$$

Note that  $\psi(a_{ij}) \in 1 \otimes \mathcal{A}_p$  and is nonzero if and only if  $q_i = q_j$ , and that  $\psi(\overline{a_{ij}}) = \overline{\psi(a_{ij})}$ . This homomorphism gives us a way of relating  $\Phi_{B^{(p)}}^L$  to  $\Phi_B^L$  via the following proposition.

**Proposition 3.1.** For any braid  $B$ ,  $\psi(\Phi_{B^{(p)}}^L) = ((\Phi_B^L)_{ij} \otimes 1) \otimes I_p$  and  $\psi(\Phi_{B^{(p)}}^R) = ((\Phi_B^R)_{ij} \otimes 1) \otimes I_p$

make consistent throughout paper

Note that here we mean the tensor product of  $\Phi_B^L$  and  $I_p$  as matrices, not as linear maps, while the tensor product of  $(\Phi_B^L \otimes I_p)_{ij}$  and 1 is a tensor product of algebra elements, so that if we divide the matrix  $\psi(\Phi_{B^{(p)}}^L)$  into  $k^2 p \times p$  blocks, the  $ij$ th block is  $(\Phi_B^L)_{ij} I_p$ .

*Proof of Theorem 1.1.* By Theorem 2.11 it suffices to prove that there exists a homomorphism  $\epsilon: \mathcal{A}_{pk} \rightarrow \mathbb{C}$  such that  $\epsilon(\Phi_{B^{(p)}B'}^L) = \epsilon(\Phi_{B^{(p)}B'}^R) = \Delta(B^{(p)}B')$ . Additionally, our assumptions imply that there exist augmentations  $\epsilon_k: \mathcal{A}_k \otimes R_0 \rightarrow \mathbb{C}$  and  $\epsilon_p: \mathcal{A}_p \otimes R_0 \rightarrow \mathbb{C}$  such that  $\epsilon_k(\Phi_B^L) = \epsilon_k(\Phi_B^R) = \Delta(B)$  and  $\epsilon_p(\Phi_{B'}^L) = \epsilon_p(\Phi_{B'}^R) = \Delta(B')$ .

We will define a homomorphism  $\delta: \mathcal{A}_p \rightarrow \mathbb{C}$  such that  $\delta = \pm \epsilon_p$ , the sign depending on the parity of  $w(B)$  and  $p$ . If  $\pi: \mathbb{C} \otimes \mathbb{C} \rightarrow \mathbb{C}$  is the homomorphism determined by  $\pi(a \otimes b) = ab$ , for  $a, b \in \mathbb{C}$ , then we set  $\epsilon = \pi \circ (\epsilon_k \otimes \delta) \circ \psi$ .

The Chain Rule theorem gives that

$$(6) \quad \pi \circ (\epsilon_k \otimes \delta) \circ \psi(\Phi_{B^{(p)}B'}^L) = \pi \circ (\epsilon_k \otimes \delta) \psi(\phi_{B^{(p)}}(\Phi_{B'}^L)) \psi(\Phi_{B^{(p)}}^L)$$

Note that since the non zero or one entries of  $\Phi_{B'}^L$  are products of  $a_{ij}$  where  $i < j \leq p$ ,  $\phi_{B^{(p)}}$  takes each of the  $a_{ij}$ 's in these products to  $a_{i+mp, j+mp}$  for some  $0 \leq m < k$  (this can be seen easily by using the alternate description

of  $\phi_{B^{(p)}}$  detailed in Section 2.1 and using that  $B^{(p)}$  is a product of braids  $\Sigma_l^{(p)}$ ). We have that  $\psi$  takes  $a_{i+mp, j+mp}$  to  $1 \otimes a_{ij}$ , however, so

$$\psi(\phi_{B^{(p)}}(\Phi_{B'}^L)) = \left(1 \otimes (\Phi_{B'}^L)_{ij}\right)$$

By Proposition 3.1

$$\psi(\Phi_{B^{(p)}}^L) = \left((\Phi_B^L)_{ij} \otimes 1\right) \otimes I_p = \left((\Phi_B^L \otimes I_p)_{ij} \otimes 1\right).$$

Returning to the right hand side of (6)

$$\begin{aligned} \pi \circ (\epsilon_k \otimes \delta) \left( \psi(\phi_{B^{(p)}}(\Phi_{B'}^L)) \psi(\Phi_{B^{(p)}}^L) \right) &= \pi \circ (\epsilon_k \otimes \delta) \left( \left(1 \otimes (\Phi_{B'}^L)_{ij}\right) \left((\Phi_B^L \otimes I_p)_{ij} \otimes 1\right) \right) \\ &= \delta(\Phi_{B'}^L) (\Delta(B) \otimes I_p). \end{aligned}$$

We are done if  $\delta$  may be defined so that  $\delta(\Phi_{B'}^L) (\Delta(B) \otimes I_p) = \Delta(B^{(p)} B')$ . When  $w(B)$  is even  $w(B^{(p)})$  is also, and further  $\Delta(B) = I_k$ . Letting  $\delta = \epsilon_p$  makes

$$\delta(\Phi_{B'}^L) (\Delta(B) \otimes I_p) = \epsilon_p(\Phi_{B'}^L) = \Delta(B') = \Delta(B^{(p)} B').$$

Suppose that  $w(B)$  is odd. We define  $g: \{1, \dots, p\} \rightarrow \{\pm 1\}$  as follows. Let  $x_1 = 1$ , and  $x_l = \text{perm}(B')(x_{l-1})$  for  $1 < l \leq p$ . Since the first  $p$  strands of  $B'$  close to a knot,  $\text{perm}(B')$  is given by the  $p$ -cycle  $(x_1 x_2 \dots x_p)$ . If  $p$  is even, we let  $g(x_1) = 1$ , and  $g(x_l) = -g(x_{l-1})$  for  $1 < l \leq p$ . If  $p$  is odd, let  $g(x_1) = g(x_2) = 1$  and  $g(x_l) = -g(x_{l-1})$  for  $2 < l \leq p$ .

Define  $\delta: \mathcal{A}_p \rightarrow \mathbb{C}$  by setting  $\delta(a_{ij}) = g(i)g(j)\epsilon_k(a_{ij})$  for  $1 \leq i \neq j \leq p$ . Fix  $i, j$  and consider a monomial  $M$  in  $(\Phi_{B'}^L)_{ij}$ , which is constant if  $i > p$  or  $j > p$  so that  $\delta(M)$  is defined. If  $i, j \leq p$ , then writing  $i' = \text{perm}(B')(i)$ , Proposition 2.2 implies  $M = c_{ij} a_{i', j_1} a_{j_1, j_2} \dots a_{j_m, j}$  for some  $j_1, \dots, j_m \in \{1, \dots, p\}$ , possibly being constant only if  $i' = j$ , implying that

$$\delta(M) = g(i')g(j) \left( \prod_{k=1}^m g(j_k)^2 \right) \epsilon_p(M) = g(i')g(j)\epsilon_p(M).$$

Note, when  $M$  is constant then  $\delta(M) = M = g(i')g(j)\epsilon_p(M)$  since  $i' = j$ . Since this holds for each monomial, we have that

$$\delta\left((\Phi_{B'}^L)_{ij}\right) = g(i')g(j)\epsilon_p\left((\Phi_{B'}^L)_{ij}\right).$$

When  $p$  is even,  $w(B^{(p)})$  is also even and so the opposite parity of  $w(B)$ . Our definition of  $g$  gives  $\delta((\Phi_{B'}^L)_{ii}) = -\epsilon((\Phi_{B'}^L)_{ii})$  for  $i \leq p$ . Thus

$$\delta(\Phi_{B'}^L) = \begin{pmatrix} (-1)^{w(B')+1} & 0 & 0 \\ 0 & -I_{p-1} & 0 \\ 0 & 0 & I_{(k-1)p} \end{pmatrix}$$

and therefore

$$\delta(\Phi_{B'}^L) (\Delta(B) \otimes I_p) = \text{diag}[(-1)^{w(B)+w(B')+1}, 1 \dots 1] = \Delta(B^{(p)} B')$$

as desired.

When  $p$  is odd,  $w(B^{(p)})$  is odd and therefore the same parity of  $w(B)$ . Our definition of  $g$  gives that  $\delta((\Phi_{B'}^L)_{11}) = \epsilon((\Phi_{B'}^L)_{11})$  and  $\delta((\Phi_{B'}^L)_{ii}) = -\epsilon((\Phi_{B'}^L)_{ii})$  for  $1 < i \leq p$ , so

$$\delta(\Phi_{B'}^L) = \begin{pmatrix} (-1)^{w(B')} & 0 & 0 \\ 0 & -I_{p-1} & 0 \\ 0 & 0 & I_{(k-1)p} \end{pmatrix}$$

and therefore

$$\delta(\Phi_{B'}^L)(\Delta(B) \otimes I_p) = \text{diag}[(-1)^{w(B)+w(B')}, 1 \dots 1] = \Delta(B^{(p)}B')$$

as desired.

There is little difference in the proof that  $\epsilon(\Phi_{B^{(p)}}^R) = \Delta(B^{(p)}B')$ , except that monomials in  $(\Phi_{B'}^R)_{ij}$  are of the form  $c_{ij}a_{i,j_1}a_{j_1,j_2} \cdots a_{j_k,j'}$  where  $j' = \text{perm}(B')(j)$ . Applying Theorem 2.11 now completes the proof.  $\square$

**3.2. Proposition 3.1 and supporting lemmas.** We use the following two lemmas to prove Proposition 3.1. Figure 5 demonstrates an example for Lemma 3.2, showing that  $\psi(\phi_{\Sigma_2^{(2)}}(a_{24})) = \phi_{\sigma_2} \otimes \text{id}(\psi(a_{24}))$ . Note that in the figure we condense elements such as  $a_{13} \otimes 1$  to  $a_{13}$  and include products of algebra elements on a single set of points in order to make the notation cleaner.

**Lemma 3.2.**  $\psi(\phi_{\Sigma_n^{\pm(p)}}(a_{ij})) = (\phi_{\sigma_n^{\pm 1}} \otimes \text{id})(\psi(a_{ij}))$  for all  $1 \leq n < k$ ,  $1 \leq i, j \leq pk$ .

**Remark 3.3.** As the map  $B_k \rightarrow \text{Aut}(\mathcal{A}_k \otimes \mathcal{A}_p)$  given by  $B \mapsto \phi_B \otimes \text{id}$  is a homomorphism, Lemma 3.2 immediately implies that  $\psi(\phi_{B^{(p)}}(a_{ij})) = (\phi_B \otimes \text{id})(\psi(a_{ij}))$  for any  $B \in B_k$ .

**Lemma 3.4** (Base Case).  $\psi(\Phi_{\Sigma_n^{\pm(p)}}^L) = \left( (\Phi_{\sigma_n^{\pm 1}}^L)_{ij} \otimes 1 \right) \otimes I_p$

$$\begin{aligned} & \psi(\phi_{\Sigma_2^{(2)}}(\cdot \curvearrowright \cdot \cdot)) \\ &= \psi(\cdot \curvearrowright \cdot) \\ &= \psi(\cdot \curvearrowright \cdot - \cdot \curvearrowright \cdot - \cdot \curvearrowright \cdot - \cdot \curvearrowright \cdot) \\ &= 0 - \curvearrowright - 0 + \curvearrowright \\ &= \phi_{\sigma_2}(\curvearrowright \cdot) \end{aligned}$$

FIGURE 5. Computing  $\psi(\phi_{\Sigma_2^{(p)}}(a_{24}))$

*Proof of Proposition 3.1.* Let  $B = \sigma_{n_1}^{q_1} \cdots \sigma_{n_r}^{q_r}$ , where  $1 \leq n_i < k$  and  $q_i = \pm 1$ . We will prove the proposition by induction on  $r$ . The base case is already taken care of by Lemma 3.4. Suppose that the proposition holds for braids of length  $r-1$ . Let  $B' = \sigma_{n_1}^{q_1} \cdots \sigma_{n_{r-1}}^{q_{r-1}}$ . Then by the Chain Rule and Lemmas 3.2 and 3.4, we have that

$$\begin{aligned} \psi(\Phi_{B(p)}^L) &= \psi\left(\phi_{B'(p)}\left(\Phi_{\Sigma_{n_r}^{q_r(p)}}^L\right) \cdot \Phi_{B'(p)}^L\right) \\ &= (\phi_{B'} \otimes \text{id})\left(\psi\left(\Phi_{\Sigma_{n_r}^{q_r(p)}}^L\right)\right) \cdot \left((\Phi_{B'}^L)_{ij} \otimes 1\right) \otimes I_p \\ &= (\phi_{B'} \otimes \text{id})\left(\left(\left(\Phi_{\Sigma_{n_r}^{q_r(p)}}^L\right)_{ij} \otimes 1\right) \otimes I_p\right) \cdot \left((\Phi_{B'}^L)_{ij} \otimes 1\right) \otimes I_p \\ &= \left((\Phi_B^L)_{ij} \otimes 1\right) \otimes I_p \end{aligned}$$

We also have then that  $\psi(\Phi_{B(p)}^R) = \left((\Phi_B^R)_{ij} \otimes 1\right) \otimes I_p$  as well, since  $\Phi_B^R = \overline{\Phi_B^L}^t$  and  $\psi(\overline{a_{ij}}) = \overline{\psi(a_{ij})}$ .  $\square$

In the proof of Lemmas 3.2 and 3.4, we will make use of some calculations of  $\phi_B(a_{ij})$  for simple braids  $B$ . Recall that  $\tau_{m,l} = \sigma_m \sigma_{m+1} \cdots \sigma_{m+l-1}$ . It can easily be checked that for all  $1 \leq m < n$ ,  $1 \leq l \leq n-m$ ,  $i < j$

$$(7) \quad \phi_{\tau_{m,l}}(a_{ij}) = \begin{cases} a_{i+1,j+1} & : m \leq i < j < m+l \\ a_{i-l,j} & : m < m+l = i < j \\ a_{i,j-l} & : i < m < m+l = j \\ a_{i+1,j-l} & : m \leq i < j = m+l \\ a_{i,j+1} - a_{i,m} a_{m,j+1} & : i < m \leq j < m+l \\ a_{i+1,j} - a_{i+1,m} a_{m,j} & : m \leq i < m+l < j \\ a_{ij} & : \text{otherwise} \end{cases}$$

We also make the following definition

Let  $X \subseteq \{1, \dots, n\}$ , and write the elements of a subset  $Y \subseteq X$  as  $y_1 < \dots < y_k$ . Define

$$A(i, j, X) = \sum_{Y \subseteq X} (-1)^{|Y|} a_{iy_1} a_{y_1 y_2} \cdots a_{y_k j}$$

and

$$A'(i, j, X) = \sum_{Y \subseteq X} (-1)^{|Y|} a_{iy_k} a_{y_k y_{k-1}} \cdots a_{y_1 j}$$

and have the following lemma

**Lemma 3.5.** Let  $X_n^{(p)} = \{(n-1)p+1, \dots, np\}$ . We have

$$\phi_{\Sigma_n^{(p)}}(a_{ij}) = \begin{cases} a_{i-p, j-p} & : np < i < j \leq (n+1)p \\ a_{i-p, j} & : np < i \leq (n+1)p < j \\ a_{i, j-p} & : i \leq (n-1)p < np < j \leq (n+1)p \\ a_{i+p, j+p} & : (n-1)p < i < j \leq np \\ A'(i+p, j-p, X_n^{(p)}) & : (n-1)p < i \leq np < j \leq (n+1)p \\ A(i, j+p, X_n^{(p)}) & : i \leq (n-1)p < j \leq np < (n+1)p \\ A'(i+p, j, X_n^{(p)}) & : (n-1)p < i \leq np < (n+1)p < j \\ a_{ij} & : \text{otherwise} \end{cases}$$

*Proof of Lemma 3.2.* Note that if  $\psi(\phi_{\Sigma_n^{(p)}}(a_{ij})) = (\phi_{\sigma_n} \otimes \text{id})(\psi(a_{ij}))$ , then

$$\psi(a_{ij}) = \psi\left(\phi_{\Sigma_n^{(p)}}\left(\phi_{\Sigma_n^{-(p)}}(a_{ij})\right)\right) = (\phi_{\sigma_n} \otimes \text{id})\left(\psi\left(\phi_{\Sigma_n^{-(p)}}(a_{ij})\right)\right)$$

And applying  $(\phi_{\sigma_n^{-1}} \otimes \text{id})$  to both sides gives

$$\psi\left(\phi_{\Sigma_n^{-(p)}}(a_{ij})\right) = (\phi_{\sigma_n^{-1}} \otimes \text{id})\psi(a_{ij})$$

Furthermore,  $\phi_B(\overline{a_{ij}}) = \overline{\phi_B(a_{ij})}$  and  $\psi(\overline{a_{ij}}) = \overline{\psi(a_{ij})}$ , so it suffices to prove the lemma for  $\Sigma_n^{(p)}$  in the case where  $i < j$ .

does this need justification?

With these restrictions, we then break the statement up into the cases from Lemma 3.5, from which the first four cases as well as the last case can be checked easily. Consider the sixth case. Lemma 3.5 gives that

$$\psi\left(\phi_{\Sigma_n^{(p)}}(a_{ij})\right) = \sum_{Y \subseteq \{np-p+1, \dots, np\}} (-1)^{|Y|} \psi(a_{iy_1} a_{y_1 y_2} \cdots a_{y_k, j+p})$$

Let  $\alpha_i = np - p + r_i$ . Note that if  $y_1 < \alpha_i$  then  $\psi(a_{iy_1}) = 0$ , and if  $y_k > \alpha_j$  then  $\psi(a_{y_k j}) = 0$ , so the sum on the right hand side can be taken over  $Y \subseteq \{\alpha_i, \alpha_i + 1, \dots, \alpha_j\}$ . Then we manipulate the sum to get

do I need to explain what I'm doing here?

$$\begin{aligned} & \sum_{Y \subseteq \{\alpha_i, \dots, \alpha_j\}} (-1)^{|Y|} \psi(a_{iy_1} a_{y_1 y_2} \cdots a_{y_k, j+p}) \\ &= \psi(a_{i, j+p} - a_{i, \alpha_i} a_{\alpha_i, j+p}) \\ &+ \sum_{y=\alpha_i+1}^{\alpha_j} \sum_{Y \subseteq \{y+1, \dots, \alpha_j\}} (-1)^{|Y|+1} \psi(a_{iy} a_{y y_1} \cdots a_{y_k, j+p}) + (-1)^{|Y|} \psi(a_{i, \alpha_i} a_{\alpha_i, y} a_{y y_1} \cdots a_{y_k, j+p}) \\ &= \psi(a_{i, j+p} - a_{i, \alpha_i} a_{\alpha_i, j+p}) \\ &+ \sum_{y=\alpha_i+1}^{\alpha_j} \sum_{Y \subseteq \{y+1, \dots, \alpha_j\}} (-1)^{|Y|} \psi(a_{i, \alpha_i} a_{\alpha_i, y} - a_{iy}) \psi(a_{y y_1} \cdots a_{y_k, j+p}) \end{aligned}$$

Note that  $r_i = r_{\alpha_i}$  and since we're in the sixth case we have  $(n-1)p < j \leq np$ , so  $q_{\alpha_i} = q_y$ . Thus  $\psi(a_{i,\alpha_i}) = a_{q_i+1,q_{\alpha_i}+1} \otimes 1 = a_{q_i+1,q_y+1} \otimes 1$  and  $\psi(a_{\alpha_i,y}) = 1 \otimes a_{r_{\alpha_i},r_y} = 1 \otimes a_{r_i,r_y}$ , so we have

$$\psi(a_{i,\alpha_i}a_{\alpha_i,y} - a_{iy}) = (a_{q_i+1,q_y+1} \otimes 1)(1 \otimes a_{r_i,r_y}) - a_{q_i+1,q_y+1} \otimes a_{r_i,r_y} = 0$$

Thus the right hand side reduces to

$$\psi(a_{i,j+p} - a_{i\alpha_i}a_{\alpha_i,j+p})$$

**Remark 3.6.** The fact that  $\psi(a_{i,\alpha_i}a_{\alpha_i,y} - a_{iy}) = 0$  and  $\psi$  behaves similarly for the analogous terms in cases 5 and 7 is the key to this proof working, and  $\psi$  is defined the way it is mainly so that this will be true. As we hinted at earlier, the homomorphism  $\rho: \mathcal{A}_{pk} \rightarrow \mathcal{A}_k$  defined to send  $a_{ij}$  to  $a_{q_i+1,q_j+1}$  if  $r_i = r_j$  and to 0 otherwise would also send these terms to 0, so Proposition 3.1 would still be true with  $\rho$  used in the place of  $\psi$ . We will need  $\psi$  for the proof of the main result, however.

Note that, since we're in the sixth case,  $q_j + 1 = n$ . If  $r_i = r_j$ , then

$$\psi(a_{i,j+p} - a_{i\alpha_i}a_{\alpha_i,j+p}) = (a_{q_i+1,n+1} - a_{q_i+1,n}a_{n,n+1}) \otimes 1 = (\phi_{\sigma_n} \otimes \text{id})(\psi(a_{ij}))$$

If  $r_i < r_j$ , then

$$\begin{aligned} \psi(a_{i,j+p} - a_{i\alpha_i}a_{\alpha_i,j+p}) &= (a_{q_i+1,n+1} \otimes a_{r_i r_j} - a_{q_i+1,n}a_{n,n+1} \otimes a_{r_i r_j}) \\ &= (a_{q_i+1,n+1} - a_{q_i+1,n}a_{n,n+1}) \otimes a_{r_i r_j} \\ &= (\phi_{\sigma_n} \otimes \text{id})(\psi(a_{ij})) \end{aligned}$$

Finally, if  $r_i > r_j$ , then

$$\psi(a_{i,j+p} - a_{i\alpha_i}a_{\alpha_i,j+p}) = 0 = (\phi_{\sigma_n} \otimes \text{id})(\psi(a_{ij}))$$

The proof for the seventh case goes exactly as the proof for the sixth case except with all  $i$ 's replaced with  $i+p$ , all  $(j+p)$ 's replaced with  $j$ , all  $y_i$ 's replaced with  $y_{k+1-i}$ , and with  $\alpha_i$  and  $\alpha_j$  swapped. The proof for the fifth case goes exactly as the proof for the seventh.  $\square$

check this

check

*Proof of Lemma 3.4.* First we will prove the lemma for  $\Sigma_n^{(p)}$ . We can extend the definition of  $\psi$  to be an algebra morphism from the free module over  $\mathcal{A}_{pk}$  generated by the symbols  $\{a_{i*} | 1 \leq i \leq pk\}$  to the free module over  $\mathcal{A}_k \otimes \mathcal{A}_p$  generated by  $\{a_{i*} | 1 \leq i \leq k\}$  by defining  $\psi(a_{i*}) = a_{i*}$  and extending it to an algebra morphism. Then the statement of the lemma is equivalent to saying that for all  $1 \leq i \leq pk$ , the coefficient of  $a_{j*}$  in  $\psi(\phi_{\Sigma_n^{(p)}}(a_{i*}))$  is equal to 0 unless  $r_j = r_i$ , in which case it is equal to the coefficient of  $a_{q_j*}$  in  $\phi_{\sigma_n}(a_{q_i*})$ . If  $q_i + 1 \neq n$ , this fact can be easily checked. In the case that  $q_i + 1 = n$ , we have  $i = (n-1)p + r_i = \alpha_i$ , so

$$\psi(\phi_{\Sigma_n^{(p)}}(a_{i*})) = \psi(A(i+p, *, \{np-p+1, \dots, np\}))$$

which is equal to

$$\psi(a_{i+p,*} - a_{i+p,\alpha_i}a_{\alpha_i,*}) = a_{i+p,*} - (a_{n+1,n} \otimes 1)a_{\alpha_i,*}$$

by the same argument that was used in Lemma 3.2. The coefficients of the  $a_{j*}$  are equal to the coefficients of the  $a_{qj*}$  in  $\phi_{\sigma_n}(a_{q_i*})$ , so we have

$$\psi\left(\Phi_{\Sigma_n^{(p)}}^L\right) = \left((\Phi_{\sigma_n}^L)_{ij} \otimes 1\right) \otimes I_p.$$

Using this fact, the Chain Rule, and Lemma 3.2, we have

$$\begin{aligned} \left((I_{pk})_{ij} \otimes 1\right) &= \psi\left(\Phi_{\Sigma_n^{-(p)}\Sigma_n^{(p)}}^L\right) \\ &= \psi\left(\phi_{\Sigma_n^{-(p)}}\left(\Phi_{\Sigma_n^{(p)}}^L\right)\right) \psi\left(\Phi_{\Sigma_n^{-(p)}}^L\right) \\ &= \left(\phi_{\sigma_n^{-1}} \otimes \text{id}\right) \left(\left((\Phi_{\sigma_n})_{ij} \otimes 1\right) \otimes I_p\right) \psi\left(\Phi_{\Sigma_n^{-(p)}}^L\right) \end{aligned}$$

But note that the Chain Rule also gives that  $\left(\left(\Phi_{\sigma_n^{-1}}^L\right)_{ij} \otimes 1\right) \otimes I_p$  is the inverse of  $\left(\phi_{\sigma_n^{-1}} \otimes \text{id}\right) \left(\left((\Phi_{\sigma_n})_{ij} \otimes 1\right) \otimes I_p\right)$ , so

$$\psi\left(\Phi_{\Sigma_n^{-(p)}}^L\right) = \left(\left(\Phi_{\sigma_n^{-1}}^L\right)_{ij} \otimes 1\right) \otimes I_p$$

which completes the proof.  $\square$

*Proof of Lemma 3.5.*

check

We will prove a more general statement than the one presented in Lemma 3.5. Let  $\kappa_{m,l} = \tau_{m+l-1,p}\tau_{m+l-2,p} \cdots \tau_{m,p}$ , and let  $X_{m,l} = \{m, \dots, m+l-1\}$ . We will prove that if  $i < j$ , then

check

$$\phi_{\kappa_{m,l}}(a_{ij}) = \begin{cases} a_{i-p,j-p} & : m+p \leq i < j < m+l+p \\ a_{i-p,j} & : m+p \leq i < m+l+p \leq j \\ a_{i,j-p} & : i < m < m+p \leq j < m+l+p \\ a_{i+l,j+l} & : m \leq i < j < m+p \\ A'(i+l, j-p, X_{m,l}) & : m \leq i < m+p \leq j < m+l+p \\ A(i, j+l, X_{m,l}) & : i < m \leq j < m+p < m+l+p \\ A'(i+l, j, X_{m,l}) & : m \leq i < m+p < m+l+p \leq j \\ a_{ij} & : \text{otherwise} \end{cases}$$

Letting  $l = p$  and  $m = (n-1)p + 1$  then gives us Lemma ?? as a special case. The first four cases as well as the eighth can be easily checked. We will prove the remaining cases by induction on  $l$ . Consider the sixth case. The base case is covered by (7). For the inductive step, we have that

$$\begin{aligned}
\phi_{\kappa_m, l}(a_{ij}) &= \phi_{\tau_{m+l-1, p}}(\phi_{\kappa_m, l-1}(a_{ij})) \\
&= \sum_{Y \subseteq \{m, \dots, m+l-2\}} (-1)^{|Y|} \phi_{\tau_{m+l-1, p}}(a_{i, y_1} a_{y_1 y_2} \cdots a_{y_k, j+l-1}) \\
&= \sum_{Y \subseteq \{m, \dots, m+l-2\}} (-1)^{|Y|} a_{i, y_1} a_{y_1 y_2} \cdots a_{y_{k-1} y_k} (a_{y_k, j+l} - a_{y_k, m+l-1} a_{m+l-1, j+l}) \\
&= \sum_{Y \subseteq \{m, \dots, m+l-1\}} (-1)^{|Y|} a_{i, y_1} a_{y_1 y_2} \cdots a_{y_k, j+l} \\
&= A(i, j+l, X_{m, l})
\end{aligned}$$

is this clear/can it be shortened?

The proof of the seventh case goes exactly as the proof of the sixth, with all  $i$ 's replaced with  $i+l$ ,  $j$ 's replaced with  $j-l$ , and  $y_i$ 's replaced with  $y_{k-i+1}$ . The proof of the fifth case goes exactly as the proof of the seventh.  $\square$

#### REFERENCES

- [Cor13a] C. Cornwell. KCH representations, augmentations, and  $A$ -polynomials, 2013. arXiv: 1310.7526.
- [Cor13b] C. Cornwell. Knot contact homology and representations of knot groups. arXiv: 1303.4943, 2013.
- [EENS13] T. Ekholm, J. Etnyre, L. Ng, and M. Sullivan. Knot contact homology. *Geom. Topol.*, 17:975–1112, 2013.
- [Kir95] R. (Ed.) Kirby. Problems in low-dimensional topology. In *Proceedings of Georgia Topology Conference, Part 2*, pages 35–473. Press, 1995.
- [Ng05] L. Ng. Knot and braid invariants from contact homology I. *Geom. Topol.*, 9:247–297, 2005.
- [Ng08] L. Ng. Framed knot contact homology. *Duke Math. J.*, 141(2):365–406, 2008.
- [Ng12] L. Ng. A topological introduction to knot contact homology, 2012. arXiv: 1210.4803.