

# Optimization from Calculus

---

Chris Cornwell

September 11 and 16, 2025

Approximations from derivatives

Stationary points

Approximations from derivatives

Stationary points


# Linear approximations

$g : \mathbb{R} \rightarrow \mathbb{R}$ , a twice-differentiable function (at least). Write  $w$  for input to  $g$ , so  $g(w)$ .

**Linear approximation** to  $g(w)$ . At a point  $(v, g(v))$  on its graph, function whose graph is the tangent line:

$$h(w) = g(v) + g'(v)(w - v).$$

- Also called first order Taylor series approximation (or Taylor polynomial) of  $g$ .
- Approximates values of  $g$ , for inputs near  $v$ .



## Second order (quadratic) approximations

To approximate  $g$  better (and in a larger interval around  $v$ ), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

## Second order (quadratic) approximations

To approximate  $g$  better (and in a larger interval around  $v$ ), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

- Incorporates both first and second derivative.

## Second order (quadratic) approximations

To approximate  $g$  better (and in a larger interval around  $v$ ), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$


- Incorporates both first and second derivative.
- **Local minimum.** If  $g'(v) = 0$  and  $g''(v) > 0$ , then  $h(w) \geq g(v)$ .  
(near  $v$ , approximation is good, values of  $g(w) \geq g(v)$ ).

## Second order (quadratic) approximations

To approximate  $g$  better (and in a larger interval around  $v$ ), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

- Incorporates both first and second derivative.
- **Local minimum.** If  $g'(v) = 0$  and  $g''(v) > 0$ , then  $h(w) \geq g(v)$ . (near  $v$ , approximation is good, values of  $g(w) \geq g(v)$ ).
- If  $g'(v) = 0$  and  $g''(v) < 0$ , then opposite of the last item is true.





## First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if  $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$ , then

$$\nabla g = \left[ \frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

## First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if  $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$ , then

$$\nabla g = \left[ \frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

**Linear approximation:** is  $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T(\mathbf{w} - \mathbf{v})$ .

## First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if  $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$ , then

$$\nabla g = \left[ \frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

**Linear approximation:** is  $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T(\mathbf{w} - \mathbf{v})$ .

- Is a (affine) linear function; graph is translation of subspace of  $\mathbb{R}^{N+1}$  that is normal to  $[-\nabla g(\mathbf{v})^T, 1]^T$

## First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if  $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$ , then

$$\nabla g = \left[ \frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

**Linear approximation:** is  $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T(\mathbf{w} - \mathbf{v})$ .

- Is a (affine) linear function; graph is translation of subspace of  $\mathbb{R}^{N+1}$  that is normal to  $[-\nabla g(\mathbf{v})^T, 1]^T$
- The graph of  $h(\mathbf{w})$  is the tangent (hyper)plane to graph of  $g(\mathbf{w})$  at the point  $(\mathbf{v}, g(\mathbf{v}))$ .

## Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**,  $\nabla^2 g$ .

## Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**,  $\nabla^2 g$ .

- The matrix  $\nabla^2 g$  (evaluated at  $\mathbf{v}$ ) is the matrix of second order partial derivatives of  $g$ :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

## Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**,  $\nabla^2 g$ .

- The matrix  $\nabla^2 g$  (evaluated at  $\mathbf{v}$ ) is the matrix of second order partial derivatives of  $g$ :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

**Second order approximation:** is

$$h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v}) + \frac{1}{2} (\mathbf{w} - \mathbf{v})^T \nabla^2 g(\mathbf{v}) (\mathbf{w} - \mathbf{v}).$$

## Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**,  $\nabla^2 g$ .

- The matrix  $\nabla^2 g$  (evaluated at  $\mathbf{v}$ ) is the matrix of second order partial derivatives of  $g$ :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

**Second order approximation:** is

$$h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v}) + \frac{1}{2} (\mathbf{w} - \mathbf{v})^T \nabla^2 g(\mathbf{v}) (\mathbf{w} - \mathbf{v}).$$

- There is something like the second derivative test in this general case; is more involved to describe. In practice, techniques using first order approximations are the most commonly used.



Approximations from derivatives

Stationary points

## Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of  $g$  is a point  $\mathbf{v}$  where  $\nabla g(\mathbf{v})$  is the zero vector. (Called the “first order condition for optimality.”)

## Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of  $g$  is a point  $\mathbf{v}$  where  $\nabla g(\mathbf{v})$  is the zero vector. (Called the “first order condition for optimality.”)

- Means that  $\mathbf{v}$  is point where all partial derivatives of  $g$  are zero.

## Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of  $g$  is a point  $\mathbf{v}$  where  $\nabla g(\mathbf{v})$  is the zero vector. (Called the “first order condition for optimality.”)

- Means that  $\mathbf{v}$  is point where all partial derivatives of  $g$  are zero.
- A minimum of  $g$  can only occur at a stationary point. However, other things can happen at a stationary point too – maximum of  $g$  or a “saddle” point.