

Optimization from Calculus

Chris Cornwell

September 11 and 16, 2025

Approximations from derivatives

Stationary points

Approximations from derivatives

Stationary points


Linear approximations

$g : \mathbb{R} \rightarrow \mathbb{R}$, a twice-differentiable function (at least). Write w for input to g , so $g(w)$.

Linear approximation to $g(w)$. At a point $(v, g(v))$ on its graph, function whose graph is the tangent line:

$$h(w) = g(v) + g'(v)(w - v).$$

- Also called first order Taylor series approximation (or Taylor polynomial) of g .
- Approximates values of g , for inputs near v .



Second order (quadratic) approximations

To approximate g better (and in a larger interval around v), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

Second order (quadratic) approximations

To approximate g better (and in a larger interval around v), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

- Incorporates both first and second derivative.

Second order (quadratic) approximations

To approximate g better (and in a larger interval around v), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$


- Incorporates both first and second derivative.
- **Local minimum.** If $g'(v) = 0$ and $g''(v) > 0$, then $h(w) \geq g(v)$.
(near v , approximation is good, values of $g(w) \geq g(v)$).

Second order (quadratic) approximations

To approximate g better (and in a larger interval around v), can use the second order Taylor polynomial. This is the function:

$$h(w) = g(v) + g'(v)(w - v) + \frac{1}{2}g''(v)(w - v)^2.$$

- Incorporates both first and second derivative.
- **Local minimum.** If $g'(v) = 0$ and $g''(v) > 0$, then $h(w) \geq g(v)$. (near v , approximation is good, values of $g(w) \geq g(v)$).
- If $g'(v) = 0$ and $g''(v) < 0$, then opposite of the last item is true.



First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$, then

$$\nabla g = \left[\frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$, then

$$\nabla g = \left[\frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

Linear approximation: is $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v})$.

First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$, then

$$\nabla g = \left[\frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

Linear approximation: is $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v})$.

- Is a (affine) linear function; graph is translation of subspace of \mathbb{R}^{N+1} that is normal to $[-\nabla g(\mathbf{v})^T, 1]^T$

First order approximation in multiple variables

Extending to the setting of multiple variables.

- Use the gradient where, if $\mathbf{w} = [w_1 \ w_2 \ \dots \ w_N]^T$, then

$$\nabla g = \left[\frac{\partial g}{\partial w_1} \ \frac{\partial g}{\partial w_2} \ \dots \ \frac{\partial g}{\partial w_N} \right]^T.$$

Linear approximation: is $h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T(\mathbf{w} - \mathbf{v})$.

- Is a (affine) linear function; graph is translation of subspace of \mathbb{R}^{N+1} that is normal to $[-\nabla g(\mathbf{v})^T, 1]^T$
- The graph of $h(\mathbf{w})$ is the tangent (hyper)plane to graph of $g(\mathbf{w})$ at the point $(\mathbf{v}, g(\mathbf{v}))$.

Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**, $\nabla^2 g$.

Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**, $\nabla^2 g$.

- The matrix $\nabla^2 g$ (evaluated at \mathbf{v}) is the matrix of second order partial derivatives of g :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**, $\nabla^2 g$.

- The matrix $\nabla^2 g$ (evaluated at \mathbf{v}) is the matrix of second order partial derivatives of g :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

Second order approximation: is

$$h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v}) + \frac{1}{2} (\mathbf{w} - \mathbf{v})^T \nabla^2 g(\mathbf{v}) (\mathbf{w} - \mathbf{v}).$$

Second order approximation in multiple variables

For second order approximations, use a matrix called the **Hessian**, $\nabla^2 g$.

- The matrix $\nabla^2 g$ (evaluated at \mathbf{v}) is the matrix of second order partial derivatives of g :

$$\nabla^2 g = \begin{bmatrix} \frac{\partial^2 g}{\partial w_1 \partial w_1} & \frac{\partial^2 g}{\partial w_1 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_1 \partial w_N} \\ \frac{\partial^2 g}{\partial w_2 \partial w_1} & \frac{\partial^2 g}{\partial w_2 \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_2 \partial w_N} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 g}{\partial w_N \partial w_1} & \frac{\partial^2 g}{\partial w_N \partial w_2} & \cdots & \frac{\partial^2 g}{\partial w_N \partial w_N} \end{bmatrix}.$$

Second order approximation: is

$$h(\mathbf{w}) = g(\mathbf{v}) + \nabla g(\mathbf{v})^T (\mathbf{w} - \mathbf{v}) + \frac{1}{2} (\mathbf{w} - \mathbf{v})^T \nabla^2 g(\mathbf{v}) (\mathbf{w} - \mathbf{v}).$$

- There is something like the second derivative test in this general case; is more involved to describe. In practice, techniques using first order approximations are the most commonly used.

Approximations from derivatives

Stationary points

Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of g is a point \mathbf{v} where $\nabla g(\mathbf{v})$ is the zero vector. (Called the “first order condition for optimality.”)

Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of g is a point \mathbf{v} where $\nabla g(\mathbf{v})$ is the zero vector. (Called the “first order condition for optimality.”)

- Means that \mathbf{v} is point where all partial derivatives of g are zero.

Stationary points versus (local) minimal

A **stationary point** (or, **critical point**) of g is a point \mathbf{v} where $\nabla g(\mathbf{v})$ is the zero vector. (Called the “first order condition for optimality.”)

- Means that \mathbf{v} is point where all partial derivatives of g are zero.
- A minimum of g can only occur at a stationary point. However, other things can happen at a stationary point too – maximum of g or a “saddle” point.

The Chain rule

The **chain rule** is very useful for understanding derivatives and gradients. The more general version of it (“suped up” from Calculus I):

¹“ Df ” and “ Dg ” mean, take the appropriate version of the derivative for the function. If the function is from \mathbb{R} to \mathbb{R} , this is the f' from Calc I; if it is multi-variable (from \mathbb{R}^N to \mathbb{R} , for some $N > 1$), then Df means ∇f . If f has vector output, there is a matrix of partial derivatives for Df .

The Chain rule

The **chain rule** is very useful for understanding derivatives and gradients. The more general version of it (“suped up” from Calculus I):

Chain rule.

If the composition $f(g(\mathbf{w}))$ is defined and each of f and g are differentiable, then the derivative of the composition is

$$Df(g(\mathbf{w})) \cdot Dg(\mathbf{w}).^1$$

¹“ Df ” and “ Dg ” mean, take the appropriate version of the derivative for the function. If the function is from \mathbb{R} to \mathbb{R} , this is the f' from Calc I; if it is multi-variable (from \mathbb{R}^N to \mathbb{R} , for some $N > 1$), then Df means ∇f . If f has vector output, there is a matrix of partial derivatives for Df .

The Chain rule

The **chain rule** is very useful for understanding derivatives and gradients. The more general version of it (“suped up” from Calculus I):

Chain rule.

If the composition $f(g(\mathbf{w}))$ is defined and each of f and g are differentiable, then the derivative of the composition is $Df(g(\mathbf{w})) \cdot Dg(\mathbf{w})$.¹

Example: given a fixed vector \mathbf{a} , set $h(\mathbf{w}) = \frac{1}{1+e^{\mathbf{a}^T \mathbf{w}}}$. This is a composition of $f(x) = \frac{1}{1+e^x}$ and $g(\mathbf{w}) = \mathbf{a}^T \mathbf{w}$. Thus,

$$\nabla h(\mathbf{w}) = f'(g(\mathbf{w})) \cdot \nabla g(\mathbf{w}) = \left(\frac{-e^{\mathbf{a}^T \mathbf{w}}}{1 + e^{\mathbf{a}^T \mathbf{w}}} \right) \mathbf{a}.$$

¹“ Df ” and “ Dg ” mean, take the appropriate version of the derivative for the function. If the function is from \mathbb{R} to \mathbb{R} , this is the f' from Calc I; if it is multi-variable (from \mathbb{R}^N to \mathbb{R} , for some $N > 1$), then Df means ∇f . If f has vector output, there is a matrix of partial derivatives for Df .

Example of stationary points

Here we look at the stationary points of the function

$$f(w_1, w_2) = w_1^3 + 3w_1w_2 + w_2^3.$$

- $\nabla f(w_1, w_2) = [3w_1^2 + 3w_2, \quad 3w_1 + 3w_2^2]^T$

Example of stationary points

Here we look at the stationary points of the function

$$f(w_1, w_2) = w_1^3 + 3w_1w_2 + w_2^3.$$

- $\nabla f(w_1, w_2) = [3w_1^2 + 3w_2, \quad 3w_1 + 3w_2^2]^T$

- If a stationary pt.,

$$3w_1^2 + 3w_2 = 0 \text{ and}$$

$$3w_1 + 3w_2^2 = 0; \text{ so, } w_2 = -w_1^2;$$

using this and simplifying,

$$w_1 + w_1^4 = 0.$$

Example of stationary points

Here we look at the stationary points of the function


$$f(w_1, w_2) = w_1^3 + 3w_1w_2 + w_2^3.$$

- $\nabla f(w_1, w_2) = [3w_1^2 + 3w_2, \quad 3w_1 + 3w_2^2]^T$

- If a stationary pt.,
 $3w_1^2 + 3w_2 = 0$ and
 $3w_1 + 3w_2^2 = 0$; so, $w_2 = -w_1^2$;
using this and simplifying,
 $w_1 + w_1^4 = 0$.

- We get: either $w_1 = 0$ (with
 $w_2 = 0$ also), or $w_1 = -1$ (with
 $w_2 = -1$ also).

See Figure on right.



Vertical lines placed over stationary points $(0, 0)$ and $(-1, -1)$

Convex functions

Function $g : \mathbb{R} \rightarrow \mathbb{R}$ is called **convex** if $g''(v) \geq 0$ for all v .²


²A different definition (which is equivalent): g is **convex at** v if the linear approximation at v has a graph that is below the graph of g , near v ; it is then **convex** if it is convex at v for all v in \mathbb{R} . This extends to multiple variables.

Convex functions

Function $g : \mathbb{R} \rightarrow \mathbb{R}$ is called **convex** if $g''(v) \geq 0$ for all v .²

Example.

The function $g(w) = e^w - w$ is a convex function, since $g''(w) = e^w$, and e^v is bigger than zero for all v .



²A different definition (which is equivalent): g is **convex at v** if the linear approximation at v has a graph that is below the graph of g , near v ; it is then **convex** if it is convex at v for all v in \mathbb{R} . This extends to multiple variables.