

Linear Regression, Method 1

Chris Cornwell

Feb 11, 2025

Outline

Overview of linear regression task

The procedure

Implementing the procedure

Examples

Outline

Overview of linear regression task

The procedure

Implementing the procedure


Examples

The goal

- ▶ Setting: have points in the plane, say n of them. Say the points are $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$.
- ▶ **Goal:** *Model* them as being “noisy” points from a line, finding “best fit” line (the closest linear model). This line is also called the **least squares regression** (LSR) line.

The goal

- Setting: have points in the plane, say n of them. Say the points are $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$.
- **Goal:** *Model* them as being “noisy” points from a line, finding “best fit” line (the closest linear model). This line is also called the **least squares regression** (LSR) line.
- **Running example:** A data set, '[Example1.csv](#)', with 50 points, is [available here](#); these points are displayed in the plot.



Outline

Overview of linear regression task

The procedure

Implementing the procedure

Examples

Finding the LSR line




Figure: Our running example

How do we find the LSR line?

Finding the LSR line




Figure: Our running example

How do we find the LSR line?

Can get the slope m , intercept b simply from using the `polyfit` function in NumPy. If x , y are the arrays with x - and y -coordinates:

```
| np.polyfit(x,y,1)
```


Finding the LSR line

But, how are the slope, intercept found?

¹Will use \mathbf{p} for this vector, for the rest of these slides.

Finding the LSR line

But, how are the slope, intercept found?

- If a slope m and intercept b existed so that $(x_1, y_1), \dots, (x_{50}, y_{50})$ were points on $y = mx + b$, then

$$y_i = mx_i + b$$

would hold for all $1 \leq i \leq 50$.

¹Will use \mathbf{p} for this vector, for the rest of these slides.

Finding the LSR line

But, how are the slope, intercept found?

- If a slope m and intercept b existed so that $(x_1, y_1), \dots, (x_{50}, y_{50})$ were points on $y = mx + b$, then

$$y_i = mx_i + b$$

would hold for all $1 \leq i \leq 50$.

1. Write those 50 equations as a matrix equation. Setting:

$$A = \begin{bmatrix} x_1 & 1 \\ x_2 & 1 \\ \vdots & \vdots \\ x_{50} & 1 \end{bmatrix}; \quad \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{50} \end{bmatrix},$$

and writing¹ $\mathbf{p} = \begin{bmatrix} m \\ b \end{bmatrix}$, the matrix equation is $A\mathbf{p} = \mathbf{y}$.

¹Will use \mathbf{p} for this vector, for the rest of these slides.

Finding the LSR line


Now the equation $A\mathbf{p} = \mathbf{y}$ does not have a solution (those points are *not* on a line).

Finding the LSR line

Now the equation $A\mathbf{p} = \mathbf{y}$ does not have a solution (those points are *not* on a line).

Next idea: (*thinking of noise being in y-coordinate direction*)

- Get vector $\hat{\mathbf{y}}$ that is *as close to \mathbf{y} as possible*, so that $A\mathbf{p} = \hat{\mathbf{y}}$ has a solution.




Finding the LSR line

Now the equation $A\mathbf{p} = \mathbf{y}$ does not have a solution (those points are *not* on a line).

Next idea: (*thinking of noise being in y-coordinate direction*)

- ▶ Get vector $\hat{\mathbf{y}}$ that is *as close to \mathbf{y} as possible*, so that $A\mathbf{p} = \hat{\mathbf{y}}$ has a solution.
- ▶ For each i , we either increase or decrease y_i by a (hopefully small) amount, $y_i \rightsquigarrow \hat{y}_i$. We make $|\mathbf{y} - \hat{\mathbf{y}}|$ as small as possible.



Finding the LSR line

Now the equation $A\mathbf{p} = \mathbf{y}$ does not have a solution (those points are *not* on a line).

Next idea: (thinking of noise being in y -coordinate direction)


- Get vector $\hat{\mathbf{y}}$ that is as close to \mathbf{y} as possible, so that $A\mathbf{p} = \hat{\mathbf{y}}$ has a solution.
- For each i , we either increase or decrease y_i by a (hopefully small) amount, $y_i \rightsquigarrow \hat{y}_i$. We make $|\mathbf{y} - \hat{\mathbf{y}}|$ as small as possible.

2. Done by solving

$$A^T A \mathbf{p} = A^T \mathbf{y} \text{ (normal equations).}$$

If $\mathbf{p} = \begin{bmatrix} \hat{m} \\ \hat{b} \end{bmatrix}$ is the solution,

then $\hat{\mathbf{y}}$ is given by $A \begin{bmatrix} \hat{m} \\ \hat{b} \end{bmatrix}$.



Normal equation

Why does solving $A^T A \mathbf{p} = A^T \mathbf{y}$ give the right thing?

Normal equation

Why does solving $A^T A \mathbf{p} = A^T \mathbf{y}$ give the right thing?

Related to orthogonal vectors in \mathbb{R}^n (in the example, \mathbb{R}^{50}).

- If $A \mathbf{p} = \hat{\mathbf{y}}$ has a solution, then $\hat{\mathbf{y}} \in \mathbb{R}^{50}$ is in column space of A .

Normal equation

Why does solving $A^T A \mathbf{p} = A^T \mathbf{y}$ give the right thing?

Related to orthogonal vectors in \mathbb{R}^n (in the example, \mathbb{R}^{50}).

- ▶ If $A \mathbf{p} = \hat{\mathbf{y}}$ has a solution, then $\hat{\mathbf{y}} \in \mathbb{R}^{50}$ is in column space of A .
- ▶ $\mathbf{y} = \mathbf{z}_1 + \mathbf{z}_2$, where \mathbf{z}_1 in null space of A^T and \mathbf{z}_2 in column space of A .

(Note: Null space of A^T orthogonal to column space of A .)

Normal equation

Why does solving $A^T A \mathbf{p} = A^T \mathbf{y}$ give the right thing?

Related to orthogonal vectors in \mathbb{R}^n (in the example, \mathbb{R}^{50}).

- ▶ If $A \mathbf{p} = \hat{\mathbf{y}}$ has a solution, then $\hat{\mathbf{y}} \in \mathbb{R}^{50}$ is in column space of A .
- ▶ $\mathbf{y} = \mathbf{z}_1 + \mathbf{z}_2$, where \mathbf{z}_1 in null space of A^T and \mathbf{z}_2 in column space of A .
(Note: Null space of A^T orthogonal to column space of A .)
- ▶ As \mathbf{z}_2 in column space, $\exists \hat{\mathbf{p}}$ so that $A \hat{\mathbf{p}} = \mathbf{z}_2$.

Normal equation

Why does solving $A^T A \mathbf{p} = A^T \mathbf{y}$ give the right thing?

Related to orthogonal vectors in \mathbb{R}^n (in the example, \mathbb{R}^{50}).

- ▶ If $A\mathbf{p} = \hat{\mathbf{y}}$ has a solution, then $\hat{\mathbf{y}} \in \mathbb{R}^{50}$ is in column space of A .
- ▶ $\mathbf{y} = \mathbf{z}_1 + \mathbf{z}_2$, where \mathbf{z}_1 in null space of A^T and \mathbf{z}_2 in column space of A .

(Note: Null space of A^T orthogonal to column space of A .)

- ▶ As \mathbf{z}_2 in column space, $\exists \hat{\mathbf{p}}$ so that $A\hat{\mathbf{p}} = \mathbf{z}_2$. But then,

$$A^T(A\hat{\mathbf{p}}) = A^T\mathbf{z}_2 = A^T(\mathbf{y} - \mathbf{z}_1) = A^T\mathbf{y}.$$

And \mathbf{z}_2 is closest, since subtracted \mathbf{z}_1 from \mathbf{y} , orthogonal to column space:

$$\mathbf{z}_2 = \hat{\mathbf{y}}.$$

Finding the LSR line

Normal equations:

$$(A^T A)\mathbf{p} = A^T \mathbf{y}.$$

Finding the LSR line

Normal equations:

$$(A^T A)\mathbf{p} = A^T \mathbf{y}.$$

Note:

- ▶ $A^T A$ is 2×2 matrix, $A^T \mathbf{y} \in \mathbb{R}^2$, and $A^T A$ is invertible as long as there exists $x_i \neq x_j$.

Finding the LSR line

Normal equations:

$$(A^T A)\mathbf{p} = A^T \mathbf{y}.$$

Note:

- ▶ $A^T A$ is 2×2 matrix, $A^T \mathbf{y} \in \mathbb{R}^2$, and $A^T A$ is invertible as long as there exists $x_i \neq x_j$.
- 3. Pretty straightforward to find solution to $(A^T A)\mathbf{p} = A^T \mathbf{y}$.

Finding the LSR line

Normal equations:

$$(A^T A)\mathbf{p} = A^T \mathbf{y}.$$

Note:

- ▶ $A^T A$ is 2×2 matrix, $A^T \mathbf{y} \in \mathbb{R}^2$, and $A^T A$ is invertible as long as there exists $x_i \neq x_j$.
- 3. Pretty straightforward to find solution to $(A^T A)\mathbf{p} = A^T \mathbf{y}$.

So, three steps:

1. Write the n equations in matrix form. (get matrix A , vector \mathbf{y})
2. Get matrix $A^T A$ and vector $A^T \mathbf{y}$ for normal equations: $A^T A \mathbf{p} = A^T \mathbf{y}$.
3. Use a method to solve normal equations for \mathbf{p} .

Outline

Overview of linear regression task

The procedure

Implementing the procedure

Examples

Solving normal equation, in pseudocode

Procedure to carry out the steps:

1. Write the n equations in matrix form. (get matrix A , vector \mathbf{y})
2. Get matrix $A^T A$ and vector $A^T \mathbf{y}$ for normal equations: $A^T A \mathbf{p} = A^T \mathbf{y}$.
3. Use a method to solve normal equations for \mathbf{p} .

Solving normal equation, in pseudocode

Procedure to carry out the steps:

1. Write the n equations in matrix form. (get matrix A , vector y)
2. Get matrix $A^T A$ and vector $A^T y$ for normal equations: $A^T A p = A^T y$.
3. Use a method to solve normal equations for p .

Given $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$, as a NumPy array (call it D , with shape $(n, 2)$):

```
A ← [x coordinates, all 1s] # 2-column matrix
y ← y coordinates
# next, get 2x2 matrix and 2-vector
Compute A.T times A; compute A.T times y
Solve normal eq'ns (numpy solve, or use inverse)
return solution
```

Outline

Overview of linear regression task

The procedure

Implementing the procedure

Examples

Result on running example

For the data (linked to above) with 50 points, the LSR line comes out close to


$$y = 1.520275x - 0.33458.$$

Result on running example

For the data (linked to above) with 50 points, the LSR line comes out close to

$$y = 1.520275x - 0.33458.$$

A plot of the line (in red), alongside the points, looks as follows.




Another example, Advertising data

In the [DataSets folder](#), the '[Advertising.csv](#)' file contains data on amounts spent (in thousands of dollars) on TV, Radio, and Newspaper advertising in 200 different markets, as well as the amounts sold in each market (in thousands of units).

Another example, Advertising data


In the [DataSets folder](#), the '[Advertising.csv](#)' file contains data on amounts spent (in thousands of dollars) on TV, Radio, and Newspaper advertising in 200 different markets, as well as the amounts sold in each market (in thousands of units).

We will look more at this data later. For now, plotted here are the columns ('TV', 'Sales').



Another example, Advertising data


If you switch the role of x- and y-coordinates, you can still do linear regression; i.e., for purpose of a thought experiment, predict the TV data as the *response*, instead of the Sales.



Another example, Advertising data

If you switch the role of x- and y-coordinates, you can still do linear regression; i.e., for purpose of a thought experiment, predict the TV data as the *response*, instead of the Sales.

The LSR line for the data is then **not** the same line, if you switch the roles of TV and Sales in the algorithm to get $\hat{\mathbf{p}}$.



Another example, Advertising data

If you switch the role of x- and y-coordinates, you can still do linear regression; i.e., for purpose of a thought experiment, predict the TV data as the *response*, instead of the Sales.

The LSR line for the data is then **not** the same line, if you switch the roles of TV and Sales in the algorithm to get $\hat{\mathbf{p}}$.

