

A spatio-temporal analysis of the environmental correlates of COVID-19 incidence in Spain

Antonio Paez^{*,a}, Fernando A. Lopez^b, Tatiane Menezes^c, Renata Cavalcanti^d,
Maira Galdino da Rocha Pitta^d

^a*School of Geography and Earth Sciences, McMaster University, 1281 Main St W, Hamilton, ON, L8S 4K1, Canada*

^b*Departamento de Metodos Cuantitativos, Ciencias Juridicas, y Lenguas Modernas, Universidad Politecnica de Cartagena, Calle Real Numero 3, 30201, Cartagena, Murcia, Spain*

^c*Departamento de Economia, Universidade Federal de Pernambuco, Av dos Economistas, s/n - Cidade Universitária, Recife - PE, 50670-901, Brasil*

^d*Núcleo de Pesquisa em Inovação Terapêutica NUPIT / UFPE, Av. Prof. Moraes Rego, 1235 - Cidade Universitária, Recife, PE, CEP 50670-901, Brazil*

Abstract

The novel SARS-CoV2 has disrupted health systems and the economy and public health interventions to slow its spread have been costly. How and when to ease restrictions to movement hinges in part on whether SARS-CoV2 will display seasonality due to variations in temperature, humidity, and hours of sunshine. Here, we address this question by means of a spatio-temporal analysis in Spain of the incidence of COVID-19, the disease caused by the virus. Use of spatial Seemingly Unrelated Regressions (SUR) allows us to model the incidence of reported cases of the disease per 100,000 population as an interregional contagion process, in addition to a function of temperature, humidity, and sunshine. In the analysis we also control for GDP per capita, percentage of older adults in the population, population density, and presence of mass transit systems. The results support the hypothesis that incidence of the disease is lower at higher temperatures and higher levels of humidity. Sunshine, in contrast, displays a positive association with incidence of the disease. Our control variables also yield interesting insights. Higher incidence is associated with higher GDP per capita and presence of mass transit systems in the province; in contrast, population density and percentage of older adults display negative associations with incidence of COVID-19.

*Corresponding Author

Email addresses: paezha@mcmaster.ca (Antonio Paez), fernando.lopez@upct.es (Fernando A. Lopez), tatiane.menezes@ufpe.br (Tatiane Menezes), renata.vcsantos@gmail.com (Renata Cavalcanti), mgrpitta@ufpe.br (Maira Galdino da Rocha Pitta)