

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow  
MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem

# Max-flow and min-cut

# Flow Network

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph

Augmenting  
path

MaxFlow  
MinCut Thm


Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

A network  $\mathcal{N} = (V, E, c, s, t)$  is  
formed by

- a digraph  $G = (V, E)$ ,
- a source vertex  $s \in V$
- a sink vertex  $t \in V$ ,
- and edge capacities  $c : E \rightarrow \mathbb{R}^+$



# A flow in a network

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\}, \sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



# A flow in a network

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\}, \sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



with value 3.

# A flow in a network

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\}, \sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



saturated

# The Maximum flow problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t, )$

QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem

# The Maximum flow problem

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t)$


QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .



# The Maximum flow problem

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t)$   
QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .



The value of the flow is  $7 = 4 + 1 + 2 = 5 + 2$ .


As  $t$  cannot receive more flow, this flow is a **maximum flow**.

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  $(s, t)$ -cut is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The capacity of a cut  $(S, T)$  is the sum of weights leaving  $S$ , i.e.,

$$c(S, T) = \sum_{u \in S} \sum_{v \in T} c(u, v)$$




$$\begin{aligned} S &= \{s, c, d\} \\ T &= \{a, b, e, t\} \\ c(S, T) &= 19 \\ (4+5)+5+(3+2) & \end{aligned}$$

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  **$(s, t)$ -cut** is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned}S &= \{s, c, d\} \\T &= \{a, b, e, t\} \\c(S, T) &= 19 \\f(S, T) &= 10 - 3 = 7\end{aligned}$$


- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  **$(s, t)$ -cut** is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned}S &= \{s, c, d\} \\T &= \{a, b, e, t\} \\c(S, T) &= 19 \\f(S, T) &= 10 - 3 = 7\end{aligned}$$


Due to the capacity constrain:  $f(S, T) \leq c(S, T)$

# Another $(s, t)$ -cut

Given  $\mathcal{N} = (V, E, c, s, t)$  a  **$(s, t)$ -cut** is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned} S &= \{s, a, c\} \\ T &= \{b, d, e, t\} \\ c(S, T) &= 19 \\ f(S, T) &= 8 - 1 = 7 \end{aligned}$$

# Changing weights effect on min cuts

Given a network  $\mathcal{N} = (V, E, s, t, c)$  assume that  $(S, T)$  is a min  $(s, t)$ -cut.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem

# Changing weights effect on min cuts

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem

Given a network  $\mathcal{N} = (V, E, s, t, c)$  assume that  $(S, T)$  is a min  $(s, t)$ -cut.

If we change the input by adding  $c > 0$  to the capacity of **every edge**, then it may happen that  $(S, T)$  is not longer a min  $(s, t)$ -cut.



# Changing weights effect on Min-Cut and Max-Flow

Given a network  $\mathcal{N} = (V, E, s, t, c)$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Changing weights effect on Min-Cut and Max-Flow

Given a network  $\mathcal{N} = (V, E, s, t, c)$ .

If we change the network by multiplying by  $c >$  the capacity of every edge, the capacity of any  $(s, t)$ -cut in the new network is  $c$  times its capacity in the original network.



# Notation

Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$

For  $v \in V$ ,  $U \subseteq V$  and  $v \notin U$ .

- $f(v, U)$  flow  $v \rightarrow U$  i.e.  $f(v, U) = \sum_{u \in U} f(v, u)$ ,
- $f(U, v)$  flow  $U \rightarrow v$  i.e.  $f(U, v) = \sum_{u \in U} f(u, v)$ ,

For a  $(s, t)$ -cut  $(S, T)$  and  $v \in S$

- $S' = S \setminus \{v\}$  and  $T' = T \cup \{v\}$
- $f_{-v}(S, T) = \sum_{u \in S'} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S'} f(v, u)$   
i.e, the contribution to  $f(S, T)$  from edges not incident with  $v$ .

# Flow conservation on $(s, t)$ -cuts

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem

Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$ . For any  $(s, t)$ -cut  $(S, T)$ ,  $f(S, T) = |f|$ .

## Proof (Induction on $|S|$ )

- If  $S = \{s\}$  then, by definition,  $f(S, T) = |f|$ .

# Flow conservation on $(s, t)$ -cuts

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem


Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$ . For any  $(s, t)$ -cut  $(S, T)$ ,  $f(S, T) = |f|$ .

## Proof (Induction on $|S|$ )

- If  $S = \{s\}$  then, by definition,  $f(S, T) = |f|$ .
- Assume it is true for  $S' = S - \{v\}$  and  $T' = T \cup \{v\}$ , i.e.  $f(S', T') = |f|$ .

# Flow conservation on $(s, t)$ -cuts

## Proof (cont.) (Induction on $|S|$ )




- IH:  $f(S', T') = |f|$ .
- Then,  $f(S, T) = f_{-v}(S, T) + f(v, T) - f(T, v)$ .
- But,  $f(S', T') = f_{-v}(S, T) + f(S', v) - f(v, S')$  as  $v \in T'$
- By flow conservation,  
$$f(S', v) + f(T, v) = f(v, S') + f(v, T)$$
- So,  $f(S', v) - f(v, S') = f(v, T) - f(T, v)$
- Therefore,  $f(S', T') = f(S, T) = |f|$

# Residual graph

Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a **flow**  $f$ .  
The **residual graph**,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:


- if  $c(u, v) - f(u, v) > 0$ , then  $(u, v) \in E_f$  and  $c_f(u, v) = c(u, v) - f(u, v) > 0$  (**forward edges**)



# Residual graph

Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a **flow**  $f$  on it, the **residual graph**,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:


- if  $f(u, v) > 0$ , then  $(v, u) \in E_f$  and  $c_f(v, u) = f(u, v)$  (**backward edges**).



# Residual graph


Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a **flow  $f$**  on it, the **residual graph**,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:

- if  $c(u, v) - f(u, v) > 0$ , then  $(u, v) \in E_f$  and  $c_f(u, v) = c(u, v) - f(u, v) > 0$  (**forward edges**)
- if  $f(u, v) > 0$ , then  $(v, u) \in E_f$  and  $c_f(v, u) = f(u, v)$  (**backward edges**).




# Residual graph

Max Flow and Min Cut  
Properties of flows and cuts  
Residual graph  
Augmenting path  
MaxFlow MinCut Thm  
Ford Fulkerson alg  
Maximum matching in Bip graphs  
Disjoint paths problem



- Notice that, if  $c(u, v) = f(u, v)$ , then there is only a backward edge.
- $c_f$  are called the **residual capacity**.


# Residual graph



- **forward edges:** There remains capacity to push more flow through this edge.
- **backward edges:** there are units of flow that can be redirected through other links.

# Augmenting paths


Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple** path  $P$  in  $G_f$  from  $s$  to  $t$

# Augmenting paths


Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple** path  $P$  in  $G_f$  from  $s$  to  $t$   $P$  might have forward and backward edges.

# Augmenting paths

Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple** path  $P$  in  $G_f$  from  $s$  to  $t$   $P$  might have forward and backward edges.
- For an augmenting path  $P$  in  $G_f$ , the **bottleneck**,  $b(P)$ , is the minimum (residual) capacity of the edges in  $P$ . In the example, for  $P = (s, a, t)$ ,  $b(P) = 1$ .

# Augmenting paths: increasing the flow

**Augment( $P, f$ )**

$b = \text{bottleneck } (P)$

**for each**  $(u, v) \in P$  **do**


**if**  $(u, v)$  is a forward edge **then**

    Increase  $f(u, v)$  by  $b$

**else**

    Decrease  $f(v, u)$  by  $b$

**return**  $f$



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths: increasing the flow

**Augment( $P, f$ )**

$b = \text{bottleneck } (P)$

**for** each  $(u, v) \in P$  **do**



**if**  $(u, v)$  is a forward edge **then**

        Increase  $f(u, v)$  by  $b$

**else**

        Decrease  $f(v, u)$  by  $b$

**return**  $f$



$G_f, P = (s, a, t), b(P) = 1$

# Augmenting paths: increasing the flow

**Augment( $P, f$ )**

$b = \text{bottleneck } (P)$

**for** each  $(u, v) \in P$  **do**




**if**  $(u, v)$  is a forward edge **then**

        Increase  $f(u, v)$  by  $b$

**else**

        Decrease  $f(v, u)$  by  $b$

**return**  $f$



# Augmenting paths: increasing the flow

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .

**Proof:** We have to prove the two flow properties.

### ■ Capacity law

- Forward edges  $(u, v) \in P$ , we increase  $f(u, v)$  by  $b$ , as  $b \leq c(u, v) - f(u, v)$  then  $f'(u, v) = f(u, v) + b \leq c(u, v)$ .
- Backward edges  $(u, v) \in P$  we decrease  $f(v, u)$  by  $b$ , as  $b \leq f(v, u)$ ,  $f'(v, u) = f(v, u) - b \geq 0$ .

# Augmenting paths: increasing the flow

## Lemma

Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .

**Proof:** We have to prove the two flow properties.

- **Conservation law**,  $\forall v \in P \setminus \{s, t\}$  let  $u$  be the predecessor of  $v$  in  $P$  and let  $w$  be its successor.
  - As the path is simple only the alterations due to  $(u, v)$  and  $(v, w)$  can change the flow that goes through  $v$ . We have three cases:
    - $(u, v)$  and  $(v, w)$  are backward edges, the flow in  $(v, u)$  and  $(w, v)$  is decremented by  $b$ . As one is incoming and the other outgoing the total balance is 0.
    - $(u, v)$  and  $(v, w)$  are forward edges, the flow in  $(u, v)$  and  $(v, w)$  is incremented by  $b$ . As one is incoming and the other outgoing the total balance is 0.

# Augmenting paths: increasing the flow

## Lemma

Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .

**Proof:** We have to prove the two flow properties.

- **Conservation law**,  $\forall v \in P \setminus \{s, t\}$  let  $u$  be the predecessor of  $v$  in  $P$  and let  $w$  be its successor.
  - As the path is simple only the alterations due to  $(u, v)$  and  $(v, w)$  can change the flow that goes through  $v$ . We have three cases:
    - $(u, v)$  is forward and  $(v, w)$  is backward, the flow in  $(u, v)$  is incremented by  $b$  and the flow in  $(w, v)$  is decremented by  $b$ . As both are incoming, the total balance is 0.
    - $(u, v)$  is backward and  $(v, w)$  is forward, the flow in  $(v, w)$  is incremented by  $b$  and the flow in  $(v, u)$  is decremented by  $b$ . As both are outgoing, the total balance is 0.

# Augmenting paths: incrementing the flow

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

Consider  $f' = \text{Augment}(P, f)$ , then  $|f'| > |f|$ .

Proof: Let  $P$  be the augmenting path in  $G_f$ . The first edge  $e \in P$  leaves  $s$ , and as  $G$  has no incoming edges to  $s$ ,  $e$  is a forward edge. Moreover  $P$  is simple  $\Rightarrow$  never returns to  $s$ . Therefore, the value of the flow increases in edge  $e$  by  $b$  units.

□

# Max-Flow Min-Cut theorem

Ford and Fulkerson (1954); Peter Elias, Amiel Feinstein and Claude Shannon (1956) (in framework of information-theory).

## Theorem

*For any  $\mathcal{N}(G, s, t, c)$ , the maximum of the flow value is equal to the minimum of the  $(S, T)$ -cut capacities.*

$$\max_f \{|f|\} = \min_{(S,T)} \{c(S, T)\}.$$

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V | \exists s \sim v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V | \exists s \sim v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $u \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(v, u) = c(v, u)$ ,

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V | \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $u \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(v, u) = c(v, u)$ ,
- Then,  $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$

# Max-Flow Min-Cut theorem: Proof


## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f\{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V | \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $u \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(v, u) = c(v, u)$ ,
- Then,  $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$
- $(S_s, V - \{S_s\})$  is a minimum capacity  $(s, t)$ -cut in  $G$ .





# Ford-Fulkerson algorithm

L.R. Ford, D.R. Fulkerson:  
*Maximal flow through a network.* Canadian J. of Math.  
1956.




Max Flow and  
Min Cut  
Properties of  
flows and cuts  
Residual graph  
Augmenting  
path  
MaxFlow  
MinCut Thm  
Ford  
Fulkerson alg  
Maximum  
matching in  
Bip graphs  
Disjoint paths  
problem

**Ford-Fulkerson**( $G, s, t, c$ )  
for all  $(u, v) \in E$  let  $f(u, v) = 0$   
 $G_f = G$   
**while** there is an  $s - t$  path  $P$  in  $G_f$  **do**  
     $f = \text{Augment}(P, G_f)$   
    Compute  $G_f$   
**return**  $f$





# Ford-Fulkerson algorithm

L.R. Ford, D.R. Fulkerson:  
*Maximal flow through a network.* Canadian J. of Math.  
1956.




Max Flow and  
Min Cut  
Properties of  
flows and cuts  
Residual graph  
Augmenting  
path  
MaxFlow  
MinCut Thm  
Ford  
Fulkerson alg  
Maximum  
matching in  
Bip graphs  
Disjoint paths  
problem

**Ford-Fulkerson**( $G, s, t, c$ )  
for all  $(u, v) \in E$  let  $f(u, v) = 0$   
 $G_f = G$   
**while** there is an  $s - t$  path  $P$  in  $G_f$  **do**  
     $f = \text{Augment}(P, G_f)$   
    Compute  $G_f$   
**return**  $f$




# FF example



- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem




# FF example



- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem



# FF example

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem




# FF example

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem






# FF example



- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem


# FF example

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem




# FF example

Max Flow and  
Min Cut  
Properties of  
flows and cuts  
Residual graph  
Augmenting  
path  
MaxFlow  
MinCut Thm  
Ford  
Fulkerson alg  
Maximum  
matching in  
Bip graphs  
Disjoint paths  
problem



Flow with max value



$\{s\}, \{a, b, t\}$  is a min  $(s, t)$ -cut

# Correctness of Ford-Fulkerson

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

Consequence of the Max-flow min-cut theorem.

## Theorem

*The flow returned by Ford-Fulkerson is the max-flow.*

# Networks with integer capacities

## Lemma (**Integrality invariant**)

Let  $\mathcal{N} = (V, E, c, s, t)$  where  $c : E \rightarrow \mathbb{Z}^+$ . At every iteration of the Ford-Fulkerson algorithm, the flow values  $f(e)$  are integers.

## Proof: (induction)

- The statement is true for the initial flow (all zeroes).
- Inductive Hypothesis: The statement is true after  $j$  iterations.
- At iteration  $j + 1$ : As all residual capacities in  $G_f$  are integers, then bottleneck  $(P, f) \in \mathbb{Z}$ , for the augmenting path found in iteration  $j + 1$ .
- Thus the augmented flow values are integers. □

# Networks with integer capacities

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem (Integrality theorem)

Let  $\mathcal{N} = (V, E, c, s, t)$  where  $c : E \rightarrow \mathbb{Z}^+$ . There exists a max-flow  $f^*$  such that  $f^*(e)$  is an integer, for any  $e \in E$ .

### Proof:

Since the algorithm terminates, the theorem follows from the integrality invariant lemma. □

# Networks with integer capacities: FF running time

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

*Let  $C$  be the min cut capacity ( $=\text{max. flow value}$ ),  
Ford-Fulkerson terminates after finding at most  $C$  augmenting  
paths.*

Proof: The value of the flow increases by  $\geq 1$  after each  
augmentation. □

# Networks with integer capacities: FF running time

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem

- The number of iterations is  $\leq C$ . At each iteration:
- Constructing  $G_f$ , with  $E(G_f) \leq 2m$ , takes  $O(m)$  time.
- $O(n + m)$  time to find an augmenting path, or deciding that it does not exist.
- Total running time is  $O(C(n + m)) = O(Cm)$
- Is that polynomial? No, only pseudopolynomial

# Networks with integer capacities: FF running time

The number of iterations of Ford-Fulkerson could be  $\Theta(C)$

- Max Flow and Min Cut
- Properties of flows and cuts
- Residual graph
- Augmenting path
- MaxFlow MinCut Thm
- Ford Fulkerson alg
- Maximum matching in Bip graphs
- Disjoint paths problem



Ford-Fulkerson can alternate between the two long paths, and require  $2M$  iterations. Taking  $M = 10^{10}$ , FF on a graph with 4 vertices can take time  $20^{10}$ .

# Networks with integer capacities: FF running time

The number of iterations of Ford-Fulkerson could be  $\Theta(C)$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual graph


Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem




Ford-Fulkerson can alternate between the two long paths, and require  $2M$  iterations. Taking  $M = 10^{10}$ , FF on a graph with 4 vertices can take time  $20^{10}$ .

# MAXIMUM MATCHING problem

Given an undirected graph  $G = (V, E)$  a subset of edges  $M \subseteq E$  is a **matching** if each node appears at most in one edge (a node may not appear at all).


A **perfect matching** in  $G$  is a matching  $M$  such that  $|M| = |V|/2$

The **MAXIMUM MATCHING** problem: Given as input a graph  $G$ , find a matching with maximum cardinality.




# Maximum matching in graphs bipartite

A graph  $G = (V, E)$  is said to be **bipartite** if  $V$  can be partitioned into  $L$  and  $R$ ,  $L \cup R = V$ ,  $L \cap R = \emptyset$ , such that every  $e \in E$  connects  $L$  with  $R$ .




The **MAXIMUM MATCHING BIPARTITE GRAPH** problem:  
Given as input a bipartite graph  $G = (L \cup R, E)$  with  $2n$  vertices, find a maximum matching.



# MAXIMUM MATCHING: flow formulation

Given a bipartite graph  $G = (L \cup R, E)$  construct  $\hat{G} = (\hat{V}, \hat{E})$ :


- Add vertices  $s$  and  $t$ :  $\hat{V} = L \cup R \cup \{s, t\}$ .
- Add directed edges  $s \rightarrow L$  with capacity 1. Add directed edges  $R \rightarrow t$  with capacity 1.
- Direct the edges  $E$  from  $L$  to  $R$ , and give them capacity  $\infty$ .
- $\hat{E} = \{s \rightarrow L\} \cup E \cup \{R \rightarrow t\}$ .



# MAXIMUM MATCHING: flow formulation

Given a bipartite graph  $G = (L \cup R, E)$  construct  $\hat{G} = (\hat{V}, \hat{E})$ :

- Add vertices  $s$  and  $t$ :  $\hat{V} = L \cup R \cup \{s, t\}$ .
- Add directed edges  $s \rightarrow L$  with capacity 1. Add directed edges  $R \rightarrow t$  with capacity 1.
- Direct the edges  $E$  from  $L$  to  $R$ , and give them capacity  $\infty$ .
- $\hat{E} = \{s \rightarrow L\} \cup E \cup \{R \rightarrow t\}$ .



# Maximum matching algorithm: Analysis


## Theorem

$\text{Max flow in } \hat{G} = \text{Max bipartite matching in } G.$

## Proof $\leq$

Given a matching  $M$  in  $G$  with  $k$ -edges,  
consider the flow  $F$  that sends 1 unit along each one of the  $k$  paths.

$f$  is a flow and has value  $k$ .




# Maximum matching algorithm: Analysis

Max flow  $\leq$  Max bipartite matching

- If there is a flow  $f$  in  $\hat{G}$ ,  $|f| = k$ , as capacities are  $\mathbb{Z}^*$   $\Rightarrow$  an integral flow exists.
- Consider the cut  $C = (\{s\} \cup L, R \cup \{t\})$  in  $\hat{G}$ .
- Let  $F$  be the set of edges in  $C$  with flow=1, then  $|F| = k$ .
- Each node in  $L$  is in at most one  $e \in F$  and every node in  $R$  is in at most one head of an  $e \in F$
- Therefore, exists a bipartite matching  $F$  in  $G$  with  $|F| \leq |f|$


□



# DISJOINT PATH problem

Given a digraph  $(G = (V, E), s, t)$ , a set of paths is **edge-disjoint** if their edges are disjoint (although them may go through some of the same vertices)


**DISJOINT PATH problem:** Given as input  $G, s, t$ , find the max number of edge disjoint paths  $s \rightsquigarrow t$



# DISJOINT PATH problem

Given a digraph  $(G = (V, E), s, t)$ , a set of paths is edge-disjoint if their edges are disjoint (although them may go through some of the same vertices)

The disjoint path problem given  $G, s, t$  find the max number of edge disjoint paths  $s \rightsquigarrow t$





# DISJOINT PATH: Max flow formulation

Assign unit capacity to every edge

## Theorem

*The max number of edge disjoint paths  $s \rightsquigarrow t$  is equal to the max flow value*



# DISJOINT PATH: Proof of the Theorem

Number of disjoint paths  $\leq$  max flow

If we have  $k$  edge-disjoint paths  $s \rightsquigarrow t$  in  $G$  then making  $f(e) = 1$  for each  $e$  in a path, we get a flow  $= k$

Number of disjoint paths  $\geq$  max flow

If max flow  $|f^*| = k \Rightarrow \exists$  0-1 flow  $f^*$  with value  $k$

$\Rightarrow \exists k$  edges  $(s, v)$  s.t.  $f(s, v) = 1$ , by **flow conservation** we can extend to  $k$  paths  $s \rightsquigarrow t$ , where each edge is a path carries flow  $= 1$ . □

If we have an undirected graph, with two distinguished nodes  $u, v$ , how would you apply the max flow formulation to solve the problem of finding the max number of disjoint paths between  $u$  and  $v$ ?