




Contents lists available at ScienceDirect

Injury

journal homepage: www.elsevier.com/locate/injury



Letter to the Editor

Towards a correct timing and dosage in PRP applications

Dear Editor,

A letter you have recently published makes reference to “Concerns about fibrosis development after scaffolded PRP therapy of muscle injuries” [1], and in our humble opinion, several errors require address.

In their reference to our article “Muscle repair: platelet-rich plasma derivatives as a bridge from spontaneity to intervention” [2] Robi and Matjaz make an inaccurate statement when they note “they combine PRP with a liquid-to-gel fibrin-based scaffold...”. In fact, there is no combination of two elements. Rather, the autologous enriched platelet plasma liquid has progressed to its more gel-like formulation which makes it a natural fibrin based scaffold in the infiltrated site. This transition from liquid PRGF (plasma rich in growth factors) formulation to gel form is accomplished with the simple addition of calcium chloride. In this form, platelets are activated and fibrinogen is cleaved, and in this way the polymerization of fibrin is achieved [3]. This liquid-to-gel fibrin-based scaffold conveys growth factors, cytokines, morphogens, and even exosomes contained in the platelets, as well as fibrinogen and other plasmatic proteins in a biologically balanced aggregate, managed and delivered in a physiological manner [4].

We would again point out to authors that PRGF muscle infiltrations are aimed at recruiting, activating and mobilizing satellite cells and resident macrophages which contribute to muscle reparation processes by cell signalling soluble factors in addition to the already activated endothelial cells, macrophages, and platelets in the injured area [5–7]. Furthermore, the gradual activation and homogeneous tissue distribution supported by a PRGF liquid-to-gel transition 3D (three dimensional) injectable scaffold may well serve as a highway for mechanical energy to transit from the environment to the cell, bring about cell-to-cell transition, promote multicellular assembly, provide mechanical support and plastic–elastic stiffness which has a drastic impact on the fate of muscle stem cells [8,9].

The author’s concern about the development of fibrosis mediated by TGF- β is widely shared with many other biologists and practitioners, and it is underpinned by ample research data in most of which TGF- β was applied alone [10]. Although transient fibrogenesis, collagen synthesis and deposition, mainly driven by the TGF- β family are crucial processes for structural and functional muscle regeneration, one of the most unsuccessful and nonfunctional outcomes of muscle regeneration is the formation of fibrotic scarring [11]. The application of PRGF in damaged tendons exhibited an absence of scarring [12,13] which might be attributed to the concurrent presence of TGF- β 1, VEGF, IGF-1 and HGF within PRGF thereby rendering this therapeutic system an anti-fibrotic formulation [14–17].

The authors mention the excellent study conducted by Reurink et al. [18] who reported no benefit for intramuscular PRP injections, as compared with isotonic saline injections, in patients with acute hamstring injuries. However, we have already suggested that the delayed application and low dosage may well have rendered PRP application ineffective as this initial window of 5 days is the time purported to shorten treatment, and appears to be crucial to muscle repair [19,20]. By the time of PRP application (a window of 5 days post injury), many of the injured microenvironmental biological targets of PRP, have either disappeared or undergone a phenotypic shift (myoblasts, endothelial cells, macrophages). Three in vivo studies, in which the treatment started either a few hours or two days after injury, reported histological acceleration or functional improvement in the treated group [21–23]. The total GF dosage applied at the site of injury throughout the treatment in these three studies was at least 2 times as high as that in the trial by Reurink et al. [18]. This is not caused by any difference in GF concentration of different PRPs (insulin-like growth factor 1 [IGF-1], 100,000 pg per millilitre versus 90,000 pg per millilitre, platelet-derived growth factor [PDGF], 20,000 pg per millilitre versus 20,000 pg per millilitre), but rather due to the fact that each application conveyed a higher GF dosage (2.5 millilitres versus 1 millilitre) and were greater in number (4–5 injections versus 2 injections).

Conflict of interest

EA and SP are scientists at BTI Biotechnology Institute ImasD, a biotechnology company that investigates the biological potential of PRGF.

References

- [1] Robi K, Matjaz V. Concerns about fibrosis development after scaffolded PRP therapy of muscle injuries: commentary on an article by Sanchez et al.: “Muscle repair: Platelet-rich plasma derivatives as a bridge from spontaneity to intervention”. *Injury* 2014.
- [2] Sanchez M, Anitua E, Delgado D, Sanchez P, Orive G, Padilla S. Muscle repair: platelet-rich plasma derivatives as a bridge from spontaneity to intervention. *Injury* 2014;45(Suppl. 4):S7–14.
- [3] Anitua E, Andia I, Ardanza B, Nurden P, Nurden AT. Autologous platelets as a source of proteins for healing and tissue regeneration. *Thromb Haemost* 2004;91:4–15.
- [4] Anitua E, Sanchez M, Orive G. Potential of endogenous regenerative technology for in situ regenerative medicine. *Adv Drug Deliv Rev* 2010;62:741–52.
- [5] Chazaud B, Brigitte M, Yacoub-Youssef H, Arnold L, Gherardi R, Sonnet C, et al. Dual and beneficial roles of macrophages during skeletal muscle regeneration. *Exerc Sport Sci Rev* 2009;37:18–22.
- [6] Grounds MD. Complexity of extracellular matrix and skeletal muscle regeneration. In: *Skeletal muscle repair and regeneration*. Springer; 2008. p. 269–302.
- [7] Shi X, Garry DJ. Muscle stem cells in development, regeneration, and disease. *Genes Dev* 2006;20:1692–708.
- [8] Discher DE, Mooney DJ, Zandstra PW. Growth factors, matrices, and forces combine and control stem cells. *Science* 2009;324:1673–7.
- [9] Gilbert PM, Havenstrite KL, Magnusson KE, Sacco A, Leonardi NA, Kraft P, et al. Substrate elasticity regulates skeletal muscle stem cell self-renewal in culture. *Science* 2010;329:1078–81.

- [10] Li Y, Foster W, Deasy BM, Chan Y, Prisk V, Tang Y, et al. Transforming growth factor-beta1 induces the differentiation of myogenic cells into fibrotic cells in injured skeletal muscle: a key event in muscle fibrogenesis. *Am J Pathol* 2004;164:1007–19.
- [11] Murphy MM, Lawson JA, Mathew SJ, Hutcheson DA, Kardon G. Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration. *Development* 2011;138:3625–37.
- [12] Sanchez M, Anitua E, Azofra J, Prado R, Muruzabal F, Andia I. Ligamentization of tendon grafts treated with an endogenous preparation rich in growth factors: gross morphology and histology. *Arthroscopy* 2010;26:470–80.
- [13] Anitua E, Sanchez M, Nurden AT, Zalduendo M, de la Fuente M, Azofra J, et al. Reciprocal actions of platelet-secreted TGF-beta1 on the production of VEGF and HGF by human tendon cells. *Plast Reconstr Surg* 2007;119:950–9.
- [14] Anitua E, Sanchez M, Merayo-Llones J, De la Fuente M, Muruzabal F, Orive G. Plasma rich in growth factors (PRGF-Endoret) stimulates proliferation and migration of primary keratocytes and conjunctival fibroblasts and inhibits and reverts TGF-beta1-induced myodifferentiation. *Investig Ophthalmol Vis Sci* 2011;52:6066–73.
- [15] Borselli C, Storrie H, Benesch-Lee F, Shvartsman D, Cezar C, Lichtman JW, et al. Functional muscle regeneration with combined delivery of angiogenesis and myogenesis factors. *Proc Natl Acad Sci U S A* 2010;107:3287–92.
- [16] Gong R. Multi-target anti-inflammatory action of hepatocyte growth factor. *Curr Opin Investig Drugs* 2008;9:1163–70.
- [17] Shavlakadze T, White J, Hoh JF, Rosenthal N, Grounds MD. Targeted expression of insulin-like growth factor-I reduces early myofiber necrosis in dystrophic mdx mice. *Mol Ther* 2004;10:829–43.
- [18] Reurink G, Goudswaard GJ, Moen MH, Weir A, Verhaar JA, Bierma-Zeinstra SM, et al. Platelet-rich plasma injections in acute muscle injury. *N Engl J Med* 2014;370:2546–7.
- [19] Dimauro I, Grasso L, Fittipaldi S, Fantini C, Mercatelli N, Racca S, et al. Platelet-rich plasma and skeletal muscle healing: a molecular analysis of the early phases of the regeneration process in an experimental animal model. *PLOS ONE* 2014;9:e102993.
- [20] Anitua E, Sanchez M, Padilla S. More on platelet-rich plasma injections in acute muscle injury. *N Engl J Med* 2014;371:1264.
- [21] Hammond JW, Hinton RY, Curl LA, Muriel JM, Lovering RM. Use of autologous platelet-rich plasma to treat muscle strain injuries. *Am J Sports Med* 2009;37:1135–42.
- [22] Wright-Carpenter T, Klein P, Schaferhoff P, Appell HJ, Mir LM, Wehling P. Treatment of muscle injuries by local administration of autologous conditioned serum: a pilot study on sportsmen with muscle strains. *Int J Sports Med* 2004;25:588–93.
- [23] Wright-Carpenter T, Opolon P, Appell HJ, Meijer H, Wehling P, Mir LM. Treatment of muscle injuries by local administration of autologous conditioned serum: animal experiments using a muscle contusion model. *Int J Sports Med* 2004;25:582–7.

S. Padilla^{a,1}

^aFoundation Eduardo Anitua, Vitoria, Spain

M. Sánchez

Arthroscopy Surgery Unit, Hospital Vithas San José, Vitoria, Spain

G. Orive*

Foundation Eduardo Anitua, Vitoria, Spain

E. Anitua^{a,b}

^aFoundation Eduardo Anitua, Vitoria, Spain

^bBiotechnology Institute (BTI), Vitoria, Spain

*Corresponding author

E-mail address: gorka.orive@ehu.es (G. Orive).

¹The authors declare the following competing financial interest(s): E. Anitua is the Scientific Director and S. Padilla are scientists at BTI Biotechnology Institute, a dental implant company, that investigates in the fields of oral implantology and PRGF-Endoret technology.