

Aufgabe 5

Sei $A = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix}$.

- Lassen Sie eine KI Ihrer Wahl, z.B. ChatGPT, mehrfach die Eigenwerte und Eigenvektoren von A berechnen.
- Bewerten Sie die KI-Lösungen.
- Kopieren Sie eine fehlerhafte KI-Antwort und markieren Sie die Fehler.
- Wie kann man erkennen, dass die Lösung falsch war, ohne die gesamte Lösung durchzugehen?

Lösung 5

Die Eigenwerte von A sind $\lambda_1 = -3$ und $\lambda_2 = 3$. Die Eigenvektoren sind

$$v_1 = \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad \text{und} \quad v_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Die folgende Ausgabe wurde am 14.06.2023 mit OpenAIs ChatGPT in der Version May 24 generiert. [1] Für eine bessere Darstellung in diesem Dokument wurden minimale Änderungen an der Formatierung vorgenommen und Umbrüche hinzugefügt. Die rot unterlegten Boxen wurden nachträglich hinzugefügt. Die Originalausgabe kann hier nachgelesen werden. [2].



Given the matrix $A = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix}$, compute its eigenvalues and eigenvectors step by step and check your results.



To compute the eigenvalues and eigenvectors of the matrix $A = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix}$, we follow these steps:



Step 1: Compute the characteristic equation. The characteristic equation is given by $\det(A - \lambda I) = 0$, where I is the identity matrix and λ is the eigenvalue we want to find.

Let's compute $A - \lambda I$:

$$A - \lambda I = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix} - \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} = \begin{pmatrix} 1 - \lambda & 2 \\ 4 & -1 - \lambda \end{pmatrix}.$$



Step 2: Calculate the determinant.
We need to find the determinant of $A - \lambda I$.

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 2 \\ 4 & -1 - \lambda \end{vmatrix} = (1 - \lambda)(-1 - \lambda) - (2 \cdot 4) = \lambda^2 - 2\lambda - 9.$$

Das charakteristische Polynom müsste $\lambda^2 - 9$ sein. Der Fehler wird in den nächsten Schritt übernommen, jedoch wird die quadratische Gleichung dennoch richtig gelöst und die Eigenwerte korrekt bestimmt:



Step 3: Solve the characteristic equation. Set the determinant equal to zero and solve for λ .

$$\lambda^2 - 2\lambda - 9 = 0$$

To solve this quadratic equation, we can factor it or use the quadratic formula.

Factoring it, we get:

$$(\lambda - 3)(\lambda + 3) = 0$$

So, the eigenvalues are $\lambda_1 = 3$ and $\lambda_2 = -3$.



Step 4: Compute the eigenvectors. To find the eigenvectors corresponding to each eigenvalue, we substitute the eigenvalues into the equation $(A - \lambda I) \cdot v = 0$ and solve for v .

For $\lambda_1 = 3$:

$$A - 3I = \begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix}$$

Reducing this matrix to row-echelon form, we get:

$$\begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

From this, we can see that $x_1 - x_2 = 0$.

Choosing $x_2 = t$ as a free variable, we get $x_1 = t$.

Therefore, the eigenvector corresponding to $\lambda_1 = 3$ is $v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$.



For $\lambda_2 = -3$:

$$A - (-3)I = \begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix}$$

Reducing this matrix to row-echelon form, we get:

$$\begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix}$$

From this, we can see that $2x_1 + x_2 = 0$.

Choosing $x_2 = t$ as a free variable, we get $x_1 = -\frac{1}{2}t$.

Therefore, the eigenvector corresponding to $\lambda_2 = -3$ is $v_2 = \begin{pmatrix} -\frac{1}{2} \\ 1 \end{pmatrix}$.



Step 5: Check the results. To check our results, we can multiply A by each eigenvector and see if we obtain the corresponding eigenvalue times the eigenvector.

For $\lambda_1 = 3$:

$$Av_1 = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \lambda_1 v_1$$

For $\lambda_2 = -3$:

$$Av_2 = \begin{pmatrix} 1 & 2 \\ 4 & -1 \end{pmatrix} \begin{pmatrix} -\frac{1}{2} \\ 1 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \end{pmatrix} = -3 \begin{pmatrix} -\frac{1}{2} \\ 1 \end{pmatrix} = \lambda_2 v_2$$

Both results confirm that the calculated eigenvalues and eigenvectors are correct.

Das Ergebnis der zweiten Prüfung müsste $\left(\frac{3}{2}, -3\right)^T$ lauten, aber die Schlussfolgerung ist korrekt.

Auf Rückfrage bzgl. des ersten Fehlers vermutet ChatGPT zunächst einen Vorzeichenfehler gemacht zu haben.



Are you sure about the -2λ in Step 2?



Apologies for the mistake. You are correct. In Step 2, the correct calculation should be:

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 2 \\ 4 & -1 - \lambda \end{vmatrix} = (1 - \lambda)(-1 - \lambda) - (2 \cdot 4) = \lambda^2 + 2\lambda - 9.$$

I apologize for the confusion caused by the incorrect sign in the previous response. Let me proceed with the correct calculation.

Using the corrected characteristic equation:

$$\lambda^2 + 2\lambda - 9 = 0$$

[...]

Erst mit dem folgenden Hinweis, kommt ChatGPT auf die richtige Fährte.



What if I told you, that the characteristic equation is in fact $\lambda^2 - 9 = 0$?



I apologize for the confusion caused by my previous incorrect responses. You are absolutely right. The characteristic equation for the matrix A should indeed be $\lambda^2 - 9 = 0$.

Let's continue with the computation of eigenvectors using the corrected characteristic equation.

For $\lambda_1 = 3$:

$$A - 3I = \begin{pmatrix} 1 - 3 & 2 \\ 4 & -1 - 3 \end{pmatrix} = \begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix}$$

Reducing this matrix to row-echelon form, we have:

$$\begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

From this, we can see that $x_1 - x_2 = 0$.

Choosing $x_2 = t$ as a free variable, we get $x_1 = t$.

Therefore, the eigenvector corresponding to $\lambda_1 = 3$ is $v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$.



For $\lambda_2 = -3$:

$$A - (-3)I = \begin{pmatrix} 1+3 & 2 \\ 4 & -1+3 \end{pmatrix} = \begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix}$$

Reducing this matrix to row-echelon form, we have:

$$\begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix}$$

From this, we can see that $2x_1 + x_2 = 0$.

Choosing $x_2 = -2t$ as a free variable, we get $x_1 = t$.

Therefore, the eigenvector corresponding to $\lambda_2 = -3$ is $v_2 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$.

Again, I apologize for the previous mistakes, and thank you for pointing them out.

Auch diese Ergebnisse kann man, wie ChatGPT in Schritt 5 richtig beschrieben hat, wie folgt überprüfen:

$$Av_i \stackrel{?}{=} \lambda_i v_i$$

Ob die Argumentation und die Zwischenschritte jedoch richtig sind, kann man allein durch die Prüfung der Endergebnisse nicht herausfinden. Wie Turpin et al. [3] gezeigt haben, dürfen die Erklärungen beim Chain-of-Thought Prompting auch nicht mit den tatsächlichen Gründen verwechselt werden, aus denen LLM eine Ausgabe generiert.

Aufgabe 6

Bestimmen Sie die Eigenwerte und Eigenvektoren der Matrizen

$$\text{a) } A = \begin{pmatrix} 2 & 4 \\ 1 & 2 \end{pmatrix} \qquad \text{b) } B = \begin{pmatrix} 3 & -1 \\ 2 & 5 \end{pmatrix}$$

Bestimmen Sie auch eventuelle komplexe Eigenwerte und Eigenvektoren. Ein komplexes Gleichungssystem können Sie wie ein reelles Gleichungssystem mit dem Gauß-Verfahren lösen.

Lösung 6

Aufgabe 7

Eine Abbildung im \mathbb{R}^2 ist wie folgt definiert: Ein gegebener Vektor x wird zuerst um 90° im Uhrzeigersinn gedreht und anschließend an der y -Achse gespiegelt.

- a) Das Vielfache welchen Vektors wird auf sich selbst abgebildet?

- b) Die Abbildung welchen Vektors (bzw. des Vielfachen diesen Vektors) zeigt genau in die entgegengesetzte Richtung?
- c) Stellen Sie die zugehörige Abbildungsmatrix A auf.
- d) Bestimmen Sie die Eigenwerte von A . Passen diese zu den Beobachtungen aus a) und b)?

Lösung 7

Aufgabe 8

Bei welchen Werten a, b hat die Matrix

$$A = \begin{pmatrix} a & -1 \\ 1 & b \end{pmatrix}$$

- a) zwei verschiedene reelle Eigenwerte?
- b) einen (doppelten) reellen Eigenwert?
- c) keine reellen Eigenwert?

Lösung 8

Literatur

- [1] OpenAI, *ChatGPT*, <https://openai.com>, Version May 24, accessed on 2023-06-14, 2021.
- [2] C. R. Thelen, *Conversation with ChatGPT: Eigenvalues & Eigenvectors*, Chat with OpenAI's ChatGPT model, Version May 24, accessed on 2023-06-14, Juni 2023. Adresse: <https://chat.openai.com/share/c48bc91a-607c-4f7f-ba5c-b763724586ce>.
- [3] M. Turpin, J. Michael, E. Perez und S. R. Bowman, *Language Models Don't Always Say What They Think: Unfaithful Explanations in Chain-of-Thought Prompting*, 2023. arXiv: [2305.04388](https://arxiv.org/abs/2305.04388) [cs.CL].