

Dark Matter Module: the physics behind

Contents

1	Dark Matter	1
1.1	Galaxy rotational velocity curve	1
1.2	Velocity distribution in galaxy clusters	5

1 Dark Matter

- **Target topics:** Dark Matter (galaxy rotation curves, clusters).
- **Learning objectives:** Analyze multiple lines of evidence for dark matter; derive galaxy rotational velocity curve from mechanics, Newton gravitation law and Kepler law; investigate galaxies and clusters composition; find the cluster mass from virial theorem reviewing kinetic energy and gravitational potential; connect classical Physics to Cosmology.
- **Physics program link:** Mechanics, Kepler's law, Newton's law of gravitation, centripetal acceleration, circular motion, centripetal force, gravitational force, energy and momentum conservation, gravitational potential energy, kinetic energy.

1.1 Galaxy rotational velocity curve

- **Objectives:** Apply the law of gravitation to derive the expected orbital velocity in a galaxy. Compare the theoretical velocity with observational data from galaxy datasets. Introduce the dark matter hypothesis as a potential solution to the discrepancy.
- **Dataset variables of galaxies:** [11]
 - Rad (radius)
 - V_{obs} (observed velocity)
 - $\text{err}V$ (velocity error)
 - V_{gas} (gas velocity)
 - V_{disk} (stellar disk velocity)
 - V_{bul} (bulge velocity)

- SB_{disk} (surface brightness disk)
- SB_{bul} (surface brightness bulge)

Classical Physics derivation:

- We model a small mass m (orbiting body) in a stable circular orbit around a large central mass M (e.g., galactic center) at a radius r .
- The condition for stable orbit requires the Gravitational Force (F_G) to equal the Centripetal Force (F_c).

Gravitational and Centripetal Forces

- **Gravitational Force (F_G):**

$$F_G = \frac{GMm}{r^2} \quad (1)$$

- **Centripetal Force (F_c):**

$$F_c = \frac{mv^2}{r} \quad (2)$$

- **Equating the Forces ($F_G = F_c$):**

$$\frac{GMm}{r^2} = \frac{mv^2}{r} \quad (3)$$

Deriving the rotational velocity (v)

- **Cancellation of m :** The mass of the orbiting body (m) cancels out, showing v is independent of m .

$$\frac{GM}{r^2} = \frac{v^2}{r} \quad (4)$$

- **Isolating v^2 :** Multiply by r to isolate v^2 .

$$v^2 = \frac{GM}{r} \quad (5)$$

- **Final Rotational Velocity (v):** Take the square root.

$$v = \sqrt{\frac{GM}{r}} \quad (6)$$

- **Physical Meaning:** This Keplerian velocity profile dictates that $v \propto 1/\sqrt{r}$, meaning orbital speed decreases as distance from the central mass increases.

Connection to Kepler's third law

- **Orbital Period (T):** Period is the circumference divided by velocity.

$$T = \frac{2\pi r}{v} \quad (7)$$

- **Kepler's relation :** Square T and substitute $v^2 = GM/r$.

$$T^2 = \frac{4\pi^2 r^2}{v^2} = \frac{4\pi^2 r^2}{GM/r} \implies T^2 = \frac{4\pi^2}{GM} r^3 \quad (8)$$

- **Physical Meaning:** This confirms Kepler's Third Law ($T^2 \propto r^3$), demonstrating that the proportionality derived from the force balance is fully consistent with the relationship between orbital period and radius.

Velocity and mass

- **Baryonic velocity (from data):** Incorporating the Mass-to-Light ratio (Υ) and conserving the signs for counter-rotating components [10]:

$$V_{\text{bar}}^2(r) = V_{\text{gas}}|V_{\text{gas}}| + \Upsilon [V_{\text{disk}}|V_{\text{disk}}| + V_{\text{bul}}|V_{\text{bul}}|] \quad (9)$$

- **Baryonic mass:**

$$M_{\text{bar}}(r) = \frac{r \max(V_{\text{bar}}^2(r), 0)}{G} \quad (10)$$

- **Total observed mass:**

$$M_{\text{obs}}(r) = \frac{r V_{\text{obs}}^2(r)}{G} \quad (11)$$

- **Dark matter velocity squared:** Derived from the difference between the observed total velocity and the baryonic prediction:

$$V_{\text{DM}}^2(r) = V_{\text{obs}}^2(r) - V_{\text{bar}}^2(r) \quad (12)$$

- **Dark matter density (from data):**

$$\rho_{\text{DM}}(r) = \frac{1}{4\pi G r^2} \frac{d}{dr} (r \cdot V_{\text{DM}}^2(r)) \quad (13)$$

NFW Dark Matter Profile

- **NFW profile for dark matter [13, 12]:**

$$\rho_{\text{NFW}}(r) = \frac{\rho_s}{(r/r_s)(1 + r/r_s)^2} \quad (14)$$

where r_s is the scale radius and ρ_s is the characteristic density, $\alpha = 1, \beta = 3, \gamma = 1$.

- **Global Fit parameters:** The structural parameters ρ_s , r_s , and the mass-to-light ratio Υ are calculated via a global χ^2 minimization on the entire rotation curve dataset to provide a robust estimation of the halo.

- **NFW dark matter mass:**

$$M_{\text{NFW}}(r) = 4\pi \int \rho_{\text{NFW}} r^2 dr = 4\pi \rho_s r_s^3 \left[\ln \left(1 + \frac{r}{r_s} \right) - \frac{r/r_s}{1 + r/r_s} \right] \quad (15)$$

- **Velocity dark matter:**

$$V_{\text{DM}}(r) = \sqrt{\frac{GM_{\text{NFW}}(r)}{r}} \quad (16)$$

- **Total simulated curve:**

$$V_{\text{sim}}(r, f) = \sqrt{\frac{GM_{\text{bar}}(r) + f \cdot M_{\text{NFW}}(r)}{r}} \quad \text{with } f \in [0, 2] \quad (17)$$

Data visualization and fit

- **Plot:**

- **X-axis:** Rad (radius, data)
- **Y-axis:**
 - * V_{bar} (red)
 - * V_{obs} (data, blue) $\pm \text{errV}$
 - * V_{sim} (green)

- **Fit χ^2/dof (chi-squared):**

$$\chi^2/\text{dof} = \frac{1}{N - P} \sum_{i=1}^N \left(\frac{V_{\text{obs}}(r_i) - V_{\text{sim}}(r_i)}{\text{errV}(r_i)} \right)^2 \quad (18)$$

where N is the number of data points and P is the number of free parameters ($P = 3$: ρ_s, r_s, Υ).

Global χ^2 Minimization

The parameters r_s , ρ_s , and the mass-to-light ratio Υ are determined simultaneously by minimizing the χ^2 over the entire rotation curve:

$$\chi^2 = \sum_{i=1}^N \left(\frac{V_{\text{obs}}(r_i) - V_{\text{tot,sim}}(r_i)}{\text{errV}(r_i)} \right)^2 \quad (19)$$

where $V_{\text{tot,sim}}(r) = \sqrt{\max(V_{\text{bar}}^2(r) + V_{\text{DM}}^2(r), 0)}$.

1.2 Velocity distribution in galaxy clusters

- **Objectives:** Help students infer the existence of dark matter in galaxy clusters using classical mechanics and virial theorem. Students will apply the virial theorem to estimate the total mass of a galaxy cluster from the kinetic and potential energy of its galaxies.
- **Dataset Coma cluster (SDSS) :** objid (galaxy ID), ra (right ascension, α), dec (declination, δ), modelmag (apparent magnitude in r band, m_r), modelmagerr, extinction, redshift (z), zErr (redshift error) [14, 1].
- **Dataset Abell Clusters:** Cluster (name), ID (galaxy ID), RAdeg (right ascension in degrees), DEdeg (declination in degrees), RV (radial velocity in km/s), e_RV (velocity error in km/s), q_RV (quality flag), bmag (apparent magnitude in B band). [15]

1. **Individual Galaxy Velocity:** Assuming non-relativistic speeds for low-redshift clusters, the radial velocity of each galaxy is derived directly from its observed redshift data:

$$v_i = c z_i \quad (20)$$

2. **Number of observed galaxies:** N

3. **Mean Cluster Velocity:** The systemic velocity of the cluster is calculated as the average of the individual radial velocities:

$$\bar{v} = \frac{1}{N} \sum_{i=1}^N v_i \quad (21)$$

4. **Observed Velocity Dispersion:** The 1-D velocity dispersion is the standard deviation of the galaxies' velocities relative to the systemic mean, providing a measure of the cluster's dynamical state:

$$\sigma_{\text{obs}} = \sqrt{\frac{1}{N} \sum_{i=1}^N (v_i - \bar{v})^2} \quad (22)$$

5. **Comoving Distance:** To determine physical scales, we integrate over the expansion history of the Universe (Eq. 15 [8]):

$$\chi(z) = \frac{c}{H_0} \int_0^z \frac{dz'}{\sqrt{\Omega_m(1+z')^3 + (1-\Omega_m)}} \quad (23)$$

6. **Angular Diameter Distance:** This converts the comoving distance into a metric that relates apparent angular size to true physical size (Eq. 18 [8]):

$$D_A(z) = \frac{\chi(z)}{1+z} \quad (24)$$

7. **Cluster Center Identification:** The dynamical center is approximated using either the coordinates of the Brightest Cluster Galaxy (BCG) or the spatial median of all cluster members

[2, 4]:

$$(\alpha_c, \delta_c) = (\alpha_{\text{BCG}}, \delta_{\text{BCG}}) \quad \text{or} \quad (\text{median}(\alpha), \text{median}(\delta))$$

8. **Angular Separation (θ):** Applying the spherical law of cosines, we compute the angular distance between the i -th galaxy and the cluster center [5]. Note that right ascension (α) and declination (δ) must be converted to radians:

$$\theta_i = \arccos(\sin \delta_c \sin \delta_i + \cos \delta_c \cos \delta_i \cos(\alpha_c - \alpha_i)) \quad (25)$$

9. **Projected Physical Radius:** The angular separation is converted into a transverse physical distance using the angular diameter distance:

$$r_{\text{proj},i} = \theta_i D_{A,\text{kpc}} \quad (26)$$

10. **Critical Density:** The density required for a spatially flat Universe:

$$\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G} \quad (27)$$

11. **Virial Mass Estimation (M_{200}):** Assuming the cluster is a relaxed, dynamically thermalized system, the virial theorem links the observed velocity dispersion to the total mass enclosed within the characteristic radius R_{200} [4]:

$$\begin{aligned} \sigma_{\text{obs}}^2 &= \frac{GM_{200}}{3R_{200}} \\ M_{200} &= \frac{4}{3}\pi 200 \rho_{\text{crit}} R_{200}^3 \end{aligned} \quad (28)$$

Dark Matter (NFW) Profile

12. **Halo Concentration Parameter:** Derived from empirical scaling relations based on cosmological simulations (Eq. 4 [6]):

$$c = A \left(\frac{M_{200}}{M_{\text{pivot}}} \right)^B (1+z)^C \quad (29)$$

where $A = 5.71$, $B = -0.084$, and $C = -0.47$.

13. **NFW Density Characteristic Factor:** An overdensity contrast relative to the critical density (Eq. 2 [13]):

$$\delta_c(c) = \frac{200}{3} \frac{c^3}{\ln(1+c) - c/(1+c)} \quad (30)$$

14. **Characteristic Radius (r_{200}):**

$$r_{200} = \left(\frac{3M_{200}}{4\pi \cdot 200 \rho_{\text{crit}}} \right)^{1/3} \quad (31)$$

15. **Characteristic Density and Scale Radius:**

$$\rho_s = \delta_c(c) \rho_{\text{crit}}, \quad r_s = \frac{r_{200}}{c}$$

16. **Enclosed Dark Matter Mass (NFW):** The total mass of dark matter contained within the projected radius $r_{\text{proj},i}$:

$$M_{\text{NFW}}(r_{\text{proj},i}) = M_{200} \frac{\ln(1+x_i) - x_i/(1+x_i)}{\ln(1+c) - c/(1+c)}, \quad x_i = \frac{r_{\text{proj},i}}{r_s} \quad (32)$$

Luminous Mass

17. **Luminosity Distance ($z \ll 1$):** For local clusters, we can approximate the luminosity distance using the linear Hubble law evaluated at the median redshift of the cluster [5]:

$$D_L \approx \frac{cz_{\text{cluster}}}{H_0}, \quad z_{\text{cluster}} = \text{median}(z_i) \quad (33)$$

18. **Distance Modulus (μ):** The difference between apparent and absolute magnitude, where D_L is expressed in parsecs (Eq. 25 [8]):

$$\mu = 5 \log_{10}(D_L) - 5 \quad (34)$$

19. **Absolute Magnitude (M_r):** Derived from the apparent magnitude in the r-band (m_r), corrected for the distance modulus and the line-of-sight Galactic extinction (A_r) [5]:

$$M_r = m_r - A_r - \left[5 \log_{10} \left(\frac{D_L}{10 \text{ pc}} \right) \right] \quad (35)$$

20. **Galaxy Luminosity (L_r):** Converted from absolute magnitude and expressed in units of Solar Luminosity ($L_{\odot}$):

$$L_r = 10^{0.4(M_{r\odot} - M_r)} \quad (36)$$

21. **Stellar Mass per Galaxy ($M_{\text{star},i}$):** The stellar mass is estimated by applying a theoretical Mass-to-Light ratio ($\Upsilon_r = M/L \approx 2$) typical for early-type galaxies in clusters [3]:

$$M_{\text{star},i} = \Upsilon_r L_r \quad (37)$$

22. **Total Stellar Mass:** The sum of the individual stellar masses of all observed cluster members:

$$M_{\text{star}} = \sum_{i=1}^N M_{\text{star},i} \quad (38)$$

23. **Intracluster Gas Mass:** The X-ray emitting hot gas constitutes a major fraction of the cluster's baryons. This is estimated using empirical scaling relations from observations (Section 3 [7]) 1.2:

$$M_{\text{gas}} = M_{500} \cdot 0.093 \left(\frac{M_{500}}{2 \cdot 10^{14} M_{\odot}} \right)^{0.21} \quad (39)$$

(where $M_{500} \approx 0.7 \cdot M_{200}$ 1.2 and $h = 0.7$)

24. **Total Baryonic (Luminous) Mass:** The sum of the stellar component and the intracluster medium (gas):

$$M_{\text{lum}} = M_{\text{star}} + M_{\text{gas}} \quad (40)$$

25. **Characteristic Baryonic Radius (r_{bar}):** Defined analogously to r_{200} :

$$r_{\text{bar}} = \left(\frac{3M_{\text{lum}}}{4\pi \cdot 200 \rho_{\text{crit}}} \right)^{1/3} \quad (41)$$

26. **Baryonic Velocity Dispersion:** The theoretical velocity dispersion predicted if the cluster's dynamics were governed solely by visible baryonic matter (no dark matter):

$$\sigma_{\text{bar}} = \sqrt{\frac{GM_{\text{lum}}}{3r_{\text{bar}}}} \quad (42)$$

Velocity Dispersion and Simulation

25. **Total simulated mass:**

$$M_{\text{tot}} = M_{\text{lum}} + f M_{200} \quad (43)$$

26. **Total simulated velocity dispersion:**

$$\sigma_{\text{sim}} = \sqrt{\frac{GM_{\text{tot}}}{3r_{200}}} \quad (44)$$

27. **Observed histogram:**

$$H_{\text{obs}} = \text{hist}(v_{\text{obs}})$$

28. **Generate normal distribution of simulated velocities:**

$$\sigma_{\text{tot}} = \text{Normal}(\mu = \sigma_{\text{sim}}, \sigma = \sigma_{\text{obs}}, N)$$

29. **Simulated histogram:**

$$H_{\text{sim}} = \text{hist}(\sigma_{\text{tot}})$$

Data visualization

- Plot histograms: number of galaxies on Y-axis, velocity on X-axis.
- Curves: observations (blue), simulated (green), baryonic (red).

Cluster Mass via Virial Theorem

1. **Angular Separation** ($\Delta\theta$) from right ascension (α) and declination (δ) data (converted to radians, where c denotes the cluster center and i the individual galaxy):

$$\Delta\theta = \arccos [\sin(\delta_c) \sin(\delta_i) + \cos(\delta_c) \cos(\delta_i) \cos(\alpha_c - \alpha_i)] \quad (45)$$

2. **Projected Radius** (R_P) from angular separation, assuming a distance $D = 100$ Mpc (radius converted to kpc):

$$R_P = D \cdot \Delta\theta \quad (46)$$

3. **Observed Velocity** (V_{obs}) from redshift data (or radial velocity RV):

$$V_{\text{obs}} = c \cdot z \quad \text{or} \quad V_{\text{obs}} = \text{RV} \quad (47)$$

4. **1-D Velocity Dispersion** ($\sigma_{1\text{D}}$) (observation):

$$\sigma_{1\text{D}} = \sqrt{\frac{1}{N} \sum_{i=1}^N (V_i - \bar{V})^2} \quad (48)$$

5. **3-D Velocity Dispersion** ($\sigma_{3\text{D}}$) (assuming isotropy):

$$\sigma_{3\text{D}} = 3 \sigma_{1\text{D}} \quad (49)$$

Baryonic components

6. **Convert Distance** D (in Mpc) to pc and compute the **Distance Modulus** (μ):

$$\mu = 5 \log_{10} \left(\frac{D_{\text{pc}}}{10 \text{ pc}} \right) \quad (50)$$

7. **Absolute Magnitude** (M) using apparent magnitude ($m = \text{bmag}$):

$$M = m - \mu \quad (51)$$

8. **Luminosity** (L) of each galaxy from absolute magnitude (M):

$$L = 10^{-0.4(M-M_{\odot})} \quad (52)$$

9. **Total Luminosity of the cluster** (L_{total}):

$$L_{\text{total}} = \sum_{i=1}^N L_i \quad (53)$$

10. **Stellar Mass** (M_{lum}) from luminosity (Υ is the mass-to-light ratio):

$$M_{\text{lum}} = \Upsilon \cdot L_{\text{total}} \quad (54)$$

11. **Gas mass:**

$$M_{\text{gas}} = M_{500} \cdot 0.093 \left(\frac{M_{500}}{2 \cdot 10^{14} M_{\odot}} \right)^{0.21} \quad (55)$$

(where $M_{500} \approx 0.7 \cdot M_{200}$ and $h = 0.7$) 1.2 1.2

12. **Total Baryonic Mass:**

$$M_{\text{lum}} = M_{\text{star}} + M_{\text{gas}} \quad (56)$$

Notes:

The formula used to estimate the intracluster gas mass derived from Equation 12 in Giodini et al. (2009) [7].

Step 1: Definition of the gas mass fraction

By definition, the gas mass fraction within the radius R_{500} is the ratio of the gas mass to the total mass enclosed within that same radius:

$$f_{500}^{\text{gas}} = \frac{M_{\text{gas}}}{M_{500}} \quad (57)$$

From this definition, the gas mass can be isolated as:

$$M_{\text{gas}} = M_{500} \cdot f_{500}^{\text{gas}} \quad (58)$$

Step 2: Giodini's empirical relation (Eq. 12)[7]

An empirical scaling relation to calculate the gas fraction f_{500}^{gas} as a function of the total mass M_{500} :

$$f_{500}^{\text{gas}} \left(\frac{h}{0.7} \right)^{3/2} = (9.3 \times 10^{-2}) \left(\frac{M_{500}}{2 \times 10^{14} M_{\odot}} \right)^{0.21} \quad (59)$$

Step 3: Applying the Hubble parameter ($h = 0.7$)

The dimensionless Hubble parameter is set to $h = 0.7$. Substituting this value into the left side of Giodini's equation simplifies the scaling factor:

$$\left(\frac{0.7}{0.7}\right)^{3/2} = 1^{3/2} = 1 \quad (60)$$

Consequently, the equation for the gas mass fraction reduces to:

$$f_{500}^{\text{gas}} = 0.093 \left(\frac{M_{500}}{2 \times 10^{14} M_{\odot}} \right)^{0.21} \quad (61)$$

Step 4: Gas mass

Substituting the simplified gas fraction expression (Eq. 61) back into the initial mass definition (Eq. 58) yields the structure of the applied formula:

$$M_{\text{gas}} = M_{500} \cdot 0.093 \left(\frac{M_{500}}{2 \cdot 10^{14} M_{\odot}} \right)^{0.21} \quad (62)$$

Notes:

The mathematical derivation to obtain the conversion factor between M_{500} and M_{200} relies on the analytical properties of the Navarro-Frenk-White (NFW) density profile.

Step 1: The NFW density profile and enclosed mass

The NFW density profile is defined as:

$$\rho(r) = \frac{\rho_s}{\frac{r}{r_s} \left(1 + \frac{r}{r_s}\right)^2} \quad (63)$$

where ρ_s is the characteristic density and r_s is the scale radius. Integrating this density profile over a spherical volume up to a radius R gives the enclosed mass $M(R)$:

$$M(R) = 4\pi\rho_s r_s^3 \left[\ln \left(1 + \frac{R}{r_s}\right) - \frac{R/r_s}{1 + R/r_s} \right] \quad (64)$$

By defining the concentration parameter as $c = R/r_s$, the mass function is usually written as $M(c) \propto f(c)$, where:

$$f(c) = \ln(1 + c) - \frac{c}{1 + c} \quad (65)$$

Step 2: Definition of mass by overdensity

In cosmological models, the mass of a halo is defined as the mass enclosed within a radius R_{Δ} where the average internal density is Δ times the critical density of the Universe (ρ_{crit}). The mass is given by:

$$M_{\Delta} = \frac{4\pi}{3} \Delta \rho_{\text{crit}} R_{\Delta}^3 \quad (66)$$

Applying this definition to $\Delta = 500$ and $\Delta = 200$, we can express the ratio of the two masses geometrically:

$$\frac{M_{500}}{M_{200}} = \frac{500}{200} \left(\frac{R_{500}}{R_{200}} \right)^3 = 2.5 \left(\frac{c_{500}}{c_{200}} \right)^3 \quad (67)$$

where $c_{500} = R_{500}/r_s$ and $c_{200} = R_{200}/r_s$.

Step 3: Equating the NFW Mass and the overdensity mass

The mass ratio must also satisfy the NFW integrated mass equation from Step 1:

$$\frac{M_{500}}{M_{200}} = \frac{f(c_{500})}{f(c_{200})} = \frac{\ln(1 + c_{500}) - \frac{c_{500}}{1+c_{500}}}{\ln(1 + c_{200}) - \frac{c_{200}}{1+c_{200}}} \quad (68)$$

Equating the geometric ratio (Step 2) to the NFW mass ratio yields the conversion equation:

$$2.5 \left(\frac{c_{500}}{c_{200}} \right)^3 = \frac{\ln(1 + c_{500}) - \frac{c_{500}}{1+c_{500}}}{\ln(1 + c_{200}) - \frac{c_{200}}{1+c_{200}}} \quad (69)$$

Step 4: Solving for a fixed concentration ($c_{200} = 5$)

Assuming a concentration parameter $c_{200} = 5$, we evaluate the denominator of the right side:

$$f(5) = \ln(1 + 5) - \frac{5}{1 + 5} = \ln(6) - \frac{5}{6} \approx 1.7917 - 0.8333 = 0.9584 \quad (70)$$

The equation becomes:

$$2.5 \left(\frac{c_{500}}{5} \right)^3 = \frac{\ln(1 + c_{500}) - \frac{c_{500}}{1+c_{500}}}{0.9584} \quad (71)$$

Solving this equation numerically for the unknown c_{500} yields:

$$c_{500} \approx 3.305 \quad (72)$$

Step 5: Mass ratio calculation

Substitute $c_{500} \approx 3.305$ back into the geometric mass ratio equation:

$$\frac{M_{500}}{M_{200}} = 2.5 \left(\frac{3.305}{5} \right)^3 = 2.5 \times (0.661)^3 \approx 2.5 \times 0.2888 \approx 0.722 \quad (73)$$

Therefore, for a halo with a concentration $c = 5$, the mass M_{500} is approximately **72.2%** of M_{200} , which justifies the standard approximation $M_{500} \approx 0.7 \cdot M_{200}$. **References in the**

Literature:

- **White (2001) [16]:** This paper provides the analytical background for comparing mass definitions and explicitly plots the ratio M_i/M_{200} for $c = 5$. In Figure 1, the curve corresponding to M_{500} is a horizontal line constant at ≈ 0.72 independent of Ω_m .
- **Hu & Kravtsov (2003) [9]:** Appendix C ("Mass Conversion") of this paper details

the analytical transformation between different overdensity mass definitions assuming an NFW profile. Equations (C1) to (C3) define the NFW enclosed mass function, and Equation (C10) provides the generalized geometric mass ratio required for the conversion.

Virial Theorem Derivation

11. **Gravitational Potential Energy** (light particle m orbiting heavy particle M):

$$U = -\frac{GMm}{r} \quad (74)$$

12. **Gravitational Force:**

$$F_G = \frac{GMm}{r^2} \quad (75)$$

13. **Centripetal Force:**

$$F_c = \frac{mv^2}{r} \quad (76)$$

14. **Equating Forces** ($F_G = F_c$) to find v^2 :

$$v^2 = \frac{GM}{r} \quad (77)$$

15. **Kinetic Energy** (K , neglecting heavy particle's kinetic energy):

$$K = \frac{1}{2}mv^2 \quad (78)$$

16. **Replacing v^2 in K :**

$$K = \frac{1}{2}m \left(\frac{GM}{r} \right) \quad (79)$$

17. **Replacing Potential Energy U in K :**

$$K = -\frac{1}{2}U \quad (80)$$

18. **Virial Theorem:**

$$2 \langle K \rangle + \langle U \rangle = 0 \quad (81)$$

Cluster Virial Mass and Density

- **Context:** In a cluster, the total potential energy is the sum of all pairwise interactions, and the total kinetic energy is the sum of the kinetic energies of all galaxies. The Virial Theorem relates these totals. For an isotropic and stable system, the velocity is replaced by the velocity dispersion (σ).

19. **Total Virial Mass of the cluster** (M_{Virial}) (using σ_{3D} and characteristic radius R):

$$M_{\text{Virial}} = \frac{3 \sigma_{1D}^2 R}{G} \quad (82)$$

20. **Total Volume** (approximated as a sphere):

$$\text{Volume} = \frac{4}{3} \pi R^3 \quad (83)$$

21. **Luminous Density** (ρ_{lum}):

$$\rho_{\text{lum}} = \frac{M_{\text{lum}}}{\text{Volume}} \quad (84)$$

22. **Total Density** (ρ_{total}):

$$\rho_{\text{total}} = \frac{M_{\text{total}}}{\text{Volume}} \quad (85)$$

Data visualization

- **X-Axis:** Radius (R).
- **Y-Axis (Quantities to Plot):**
 1. ρ_{lum} (Luminous Density, Red)
 2. ρ_{total} (Total Density, Data Blue)
 3. V_{sim} (Total Simulated Velocity, Red)
 4. V_{obs} (Observed Velocity, Blue data points)

References

- [1] Mert Alkan. *Coma Cluster Dataset*. <https://www.kaggle.com/datasets/mertalkan98/coma-cluster>. 2021.
- [2] Neta A. Bahcall and Craig L. Sarazin. “Cluster Velocity Dispersions and Masses”. In: *The Astrophysical Journal* 258 (1982). virial of clusters and application with velocity dispersion, pp. 1–9. DOI: 10.1086/160050.
- [3] E. F. Bell et al. “The Optical and Near-Infrared Properties of Galaxies: I. Luminosity and Stellar Mass Functions”. In: *The Astrophysical Journal Supplement Series* 149 (2003), pp. 289–312.
- [4] R.G. Carlberg et al. “Galaxy Cluster Virial Masses and Omega”. In: *apj* 462 (May 1996), p. 32. DOI: 10.1086/177125. arXiv: astro-ph/9509034 [astro-ph].
- [5] B. W. Carroll and D. A. Ostlie. *An Introduction to Modern Astrophysics*. Cambridge University Press.

- [6] A. R. Duffy et al. “Dark Matter Halo Concentrations in the Wilkinson Microwave Anisotropy Probe Year 5 Cosmology”. In: *Monthly Notices of the Royal Astronomical Society* 390 (2008), pp. L64–L68.
- [7] S. Giodini et al. “Stellar and Total Baryon Mass Fractions in Groups and Clusters Since Redshift 1”. In: *The Astrophysical Journal* 703.1 (2009), pp. 982–993. DOI: 10.1088/0004-637X/703/1/982. URL: <https://doi.org/10.1088/0004-637X/703/1/982>.
- [8] D. W. Hogg. “Distance measures in cosmology”. In: *arXiv:astro-ph/9905116* (1999). Sec 2.3: relation redshift and radial velocity. URL: <https://arxiv.org/abs/astro-ph/9905116>.
- [9] Wayne Hu and Andrey V. Kravtsov. “Sample Variance Considerations for Cluster Surveys”. In: *The Astrophysical Journal* 584.2 (Feb. 2003), pp. 702–715. DOI: 10.1086/345846.
- [10] E. V. Karukes and P. Salucci. “The Universal Rotation Curve of Dwarf Disk Galaxies”. In: *Monthly Notices of the Royal Astronomical Society* 465 (2016), pp. 4703–4722.
- [11] Federico Lelli, Stacy S. McGaugh, and James M. Schombert. “SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves”. In: *aj* 152.6, 157 (Dec. 2016). <https://astroweb.cwru.edu/SPARC/>, p. 157. DOI: 10.3847/0004-6256/152/6/157. arXiv: 1606.09251 [astro-ph.GA].
- [12] J. F. Navarro, C. S. Frenk, and S. D. M. White. “The Structure of Cold Dark Matter Halos”. In: *The Astrophysical Journal* 462 (1996), p. 563.
- [13] J. F. Navarro, C. S. Frenk, and S. D. M. White. “A Universal Density Profile from Hierarchical Clustering”. In: *The Astrophysical Journal* 490 (1997), p. 493.
- [14] SDSS Collaboration. *Sloan Digital Sky Survey: Science and Data-Release Publications*. Sloan Digital Sky Survey. 2023. URL: <https://www.sdss.org/science/data-release-publications> (visited on 11/13/2026).
- [15] M.J. Way et al. “Redshifts in the Southern Abell Redshift Survey Clusters. I. The Data”. In: *aj* 130.5 (2005), pp. 2012–2018. DOI: 10.1086/462418. arXiv: astro-ph/0509736 [astro-ph].
- [16] Martin White. “The mass of a halo”. In: *Astronomy & Astrophysics* 367 (2001), pp. 27–32. DOI: 10.1051/0004-6361:20000357. arXiv: astro-ph/0011495.