


CosmoLattice

School 2022 - Valencia

Non-linear dynamics of
Axion-inflation

Joanes Lizarraga & Ander Uriol



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...


Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

$\phi \rightarrow$ Pseudo-scalar axion field



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$


Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad \tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:


$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad \tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]
...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$


$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

Interaction only through this vertex
Axion coupling!

Λ Coupling constant $[m_{\text{pl}}^{-1}]$

[M. M. Anber, L. Sorbo (0908.4089)]
[J. Cook, L. Sorbo (1109.0022)]
[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]
...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$


$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

Generated from
external mechanism
breaks the shift
symmetry explicitly

Interaction only through this vertex
Axion coupling!

Λ Coupling constant $[m_{\text{pl}}^{-1}]$


- [M. M. Anber, L. Sorbo (0908.4089)]
[J. Cook, L. Sorbo (1109.0022)]
[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$


Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$


Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Vectorial form


$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time

3 vertex



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Our choice of
the potential,
also:

$$V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3}}\phi} \right)^2$$

$$V(\phi) = V_0 \left(1 - \left(\frac{\phi}{\eta} \right)^4 \right)^2$$


$$V(\phi) = \dots$$

3 vertex

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Continuum equations of motion:


Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction
term in the EoMs



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$


CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction
term in the EoMs

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$


No Hubble friction term in the EoMs

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction term in the EoMs


$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$

Non-linear terms



Basics of the model: linear regime

Let's go now deep inside inflation


Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+


dominated by the potential

$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \cancel{\vec{\nabla} \phi} \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \cancel{\vec{\nabla} \phi} \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+

dominated by the potential


$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \cancel{\vec{\nabla} \phi} \times \tilde{\vec{E}}$$

- No backreaction in inflation's dynamics
- No generation of inhomogeneities

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \cancel{\vec{\nabla} \phi} \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+

dominated by the potential

$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

- No backreaction in inflation's dynamics
- No generation of inhomogeneities

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$

Gauge fields get excited by this term



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law


$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

$$\left(\begin{array}{l} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, \quad G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, \quad V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, \quad G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{array} \right)$$



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law


$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

$$\left(\begin{array}{l} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, \quad G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, \quad V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, \quad G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{array} \right)$$



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$


Gauge fields evolve in
the background set by
the axion field

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \left\langle -2K_\phi + V + K_A - G_A \right\rangle$$

$$\begin{pmatrix} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, & G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, & V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, & G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{pmatrix}$$



Basics of the model: linear regime

An analytical solution can be found


Changing to the helicity basis: + Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i \vec{k} \cdot \vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$



Basics of the model: linear regime

An analytical solution can be found

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$


Changing to the helicity basis: + Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k} \cdot \vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis: + Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states


$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$



$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis: + Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$


$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$



$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis: + Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$




$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis:

+

Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$


$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis:

+

Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$


$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability

$$|A^+| \gg |A^-|$$



Basics of the model: linear regime

Analytical solution:


[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by

ξ

Not accurate towards the end
of inflation



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$


Instability controlled by

Deep inside inflation $\xi \rightarrow \text{const}$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$

ξ

Not accurate towards the end
of inflation



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

Topological term:

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by

Deep inside inflation

$$\xi \rightarrow \text{const}$$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$

Appears in EoMs


$$\frac{1}{a^3} \langle \vec{E} \cdot \vec{B} \rangle \simeq 2.4 \cdot 10^{-4} \frac{H^4}{|\xi|^4} e^{2\pi|\xi|}$$

EM energy:

$$\frac{1}{2a^4} \langle a^2 E^2 + B^2 \rangle \simeq 1.4 \cdot 10^{-4} \frac{H^4}{|\xi|^3} e^{2\pi|\xi|}$$

ξ

Not accurate towards the end
of inflation



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

Topological term:

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by ξ

Deep inside inflation

$$\xi \rightarrow \text{const}$$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$

ξ

Not accurate towards the end of inflation


Appears in EoMs

$$\frac{1}{a^3} \langle \vec{E} \cdot \vec{B} \rangle \simeq 2.4 \cdot 10^{-4} \frac{H^4}{|\xi|^4} e^{2\pi|\xi|}$$

EM energy:

$$\frac{1}{2a^4} \langle a^2 E^2 + B^2 \rangle \simeq 1.4 \cdot 10^{-4} \frac{H^4}{|\xi|^3} e^{2\pi|\xi|}$$


Amplified around the Hubble scale at each time




Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds
↓
0 eFolds







Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds
↓
0 eFolds






Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds


0 eFolds




Basics of the model: linear regime

[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

Energy components




Topological term vs slow-roll




Basics of the model: linear regime

[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]


Energy components



Topological term vs slow-roll



Backreaction effect cannot be neglected




Basics of the model: linear regime

Linear regime

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$



Basics of the model: linear regime

Linear regime

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

Basics of the model: linear regime

Linear regime


Dynamical equations:

$$\begin{aligned} \dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}} \end{aligned}$$

Neglected

Expansion equation:

$$\begin{aligned} \dot{\pi}_a &= \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) = \\ &\quad \frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle \end{aligned}$$



Basics of the model

Beyond the linear regime: backreaction


Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Basics of the model

Beyond the linear regime: backreaction


Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) = \frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Backreaction terms



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

- [G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]
[V. Domcke, V. Guidetti, Y. Welling A. Wstphal (2002.02952)]
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\begin{aligned} \tilde{\pi}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}} \end{aligned}$$


*1st approximation
homogeneous*

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

- [G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]
- [V. Domcke, V. Guidetti, Y. Welling A. Wstphal (2002.02952)]
- [E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:


$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

*1st approximation
homogeneous*

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

- [G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]
- [V. Domcke, V. Guidetti, Y. Welling A. Wstphal (2002.02952)]
- [E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

1st approximation
homogeneous

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} \left| A_+(\vec{k}, \tau) \right|^2$$


Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]

[V. Domcke, V. Guidetti, Y. Welling A. Wstphal (2002.02952)]

[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction


Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \tilde{\vec{E}} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$
$$\langle K_A + G_A \rangle = \frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

1st approximation
homogeneous

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \tilde{\vec{E}} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

Due to
backreaction
not only
slow-roll

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$
$$\frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$

Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \vec{E} \cdot \vec{B} \rangle,$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

1st approximation
homogeneous
Due to
backreaction
not only
slow-roll

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

$$\frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$

Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

*1st approximation
homogeneous*

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \vec{E} \cdot \vec{B} \rangle,$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

Due to
backreaction
not only
slow-roll


[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

$$\frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$




Why in the lattice?

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\begin{aligned}\dot{\pi}_a &= \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) = \\ &\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle\end{aligned}$$



Why in the lattice?

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = -a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}$$

No gradient terms induced
(yet)

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields


Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:


$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} - \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:


$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Inhomogeneous source:
Induce gradients not
tractable analytically

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$



$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

\forall D.o.F will be simultaneously
evolved

- Track of all inhomogeneities
- Full backreaction effects included

Discretisation procedure

[D. G. Figueroa & M. Shaposhnikov (1705.09629)]



ϕ @ sites
 A_μ @ links

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

2- Bianchi identities


$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu}\tilde{F}^{\mu\nu} = \partial_\mu K^\mu$$

Exact lattice shift symmetry

4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]


Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

The only one
that satisfies
all conditions



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$


The only one
that satisfies
all conditions


Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$





Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$



Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

2- Bianchi identities


$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu} \tilde{F}^{\mu\nu} = \partial_\mu K^\mu$$

Exact lattice shift symmetry

4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$



Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$


2- Bianchi identities

$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu} \tilde{F}^{\mu\nu} = \text{Exact lattice shift symmetry}$$


4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{\frac{0}{2}}} \left(\Delta_i^+ \phi_{+\hat{\frac{0}{2}}} \right)^2 - \frac{1}{2}a_{+\hat{\frac{0}{2}}}^3 m^2 \phi_{+\hat{\frac{0}{2}}} \right. \\ & + \frac{1}{2}a_{+\hat{\frac{0}{2}}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$



Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a (\vec{\nabla}\phi)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} \left(\frac{1}{2}a_{+\frac{\hat{0}}{2}} \left(\Delta_i^+ \phi_{+\frac{\hat{0}}{2}} \right)^2 - \frac{1}{2}a_{+\frac{\hat{0}}{2}} m^2 \phi_{+\frac{\hat{0}}{2}} \right. \right. \\ & + \frac{1}{2}a_{+\frac{\hat{0}}{2}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \right] \end{aligned}$$

Ready for variational principle




Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L , \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi , \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$



Discretisation procedure

Common kick-drift scheme


Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:


$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+\frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+\frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:


$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+ \frac{\hat{0}}{2}} = \tilde{E}_{i,- \frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+ \frac{\hat{0}}{2}} = \phi_{- \frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+ \hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+ \frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+ \frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+ \frac{\hat{0}}{2}} - a_{+ \frac{\hat{0}}{2}}^3 m^2 \phi_{+ \frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+ \frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{- \frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+ \frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+ \frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+\frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+\frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$


$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

Alternative approach in:
[A. Caravano, E. Komatsu, K. D. Lozanov, J. Weller (2204.12874)]

$$\sum_i \Delta_i^- \tilde{E}_{i,+\frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+\frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$b_{+\hat{0}} = b + dt\mathcal{K}_a^L$$


Expansion equations:

$$a_{+\frac{\hat{0}}{2}} = a_{-\frac{\hat{0}}{2}} + dtb$$

$$\mathcal{K}_a^L = -\frac{a_{+\hat{0}/2}}{6m_{pl}^2}(\rho_L + 3p_L)_{+\hat{0}/2}$$

Hubble's constraint:

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2}\rho_L$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$b_{+\hat{0}} = b + dt \mathcal{K}_a^L$$

Expansion equations:

$$\mathcal{K}_a^L = -\frac{a_{+\hat{0}/2}}{6m_{pl}^2} (\rho_L + 3p_L)_{+\hat{0}/2}$$

$$(\rho_L + 3p_L)_{+\hat{0}/2} = 2(\bar{K}_\phi^L + \bar{K}_{\phi,+\hat{0}}^L) - 2\bar{V}_{\phi,+\hat{0}/2}^L + 2K_A^L + (G_A^L + G_{A,+\hat{0}}^L)$$


$$\rho_L = \bar{K}_\phi^L + \frac{1}{2}(\bar{G}_{\phi,-\hat{0}/2}^L + \bar{G}_{\phi,+\hat{0}/2}^L) + \frac{1}{2}(\bar{V}_{\phi,-\hat{0}/2}^L + \bar{V}_{\phi,+\hat{0}/2}^L) + \frac{1}{2}(\bar{K}_{A,-\hat{0}/2}^L + \bar{K}_{A,+\hat{0}/2}^L) + \bar{G}_A^L$$

Hubble's constraint:

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2} \rho_L$$

$$\bar{K}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2a^6} \tilde{\pi}_\phi^2, \quad \bar{G}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2} \sum_i (\Delta_i^+ \phi_{+\hat{0}/2})^2, \quad \bar{V}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2} m^2 \phi_{+\hat{0}/2}^2$$

$$\bar{K}_A^L = \frac{1}{N^3} \sum_{\vec{n}} \sum_i \frac{1}{2a_{+\hat{0}/2}^4} \tilde{E}_{i,+\hat{0}/2}^2, \quad \bar{G}_A^L = \frac{1}{N^3} \sum_{\vec{n}} \sum_i \frac{1}{4a^4} (\Delta_i^+ A_j - \Delta_j^+ A_i)^2$$




Discretisation procedure

Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \frac{\hat{0}}{2}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$



Discretisation procedure


Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \frac{\hat{0}}{2}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$

Non-symplectic,
Conjugate momenta in kernels!
[Remember 4th lecture & 1st Topical]

$$\dot{\tilde{E}}_i = \mathcal{K}_{i,A}^L \left[a, b, \phi, A_i, \tilde{E}_i \right]$$



Discretisation procedure


Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \frac{\hat{0}}{2}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$

Non-symplectic,
Conjugate momenta in kernels!
[Remember 4th lecture & 1st Topical]

$$\dot{\tilde{E}}_i = \mathcal{K}_{i,A}^L \left[a, b, \phi, A_i, \tilde{E}_i \right]$$




Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L , \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi , \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$



Discretisation procedure

Common kick-drift scheme


Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\mathcal{K}_\phi^L = a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)$$
$$\mathcal{K}_{i,A}^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}$$

Implicit scheme required

We need the value of the gauge's canonical momentum @ 0/2 for obtaining its value @ 0/2!



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a} A_{i,+\hat{0}}$$
$$A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\mathcal{K}_\phi^L = a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi$$
$$\mathcal{K}_{i,A}^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- + \frac{1}{8a} \sum_{\pm j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}$$


Implicit scheme required

We need the value of the gauge's canonical momentum @ 0/2 for obtaining its value @ 0/2!



ANDER URIO


Our Current Work



Explicit-in-time integrator

Joanes

Ander




Explicit-in-time integrator

Joanes

Ander

If momenta appear in kernels



Explicit-in-time integrator


Joanes

If momenta appear in kernels



Ander

Not solvable by explicit
symplectic integrators



Explicit-in-time integrator

Joanes


Ander

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV




Explicit-in-time integrator

Joanes

Ander


If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV


Why?



Explicit-in-time integrator

Joanes

If momenta appear in kernels




Ander

Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV


Why?

Ander

Solution



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Ander

Non-symplectic integrators

Not solvable by explicit
symplectic integrators


i.e. LF or VV

Why?

Solution



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

Ander

Non-symplectic integrators




Solution



Runge-Kutta



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

Ander

Non-symplectic integrators



Solution




Runge-Kutta



(explicit version)

$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$

Solution



Ander


Non-symplectic integrators



(explicit version)

Runge-Kutta


No longer!



Explicit-in-time integrator

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$




Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$






Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{aligned} x(t_{n+1}) &= x(t_n) + dt (b_1 G_1 + b_2 G_2) \end{aligned} \right\}$$




Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$



Explicit-in-time integrator


RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$



Explicit-in-time integrator

RK2 & 2 stages


1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \quad b_2 c_2 = 1/2 , \quad b_2 a_{21} = 1/2$$

Modified Euler



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$


$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$

Modified Euler



$$b_1 = 1/2 , \ b_2 = 1/2 , \ c_2 = 1 , \ a_{21} = 1$$



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$


$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$

Modified Euler




$$b_1 = 1/2 , \ b_2 = 1/2 , \ c_2 = 1 , \ a_{21} = 1$$

$$\boxed{\begin{array}{l} x(t_{n+1}) = x(t_n) + \frac{dt}{2} (G_1 + G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + dt, x(t_n) + G_1 dt) \end{array}}$$




Explicit-in-time integrator



Explicit-in-time integrator

2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$




Explicit-in-time integrator

2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$


1st order ODE RK2 ?



Explicit-in-time integrator


2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$





Coupled 1st order ODEs

1st order ODE RK2 ?





Explicit-in-time integrator







Explicit-in-time integrator






Explicit-in-time integrator





Explicit-in-time integrator


Axion-Inflation using RK2



Explicit-in-time integrator

Axion-Inflation using RK2


$$t_n \equiv (1)$$



Explicit-in-time integrator

Axion-Inflation using RK2


$$t_n \equiv (1) \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n) , \\ A_i^{(1)} = A_i(t_n) , \\ a^{(1)} = a(t_n) , \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n) , \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n) , \\ \pi_a^{(1)} = \pi_a(t_n) , \end{array} \right.$$



Explicit-in-time integrator

Axion-Inflation using RK2


$$t_n \equiv (1) \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n) , \\ A_i^{(1)} = A_i(t_n) , \\ a^{(1)} = a(t_n) , \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n) , \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n) , \\ \pi_a^{(1)} = \pi_a(t_n) , \end{array} \right. \quad \xrightarrow{\text{dotted arrow}} \quad \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right.$$



Explicit-in-time integrator


Axion-Inflation using RK2


$$\begin{aligned} t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\ & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\ & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right. \end{aligned}$$



Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator

Axion-Inflation using RK2






Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}
 t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right.
 \end{aligned}$$

$$\begin{aligned}
 t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.
 \end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}
 t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right.
 \end{aligned}$$


$$\begin{aligned}
 t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.
 \end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2


$$\begin{aligned} t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t , \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t , \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t , \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t , \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t , \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t . \end{array} \right. \\ & \quad \xrightarrow{\text{dotted blue arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\ & \quad \xrightarrow{\text{dotted blue arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right] , \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right] , \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right] . \end{array} \right. \end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$


$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$

$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$

$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$

More intermediate
kernels





Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$






Explicit-in-time integrator


Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$



More intermediate
kernels





Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$







Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$







Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$


$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$





Explicit-in-time integrator

Axion-Inflation using RK2



Explicit-in-time integrator

Axion-Inflation using RK2


Dynamical equations

$$\mathcal{K}_\phi^L = a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)}$$

$$\mathcal{K}_A^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right)$$

$$+ \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right] \right\}$$

$$\mathcal{K}_a^L = -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)$$



Explicit-in-time integrator


Axion-Inflation using RK2

Dynamical equations

$$\begin{aligned}\mathcal{K}_\phi^L &= a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)} \\ \mathcal{K}_A^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\} \\ \mathcal{K}_a^L &= -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)\end{aligned}$$

Constraints

$$\begin{aligned}\sum_i \Delta_i^- \tilde{E}_i &= -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)} \\ \pi_a^2 &= \frac{a^2}{3m_{pl}^2} \rho_L\end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2

No temporal semi-sums

Dynamical equations

$$\mathcal{K}_\phi^L = a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)}$$

$$\mathcal{K}_A^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right)$$


$$+ \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\}$$

$$\mathcal{K}_a^L = -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)$$

Constraints

$$\sum_i \Delta_i^- \tilde{E}_i = -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)}$$

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2} \rho_L$$



Explicit-in-time integrator


Axion-Inflation using RK2

Dynamical equations

$$\begin{aligned}\mathcal{K}_\phi^L &= a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)} \\ \mathcal{K}_A^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\} \\ \mathcal{K}_a^L &= -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)\end{aligned}$$

Constraints


$$\begin{aligned}\sum_i \Delta_i^- \tilde{E}_i &= -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)} \\ \pi_a^2 &= \frac{a^2}{3m_{pl}^2} \rho_L\end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D .G. Figueroa (1812.03132)]







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D .G. Figueroa (1812.03132)]






Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme






Explicit-in-time integrator


Axion-Inflation using RK2: validation of the scheme

Spectrum of A

$$\frac{k^3}{2\pi^2 m^2} |A_k|^2$$

Preheating example





Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D .G. Figueroa (1812.03132)]

Spectrum of A

$$\frac{k^3}{2\pi^2 m^2} |A_k|^2$$

Preheating example


Comoving momentum


$$k/m$$

Solid: Canivete and Figueroa
Scatter: Our scheme

LF implicit

RK2 explicit







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]






Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]




Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]






Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


Gauss constraint
observable





Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time




Initial Conditions in the Linear Regime

$$\dot{\tilde{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$


Helicity basis



Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis




Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



$$|A^+| \gg |A^-|$$



Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time


$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



$$|A^+| \gg |A^-|$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$


$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Deep inside Hubble $k \gg aH$



$$|A^+| \gg |A^-|$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

Deep inside Hubble $k \gg aH$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0 \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$

$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$

$|A^+| \gg |A^-|$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution


Deep inside Hubble $k \gg aH$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$

$$|A^+| \gg |A^-|$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$


Bunch-Davies solution

Deep inside Hubble $k \gg aH$


$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$




$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



$|A^+| \gg |A^-|$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution


Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



$|A^+| \gg |A^-|$

Cosmic time

$\tau \simeq -1/aH$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$

Cosmic time

$\tau \simeq -1/aH$

$$A_+(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time

$\tau \simeq -1/aH$

$$A_+(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$

Time derivative





Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time

$\tau \simeq -1/aH$

$$A_+(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$

Time derivative

$$E_+(k, t) \simeq \frac{i}{a} \sqrt{\frac{k}{2}} e^{ik/a(t)H(t)}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time

$\tau \simeq -1/aH$

$$A_\pm(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$

Time derivative


$$E_\pm(k, t) \simeq \frac{i}{a} \sqrt{\frac{k}{2}} e^{ik/a(t)H(t)}$$




Initial Conditions in the Linear Regime


Helicity basis

BD solutions for A^\pm, E^\pm





Initial Conditions in the Linear Regime







Initial Conditions in the Linear Regime






Initial Conditions in the Linear Regime






Initial Conditions in the Linear Regime




$$k_i \varepsilon_i^\pm(\hat{k}) = 0 , \quad \epsilon_{ijk} k_j \varepsilon_i^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k}) ,$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}) , \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(-\hat{k}) = \delta_{\lambda\lambda'}$$




Initial Conditions in the Linear Regime



$$k_i \varepsilon_i^\pm(\hat{k}) = 0 , \quad \epsilon_{ijk} k_j \varepsilon_i^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k}) ,$$


$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}) , \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(-\hat{k}) = \delta_{\lambda\lambda'}$$

$$A_i^\pm = P_{ij}^\pm A_j$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?




Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?



Initial Conditions in the Linear Regime


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

2 eigenvalues: 1 and -1

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

Transverse projector


$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

How to build the
Helicity Projector?



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?




Helicity projector


$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?




Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector


$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



$$A_i^{\pm} = P_{ij}^{\pm} A_j$$
$$E_i^{\pm} = P_{ij}^{\pm} E_j$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$




Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$




$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



$$A_i^{\pm} = P_{ij}^{\pm} A_j$$
$$E_i^{\pm} = P_{ij}^{\pm} E_j$$

Gauss


$$\vec{\nabla} \cdot \vec{E} = 0$$



Initial Conditions in the Linear Regime

Two IC options






Initial Conditions in the Linear Regime

Two IC options

One option

Only project the + helicity states






Initial Conditions in the Linear Regime

Two IC options

One option

Only project the + helicity states

Why?



Initial Conditions in the Linear Regime


Two IC options

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced



Initial Conditions in the Linear Regime

Two IC options


One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced

Loss of information!



Initial Conditions in the Linear Regime

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced




Two IC options

Our option

Project both states + and -





Initial Conditions in the Linear Regime


Two IC options

One option

Only project the + helicity states

Why?


A^+ is exponentially enhanced



Our option

Project both states + and -

Why?



Initial Conditions in the Linear Regime

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced




Two IC options

Our option

Project both states + and -

Why?

Main objective: non-linear regime



Initial Conditions in the Linear Regime


One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced

Loss of information!



Two IC options

Our option


Project both states + and -

Why?

Main objective: non-linear regime




Can A^- become relevant?



Initial Conditions in the Linear Regime

Gauge initialization process




Initial Conditions in the Linear Regime

Gauge initialization process

1

Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$




Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF



Initial Conditions in the Linear Regime

Gauge initialization process


1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

2 Add fluctuations to A_i with RGF



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables


$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

- 2 Add fluctuations to A_i with RGF

- 4 Repeat the process with A_-



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$


- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

- 4 Repeat the process with A_-

Fourier



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF


- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$


Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT



EOMs evolved in real space!



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$


Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT

- 6 Repeat initialization with E^\pm BD solutions

EOMs evolved in real space!



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

Fourier

- 4 Repeat the process with A_-


- 5 Go back to real space via FFT

- 6 Repeat initialization with E^\pm BD solutions


EOMs evolved in real space!


⋮ We will have an amplitude of 2 BD in A_i ⋮

$$\sum_i |A_i|^2 = |A_+|^2 + |A_-|^2$$





Simulations in the Linear Regime







Simulations in the Linear Regime







Simulations in the Linear Regime







Simulations in the Linear Regime







Simulations in the Linear Regime






Simulations in the Linear Regime







Simulations in the Linear Regime




Simulations in the Linear Regime



Simulations in the Linear Regime






Non-linear evolution: example


Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

@





Non-linear evolution: example

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$


Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

@







Non-linear evolution: example

Gauge PS

A^+

Linear







Non-linear evolution: example

Gauge PS

A^+

Linear





Non-linear evolution: example


Gauge PS


A^+

Linear



Non-linear







Non-linear evolution: example


Gauge PS


A^+

Linear



Non-linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear

Non-linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear


Non-linear






Non-linear evolution: example


What's
happening in
those extra
folds?






Non-linear evolution: example

What's
happening in
those extra
folds?







Non-linear evolution: example

What's
happening in
those extra
folds?

A_{tot}

A^+






Non-linear evolution: example

What's
happening in
those extra
folds?

A_{tot}

A^+

A^-




- 6 eFolds
- 5 eFolds
- 4 eFolds
- 3 eFolds
- 2 eFolds
- 1 eFolds
- 0 eFolds
- 1 eFolds
- 2 eFolds
- 3 eFolds
- 4 eFolds


Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear






Summary & Conclusions

- Successful implementation of axion-inflation in CL


$$\frac{\phi}{\Lambda} F \tilde{F}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simpletic integrators needed: RK
 - Validation: preheating

$$\frac{\phi}{\Lambda} F \tilde{F}$$




Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating




$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime


$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime


$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions


- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime
- Preliminari non-linear sims already show inhomogeneities non-negligible

$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simpletic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime
- Preliminär non-linear sims already show inhomogeneities non-negligible



Ready to study non-linear phenomenology!

$$\frac{\phi}{\Lambda} F \tilde{F}$$

Thank you!