shear_orientation: ionic census and conditional T/n versus elapsed_time
Inherited atomic ions; T/n all-bare instant-thermalized passive totals, no T charge projection
Auxiliary axis uses reference clock: quotient_L16_S0_Am0_Ar0; other cases have different z(t).

n: nuclear total only

H1: saved ionic census Y_q

D: saved ionic census Y_q

T: nuclear total only

He3: saved ionic census Y_q

He4: saved ionic census Y_q

Li6: saved ionic census Y_q

Li7: saved ionic census Y_q

Reference redshift z

Proper elapsed time since z=20 (Myr)

Y = n_species/n_B (log scale)

Y_Aq = n_Aq/n_B (log scale)

Initial n=0; old n₀ hard-zero

Initial T=0; T charge states unresolved; old T₀/T₁ hard-zero

Saved-model exact zero: charge 2, 3

Σ₀=≈0, Pₘ₀=≈0, P_r0=≈0
Σ₀=+6.00e-01, Pₘ₀=≈0, P_r0=≈0
Σ₀=-6.00e-01, Pₘ₀=≈0, P_r0=≈0
Σ₀=+6.00e-01, Pₘ₀=+1.03e-03, P_r0=-2.25e-03
Σ₀=-6.00e-01, Pₘ₀=+1.03e-03, P_r0=-2.25e-03
Σ₀=+6.00e-01, Pₘ₀=+1.03e-03, P_r0=+4.49e-03
ionic charge 0
ionic charge 1
ionic charge 2
ionic charge 3