sigma: ionic census and conditional T/n versus redshift
Inherited atomic ions; T/n all-bare instant-thermalized passive totals, no T charge projection
Auxiliary axis uses reference clock: quotient_L16_S0_Am0_Ar0; other cases have different z(t).

n: nuclear total only

H1: saved ionic census Y_q

D: saved ionic census Y_q

T: nuclear total only

He3: saved ionic census Y_q

He4: saved ionic census Y_q

Li6: saved ionic census Y_q

Li7: saved ionic census Y_q

Reference elapsed time (Myr)

Redshift z

Y = n_species/n_B (log scale)

Y_Aq = n_Aq/n_B (log scale)

Initial n=0; old n_0 hard-zero

Initial T=0; T charge states unresolved; old T_0/T_1 hard-zero

Saved-model exact zero: charge 2, 3

Σ_0=≈0, P_{m0}=≈0, P_{r0}=≈0
Σ_0=+1.00e-02, P_{m0}=≈0, P_{r0}=≈0
Σ_0=+5.00e-02, P_{m0}=≈0, P_{r0}=≈0
Σ_0=+2.00e-01, P_{m0}=≈0, P_{r0}=≈0
Σ_0=+6.00e-01, P_{m0}=≈0, P_{r0}=≈0
ionic charge 0
ionic charge 1
ionic charge 2
ionic charge 3