

ALTERNATIVE WEARABLE FORM FACTORS AND DISTRIBUTED NODE VARIANTS

GENERALIZED WEARABLE IMPLEMENTATION

Although Node 5 has been described with reference to ring-based and chest-mounted embodiments, the contextual physiology monitoring architecture is not limited to any single wearable form factor. The disclosed system encompasses any body-coupled device capable of acquiring systemic physiological signals used to interpret intimate physiology.

BODY-PLACEMENT VARIABILITY

Node 5 may be positioned at various anatomical locations including wrist, forearm, upper arm, neck, clavicle region, torso, waist, hip, thigh, ankle, or other body regions capable of providing meaningful physiological reference signals. Selection of placement may depend on signal quality, user comfort, intended monitoring duration, or integration with additional wearable systems.

TEXTILE AND GARMENT INTEGRATION

In certain embodiments, Node 5 sensors may be integrated directly into garments or wearable textiles including shirts, undergarments, sleepwear, athletic clothing, compression garments, or specialized biosensing fabrics. Textile-integrated sensing systems may distribute multiple sensing regions across the body to collectively form a unified contextual physiology node.

DISTRIBUTED SENSOR NETWORKS

Node 5 may comprise multiple physically separated sensing modules operating cooperatively as a single logical contextual physiology device. Distributed sensors may communicate wirelessly or through conductive textile pathways to aggregate physiological signals into a unified context model.

MODULAR AND DETACHABLE SYSTEMS

Embodiments may include modular sensing units capable of attachment, removal, replacement, or reconfiguration depending on monitoring objectives. Modules may be interchangeable across different wearable platforms while maintaining functional equivalence within the contextual physiology architecture.

FUTURE FORM FACTOR COVERAGE

The scope of the disclosed invention expressly includes future wearable technologies, novel body-interface mechanisms, implant-adjacent systems, augmented-reality-integrated wearables, or other emerging human-device interaction platforms capable of performing substantially similar contextual physiology monitoring functions.