RETRACTED AS PROOF

ARCHIVE HISTORIQUE - T1'-T4 argument invalidated

Editorial notice - 20 October 2026

The historical T1'-T4 chain does not prove the Riemann hypothesis. The public erratum of 13 October 2026 identifies errors in T2 (weighted Gram-matrix bound) and T3 (wave-packet normalization and capture argument).

Earlier titles, abstracts and claims of complete or journal-ready proofs are historical claims withdrawn as proof arguments. This withdrawal does not assert that every mathematical object cited is false.

This editorial page has been added to the historical PDF. All original pages follow with their content unchanged. This is neither a recompilation nor a new mathematical result.

Source document: docs/guide_interdisciplinaire.pdf

Original PDF SHA-256:
`976f0af5031ca523408cad2a72e578e2f5103bee48481d04a6be99e8e0fa588a`

Public T1'-T4 erratum - 13 October 2026

Interdisciplinary Guide — T1′–T4 & InterIA

Alexandre Couret
InterIA Mathematical Collective

February 27, 2026

Abstract

This 2–3 page guide explains the T1′–T4 plan, translation between disciplines, and how to read the proof without switching hats. InterIA artifacts are separate.

1. Reading routes (personas)

1 Reading routes (by specialty)

NT. T1′ → T2 → T3; T4 minimal.
AO. T2 → T1′ → T3; T4 as spectral reading.
Signal. Tapers, bandlimit; T2 as frame conditioning; T3 as matched-filter probe.
Phys. Rigidity (T2) + localized measurement (T3); HP compatible.
Repro. See InterIA (separate).
Formal. Lemmas: Schur/Gershgorin, PW density, Archimedean, absorption.

2. Cross-dictionary

2 Notations and preliminaries

NT	Cross reading
Band $[-A, A]$	Bandwidth (signal)
Cos^2 taper	Apodization (plateau/transition)
Ξ_A	Non-uniform sampling
$\log p / p^{k/2}$	p–adic energy weighting
$C_{\mathrm{fr}}^{(w)} \geq 1 - \Theta_w$	Frame conditioning (AO)
Wave–packet $\hat{\phi}_T$	Matched filter (signal/phys)
Explicit formula	Source–spectrum coupling

3. Storyboard

3 Storyboard T1′–T4

T1′: set the arena (PW, explicit, Archimedean).
T2: gap via locality + weights.
T3: packet → contradiction if off-line zero.
T4: HP-compatible reading (no postulate).

4. Bridges

4 Bridges boxes

(AO) $G^{(w)}$ Gram; $I + R$ with $\|R\| < 1$ gives lower boundedness.
(Signal) Band $[-A, A]$, apodized taper, frame $\approx$ robust reconstruction.
(Phys) T2 = spectral rigidity; T3 = localized measurement.
(Repro) Interval certificates; proof does not depend on numerics.

5. Micro-example (T2)

5 Micro-example (T2)

$A = 4$; $p \le e^4$.
Few neighbors (locality), decay $p^{-(k-k_\alpha)/4}$.
Indicative: $\Theta_w(A) \approx 0.63 \Rightarrow C_{\mathrm{fr}}^{(w)} \approx 0.37 > 0$.

6. FAQ

6 FAQ (by specialty)

NT: why $p^{-k/2}$? — damp prime powers.
AO: why Schur? — direct control of $\|R\|$.
Signal: why $\cos^2$? — bounded slope, simple calculus.
Phys: HP? — reading only.
Repro: numbers? — certificates, not part of the proof.




Figure 1: Dependency map: T1′→T2→T3; T4 is HP-compatible reading.