


9-trees: 树 (Trees)


魏恒峰


hfwei@hnu.edu.cn

2026 年 05 月 12 日











- (1) How many **trees** are there with n labeled vertices?
- (2) Draw all homomorphically **irreducible** trees with $n = 10$.



- (1) How many **trees** are there with n labeled vertices? n^{n-2}
- (2) Draw all homomorphically **irreducible** trees with $n = 10$.



(1) How many **trees** are there with n labeled vertices? n^{n-2}

(2) Draw all homomorphically **irreducible** trees with $n = 10$. 10

Definition (Tree (树))



A **tree** is a **connected acyclic undirected** graph.




Definition (Tree (树))


A **tree** is a **connected acyclic undirected** graph.

Definition (Forest (森林))

A **forest** is a **acyclic undirected** graph.







Cayley Graph (4-regular tree)

Definition (Internal Vertex (内部顶点); Leaf (叶子))

In a tree T with ≥ 2 vertices, for a vertex v in T , if

$$\deg(v) = 1$$


then v is called a **leaf**; otherwise, v is an **internal vertex**.

Definition (Internal Vertex (内部顶点); Leaf (叶子))

In a tree T with ≥ 2 vertices, for a vertex v in T , if

$$\deg(v) = 1$$

then v is called a **leaf**; otherwise, v is an **internal vertex**.




Definition (Internal Vertex (内部顶点); Leaf (叶子))

In a tree T with ≥ 2 vertices, for a vertex v in T , if

$$\deg(v) = 1$$

then v is called a **leaf**; otherwise, v is an **internal vertex**.



Lemma


Any tree T with ≥ 2 vertices contains ≥ 1 leaf.

Definition (Internal Vertex (内部顶点); Leaf (叶子))

In a tree T with ≥ 2 vertices, for a vertex v in T , if

$$\deg(v) = 1$$

then v is called a **leaf**; otherwise, v is an **internal vertex**.



Lemma

Any tree T with ≥ 2 vertices contains ≥ 1 leaf.

Otherwise, $\forall v \in V \deg(v) \geq 2 \implies T$ has cycles.

Lemma

Any tree T with ≥ 2 vertices contains ≥ 2 leaves.

Lemma

Any tree T with ≥ 2 vertices contains ≥ 2 leaves.

Consider the **two endpoints** of any **maximal** path in T .

Lemma

Any tree T with ≥ 2 vertices contains ≥ 2 leaves.

Consider the **two endpoints** of any **maximal** path in T .

They are leaves of T .

Lemma

*Deleting a **leaf** from a tree T with n vertices produces a **tree** with $n - 1$ vertices.*

Lemma

Deleting a *leaf* from a tree T with n vertices produces a *tree* with $n - 1$ vertices.



$G' = G - v$ is **connected** and **acyclic**.

Lemma

Deleting a *leaf* from a tree T with n vertices produces a *tree* with $n - 1$ vertices.



$G' = G - v$ is **connected** and **acyclic**.

A leaf does *not* belong to any paths connecting two other vertices.

Lemma

Deleting a *leaf* from a tree T with n vertices produces a *tree* with $n - 1$ vertices.



$G' = G - v$ is **connected** and **acyclic**.

A leaf does *not* belong to any paths connecting two other vertices.

This lemma can be used in induction for trees!

Theorem ((We call it) Characterization of Trees)

Let T be an undirected graph with n vertices.

Then the following statements are *equivalent*:

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges;
- (3) T is connected, and has $m = n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path;
- (6) T is acyclic, but the addition of any edge creates exactly one cycle.

Theorem ((We call it) Characterization of Trees)

Let T be an undirected graph with n vertices.

Then the following statements are *equivalent*:

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges;
- (3) T is connected, and has $m = n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path;
- (6) T is acyclic, but the addition of any edge creates exactly one cycle.

$$(1) \implies (2) \implies (3) \implies (4) \implies (5) \implies (6) \implies (1)$$

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Induction Hypothesis: Any trees with $n - 1$ vertices has $n - 2$ edges.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Induction Hypothesis: Any trees with $n - 1$ vertices has $n - 2$ edges.

Induction Step: Consider a tree T with $n \geq 2$ vertices.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Induction Hypothesis: Any trees with $n - 1$ vertices has $n - 2$ edges.

Induction Step: Consider a tree T with $n \geq 2$ vertices.

T has a leaf v .

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Induction Hypothesis: Any trees with $n - 1$ vertices has $n - 2$ edges.

Induction Step: Consider a tree T with $n \geq 2$ vertices.

T has a leaf v .

For $T' = T - v$, $m(T') = (n - 1) - 1 = n - 2$.

Theorem (Characterization of Trees)

- (1) T is a tree;
- (2) T is acyclic, and has $m = n - 1$ edges.

By induction on the number n of vertices of trees.

Basis Step: $n = 1$. $m = 0 = n - 1$.

Induction Hypothesis: Any trees with $n - 1$ vertices has $n - 2$ edges.

Induction Step: Consider a tree T with $n \geq 2$ vertices.

T has a leaf v .

For $T' = T - v$, $m(T') = (n - 1) - 1 = n - 2$.

$$m(T) = (n - 2) + 1 = n - 1.$$

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

By Contradiction.

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

By Contradiction.

Suppose that T is *disconnected*.

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

By Contradiction.

Suppose that T is *disconnected*.

T is a forest, consisting of $k \geq 2$ trees $T_1, T_2, \dots$.

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

By Contradiction.

Suppose that T is *disconnected*.

T is a forest, consisting of $k \geq 2$ trees $T_1, T_2, \dots$.

By (2), for each T_i , $m(T_i) = n(T_i) - 1$.

Theorem (Characterization of Trees)

- (2) T is acyclic, and has $n - 1$ edges;
- (3) T is connected, and has $n - 1$ edges.

By Contradiction.

Suppose that T is *disconnected*.

T is a forest, consisting of $k \geq 2$ trees $T_1, T_2, \dots$.

By (2), for each T_i , $m(T_i) = n(T_i) - 1$.


$$m(T) = \sum_{i=1}^k m(T_i) = n - k \neq n - 1.$$

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.



Definition (Bridge (桥))

A **bridge** of a graph G is an **edge** e such that

$$c(G - e) > c(G).$$

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.

Consider any edge e of T .

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.

Consider any edge e of T .

$$m(T - e) = (n - 1) - 1 = n - 2.$$

Theorem (Characterization of Trees)

- (3) T is connected, and has $n - 1$ edges;
- (4) T is connected, and each edge is a *bridge*.

Consider any edge e of T .

$$m(T - e) = (n - 1) - 1 = n - 2.$$

$T - e$ must be disconnected.

Theorem (Characterization of Trees)

- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path.

Theorem (Characterization of Trees)

- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path.

Consider any two vertices u and v .

Theorem (Characterization of Trees)

- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path.

Consider any two vertices u and v .

T is connected $\implies u$ and v are connected by ≥ 1 path.

Theorem (Characterization of Trees)

- (4) T is connected, and each edge is a *bridge*;
- (5) Any two vertices of T are connected by exactly one path.

Consider any two vertices u and v .

T is connected $\implies u$ and v are connected by ≥ 1 path.

If u and v are connected by two paths,
the edges on these two paths are not bridges.

Theorem (Characterization of Trees)

- (5) *Any two vertices of T are connected by exactly one path;*
- (6) *T is acyclic, but the addition of any edge creates exactly one cycle.*

Theorem (Characterization of Trees)

- (5) *Any two vertices of T are connected by exactly one path;*
- (6) *T is acyclic, but the addition of any edge creates exactly one cycle.*

If T has a cycle C ,
any two vertices in C is connected by ≥ 2 paths.

Theorem (Characterization of Trees)

- (5) *Any two vertices of T are connected by exactly one path;*
- (6) *T is acyclic, but the addition of any edge creates exactly one cycle.*

If T has a cycle C ,

any two vertices in C is connected by ≥ 2 paths.

Consider the addition of edge $\{u, v\}$ to T .

It creates a cycle, consisting of $\{u, v\}$ and the path from u to v .

Theorem (Characterization of Trees)

- (5) *Any two vertices of T are connected by exactly one path;*
- (6) *T is acyclic, but the addition of any edge creates exactly one cycle.*

If T has a cycle C ,

any two vertices in C is connected by ≥ 2 paths.

Consider the addition of edge $\{u, v\}$ to T .

It creates a cycle, consisting of $\{u, v\}$ and the path from u to v .

Lemma

If two distinct cycles of a graph G share a common edge e , then G has a cycle that does not contain e .

Theorem (Characterization of Trees)

- (6) T is acyclic, but the addition of any edge creates exactly one cycle;
- (1) T is a tree.

Theorem (Characterization of Trees)

- (6) T is acyclic, but the addition of any edge creates exactly one cycle;
- (1) T is a tree.

Suppose that T is disconnected.

Theorem (Characterization of Trees)

- (6) T is acyclic, but the addition of any edge creates exactly one cycle;
- (1) T is a tree.

Suppose that T is disconnected.

T is a forest, consisting of ≥ 2 trees $T_1, T_2, \dots$

Theorem (Characterization of Trees)


- (6) T is acyclic, but the addition of any edge creates exactly one cycle;
- (1) T is a tree.

Suppose that T is disconnected.

T is a forest, consisting of ≥ 2 trees $T_1, T_2, \dots$

Choose $u \in V(T_1), v \in V(T_2)$.

$T + \{u, v\}$ does **not** create cycles.




Spanning Trees (trees in graphs)

Definition (Subgraph (子图))

A graph S is a **subgraph** of G if


$$V(S) \subseteq V(G) \wedge E(S) \subseteq E(G) \wedge \bigcup E(S) \subseteq V(S)$$



Definition (Subgraph (子图))

A graph S is a **subgraph** of G if

$$V(S) \subseteq V(G) \wedge E(S) \subseteq E(G) \wedge \bigcup E(S) \subseteq V(S)$$



Definition (Induced Subgraph (诱导子图))

A graph S is an **induced subgraph** of G if S is a **subgraph** of G such that


$$\{\{u, v\} \in E(G) \mid u \in V(S), v \in V(S)\} \subseteq E(S).$$

Definition (Spanning Tree (生成树))

A **spanning tree** T of an **undirected** graph G is a **subgraph** that is a **tree** with all vertices of G .



Definition (Spanning Tree (生成树))

A **spanning tree** T of an **undirected** graph G is a **subgraph** that is a **tree** with all vertices of G .



Definition (Spanning Tree (生成树))

A **spanning tree** T of an **undirected** graph G is a **subgraph** that is a **tree** with all vertices of G .



Theorem

Every connected undirected graph G admits a spanning tree.

Theorem

Every connected undirected graph G admits a spanning tree.

Repeatedly deleting edges in cycles until the graph is acyclic.

Theorem

Every connected undirected graph G admits a spanning tree.

Repeatedly deleting edges in cycles until the graph is acyclic.


The remaining graph is a spanning tree of G .

Definition (Minimum Spanning Tree (MST; 最小生成树))

A **minimum spanning tree** T of an **edge-weighted** undirected graph G is a spanning tree with **minimum** total weight of edges.


Definition (Minimum Spanning Tree (MST; 最小生成树))

A **minimum spanning tree** T of an **edge-weighted** undirected graph G is a spanning tree with **minimum** total weight of edges.



Definition (Minimum Spanning Tree (MST; 最小生成树))

A **minimum spanning tree** T of an **edge-weighted** undirected graph G is a spanning tree with **minimum** total weight of edges.



Existence?

Uniqueness?




Algorithms?


Theorem

Every weighted connected undirected graph G admits a minimum spanning tree.

Theorem

Every weighted connected undirected graph G *admits* a minimum spanning tree.






Joseph Kruskal (1928 ~ 2010)

Repeatedly adding the next lowest-weight edge that will not form a cycle until $n - 1$ edges are added.

Repeatedly adding the next lowest-weight edge that will not form a cycle until $n - 1$ edges are added.






Robert C. Prim (1921 ~ 2021)

Repeatedly adding the **cheapest** possible edge
from **the partially built tree** to another vertex,
until $n - 1$ edges are added.

Repeatedly adding the **cheapest** possible edge from **the partially built tree** to another vertex, until $n - 1$ edges are added.



Theorem (Cut Property)


X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$



e : A *lightest* edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of *some* MST T' of G .

(a)





(b)

Edges X :MST T :

(c)

The cut:

MST T' :

Theorem (Cut Property)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e : A lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of *some* MST T' of G .

Theorem (Cut Property)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$


e : A lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of *some* MST T' of G .


Correctness of Kruskal's and Prim's algorithms.

By Exchange Argument.

By Exchange Argument.




By Exchange Argument.




If $e \in T$, we are done.

By Exchange Argument.



If $e \in T$, we are done. Otherwise, construct T' .


By Exchange Argument.



If $e \in T$, we are done. Otherwise, construct T' .

$$T' = \underbrace{\underbrace{T}_{X \subseteq T} + \{e\} - \{e'\}}_{\text{if } e \notin T}$$

By Exchange Argument.



If $e \in T$, we are done. Otherwise, construct T' .

$$T' = \underbrace{T + \{e\} - \{e'\}}_{\substack{\text{if } e \notin T \\ X \subseteq T}}$$

$w(e) \leq w(e') \implies T'$ is also an MST

Theorem (Cut Property)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e : *A* lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of *some* MST T' of G .

Theorem (Cut Property)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e : *A* lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of *some* MST T' of G .

“a” $\rightarrow$ “the” $\implies$ “some” $\rightarrow$ “all”

Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be *a* maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Choose any MST T of G .

Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Choose any MST T of G .

If $e \notin T$, we are done. Otherwise, construct T' .


Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Choose any MST T of G .

If $e \notin T$, we are done. Otherwise, construct T' .




Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be a maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Choose any MST T of G .

If $e \notin T$, we are done. Otherwise, construct T' .



$$T' = T - \underbrace{\{e\}}_{\text{if } e \in T} + \{e'\}$$

Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be *a* maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

Theorem (Cycle Property)

- ▶ Let C be any cycle in connected undirected G
- ▶ Let $e = (u, v)$ be *a* maximum-weight edge in C

Then $\exists$ MST T of $G : e \notin T$.

“a” $\rightarrow$ “the” $\Rightarrow$ “ $\exists$ ” $\rightarrow$ “ $\forall$ ”

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

*If each edge has a **distinct** weight, then there is a **unique** MST of G .*

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

*If each edge has a **distinct** weight, then there is a **unique** MST of G .*

By Contradiction.

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

If each edge has a *distinct* weight, then there is a *unique* MST of G .

By Contradiction.

$$\exists \text{ MSTs } T_1 \neq T_2$$

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

If each edge has a *distinct* weight, then there is a *unique* MST of G .

By Contradiction.

$$\exists \text{ MSTs } T_1 \neq T_2$$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

If each edge has a *distinct* weight, then there is a *unique* MST of G .

By Contradiction.

$$\exists \text{ MSTs } T_1 \neq T_2$$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

$$e = \min \Delta E$$

Theorem (Uniqueness of MST)

Let G be an edge-weighted undirected graph.

If each edge has a *distinct* weight, then there is a *unique* MST of G .


By Contradiction.


$$\exists \text{ MSTs } T_1 \neq T_2$$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$


$$e = \min \Delta E$$

Suppose that $e \in T_1 \setminus T_2$ (w.l.o.g)






$$T_2 + \{e\} \Rightarrow C$$




$$T_2 + \{e\} \Rightarrow C$$

$$\exists (e' \in C) \notin T_1$$



$$T_2 + \{e\} \implies C$$

$$\exists(e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E$$



$$T_2 + \{e\} \Rightarrow C$$


$$\exists(e' \in C) \notin T_1 \Rightarrow e' \in T_2 \setminus T_1 \Rightarrow e' \in \Delta E \Rightarrow w(e') > w(e)$$


Condition for Uniqueness of MST

Unique MST $\not\Rightarrow$ Distinct weights

Condition for Uniqueness of MST

Unique MST $\not\Rightarrow$ Distinct weights







Theorem (Cayley's Formula)

The number T_n of *labeled* trees on $n \geq 2$ vertices is n^{n-2} .


Theorem (Cayley's Formula)

The number T_n of *labeled* trees on $n \geq 2$ vertices is n^{n-2} .





Arthur Cayley (1821 ~ 1895)



Chapter 33: Cayley's formula for the number of trees

By Double Counting.

— Jim Pitman

By Double Counting.

— Jim Pitman

[https://en.wikipedia.org/wiki/Double_counting_\(proof_technique\)#Counting_trees](https://en.wikipedia.org/wiki/Double_counting_(proof_technique)#Counting_trees)

By Double Counting.

— Jim Pitman

[https://en.wikipedia.org/wiki/Double_counting_\(proof_technique\)#Counting_trees](https://en.wikipedia.org/wiki/Double_counting_(proof_technique)#Counting_trees)


How many ways are there of forming a **directed rooted tree** from an **empty graph** by adding **directed edges** one by one?

Definition (Rooted Tree (有根树))

A **rooted tree** is a **tree** where one vertex has been **designated the root**.


Definition (Rooted Tree (有根树))

A **rooted tree** is a **tree** where one vertex has been **designated the root**.



Definition (Rooted Tree (有根树))

A **rooted tree** is a **tree** where one vertex has been **designated the root**.



Definition (Directed Rooted Tree (有向有根树))

A **directed rooted tree** is a **rooted tree** where all edges directed **away from** or **towards** the root.

Choose one of the T_n labeled trees on n vertices.

Choose one of the T_n labeled trees on n vertices.

Choose one of its n vertices as root.

Choose one of the T_n labeled trees on n vertices.

Choose one of its n vertices as root.

Choose one of the $(n-1)!$ possible sequences
in which to add its $n-1$ directed edges.

$$T_n n(n-1)! = T_n n!$$

Suppose that we have added $n - k$ directed edges.

Suppose that we have added $n - k$ directed edges.

We obtain a rooted forest with k trees.

Suppose that we have added $n - k$ directed edges.


We obtain a rooted forest with k trees.

There are $n(k - 1)$ choices for the next edge to add.

Suppose that we have added $n - k$ directed edges.

We obtain a rooted forest with k trees.


There are $n(k - 1)$ choices for the next edge to add.



Suppose that we have added $n - k$ directed edges.

We obtain a rooted forest with k trees.

There are $n(k - 1)$ choices for the next edge to add.



$$\prod_{k=2}^n n(k-1) = n^{n-1}(n-1)! = n^{n-2}n!$$

$$T_n n! = n^{n-2} n!$$

$$T_n n! = n^{n-2} n!$$

$$T_n = n^{n-2}$$

$$T_n n! = n^{n-2} n!$$

$$T_n = n^{n-2}$$



Definition (Irreducible Tree)


An **irreducible tree** is a tree T where

$$\forall v \in V(T) \deg(v) \neq 2.$$

Definition (Irreducible Tree)

An **irreducible tree** is a tree T where



$$\forall v \in V(T) \deg(v) \neq 2.$$



Definition (Irreducible Tree)

An **irreducible tree** is a tree T where



$$\forall v \in V(T) \deg(v) \neq 2.$$



Definition (Irreducible Tree)


An **irreducible tree** is a tree T where

$$\forall v \in V(T) \deg(v) \neq 2.$$



Homeomorphically Irreducible Trees of size $n = 10$

Thank
You!



hfwei@hnu.edu.cn