

14-subgroup: 子群 (Subgroup)


魏恒峰

hfwei@hnu.edu.cn

2026 年 06 月 16 日




Lagrange's Theorem



Joseph-Louis Lagrange (1736 ~ 1813)

Lagrange's Theorem



Help us understand the structure of a group via its subgroups

Definition (Coset (陪集))

Suppose that $H \leq G$. For $a \in G$,

$$aH = \{ah \mid h \in H\}, \quad Ha = \{ha \mid h \in H\},$$

is called the **left coset** (左陪集) and **right coset** of H in G , respectively.

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$H = \{(1), (1\ 2)\} \leq S_3$$

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$H = \{(1), (1\ 2)\} \leq S_3$$

$$(1)H = H = (1\ 2)H \leq S_3$$

$$(1\ 3)H = \{(1\ 3), (1\ 2\ 3)\} = (1\ 2\ 3)H$$

$$(2\ 3)H = \{(2\ 3), (1\ 3\ 2)\} = (1\ 3\ 2)H$$

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$A_3 = \{(1), (1\ 2\ 3), (1\ 3\ 2)\} \leq S_3$$

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$A_3 = \{(1), (1\ 2\ 3), (1\ 3\ 2)\} \leq S_3$$

$$(1)A_3 = (1\ 2\ 3)A_3 = (1\ 3\ 2)A_3 = A_3 \leq S_3$$

$$(1\ 2)A_3 = (1\ 3)A_3 = (2\ 3)A_3 = \{(1\ 2), (1\ 3), (2\ 3)\}$$

Theorem

Suppose that $H \leq G$, $a, b \in G$.

(1)

$$a \in aH$$

(2)

$$aH = H \iff a \in H$$

(3)

$$aH \leq G \iff a \in H$$

(4)

$$aH = bH \iff a^{-1}b \in H$$

(5)

$$\forall a, b \in G (aH = bH) \vee (aH \cap bH = \emptyset)$$

(6)

$$|aH| = |H| = |bH|$$

$$aH = H \iff a \in H$$

$$aH = H \iff a \in H$$

$$aH = H \implies a \in H$$

$$aH = H \iff a \in H$$

$$aH = H \implies a \in H$$

$$a \in aH = H$$

$$aH = H \iff a \in H$$

$$aH = H \implies a \in H$$

$$a \in H \implies aH = H$$

$$a \in aH = H$$

$$aH = H \iff a \in H$$

$$aH = H \implies a \in H$$

$$a \in H \implies aH = H$$

$$a \in aH = H$$

$$aH \subseteq HH \triangleq \{hh' \mid h, h' \in H\} = H$$

$$aH = H \iff a \in H$$

$$aH = H \implies a \in H$$

$$a \in H \implies aH = H$$

$$a \in aH = H$$

$$aH \subseteq HH \triangleq \{hh' \mid h, h' \in H\} = H$$

$$H = (aa^{-1})H = a(a^{-1}H) \subseteq aH$$

$$aH \leq G \iff a \in H$$

$$aH \leq G \iff a \in H$$

$$a \in H \implies aH \leq G$$

$$aH \leq G \iff a \in H$$

$$a \in H \implies aH \leq G$$

$$a \in H \implies aH = H \leq G$$

$$aH \leq G \iff a \in H$$

$$a \in H \implies aH \leq G$$

$$aH \leq G \implies a \in H$$

$$a \in H \implies aH = H \leq G$$

$$aH \leq G \iff a \in H$$

$$a \in H \implies aH \leq G$$

$$aH \leq G \implies a \in H$$

$$a \in H \implies aH = H \leq G \qquad a \in aH \implies a^{-1} \in aH \implies e \in aH$$

$$aH \leq G \iff a \in H$$

$$a \in H \implies aH \leq G$$

$$aH \leq G \implies a \in H$$

$$a \in H \implies aH = H \leq G$$

$$a \in aH \implies a^{-1} \in aH \implies e \in aH$$

$$\exists h \in H \ e = ah \implies a = h^{-1} \in H$$

$$aH = bH \iff a^{-1}b \in H$$

$$aH = bH \iff a^{-1}b \in H$$

$$a^{-1}b \in H \iff a^{-1}bH = H$$

$$aH = bH \iff a^{-1}b \in H$$

$$a^{-1}b \in H \iff a^{-1}bH = H$$

$$aH = bH \implies a^{-1}aH = a^{-1}bH \implies a^{-1}bH = H$$

$$aH = bH \iff a^{-1}b \in H$$

$$a^{-1}b \in H \iff a^{-1}bH = H$$

$$aH = bH \implies a^{-1}aH = a^{-1}bH \implies a^{-1}bH = H$$

$$a^{-1}bH = H \implies a(a^{-1}bH) = aH \implies bH = aH$$

$$\forall a, b \in G \ (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$\forall a, b \in G (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$\forall a, b \in G (aH \cap bH \neq \emptyset \rightarrow aH = bH)$$

$$\forall a, b \in G (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$\forall a, b \in G (aH \cap bH \neq \emptyset \rightarrow aH = bH)$$

Take any $g \in aH \cap bH$.

$$\forall a, b \in G (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$\forall a, b \in G (aH \cap bH \neq \emptyset \rightarrow aH = bH)$$

Take any $g \in aH \cap bH$.

$$\exists h_1, h_2 \in H (ah_1 = g = bh_2)$$

$$\forall a, b \in G \ (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$\forall a, b \in G \ (aH \cap bH \neq \emptyset \rightarrow aH = bH)$$

Take any $g \in aH \cap bH$.

$$\exists h_1, h_2 \in H \ (ah_1 = g = bh_2)$$


$$aH = a(h_1H) = (ah_1)H = (bh_2)H = b(h_2H) = bH$$

$$|aH| = |H| = |bH|$$

$$|aH| = |H| = |bH|$$

$$\sigma : aH \rightarrow H, \quad ah \mapsto h$$

A **balanced partition** of G by its subgraph H



$$\forall a, b \in G \ (aH = bH) \vee (aH \cap bH = \emptyset)$$

$$|aH| = |H| = |bH|$$

Definition (Index (H 在 G 中的指标/指数))

$$G/H = \{gH \mid g \in G\}$$

$$[G : H] \triangleq |G/H|$$

Definition (Index (H 在 G 中的指标/指数))

$$G/H = \{gH \mid g \in G\}$$

$$[G : H] \triangleq |G/H|$$

Theorem (Lagrange's Theorem)

Suppose that $H \leq G$. Then

$$|G| = [G : H] \cdot |H|$$

Theorem

$$H \leq G \implies |H| \mid |G|$$

Theorem

$$H \leq G \implies |H| \mid |G|$$

There are *no* subgroups of order 5, 7, or 8 of a group of order 12.

Theorem

$$H \leq G \implies |H| \mid |G|$$

There are *no* subgroups of order 5, 7, or 8 of a group of order 12.

Theorem

Let G be a finite group, $a \in G$.

Then the order of a divides the order of G .

Theorem

$$H \leq G \implies |H| \mid |G|$$

There are *no* subgroups of order 5, 7, or 8 of a group of order 12.

Theorem


Let G be a finite group, $a \in G$.

Then the order of a divides the order of G .

Theorem

- ▶ There are only 2 groups of order 4.
- ▶ There are only 2 groups of order 6.

Thank
You!



hfwei@hnu.edu.cn