8-paths-cycles: 路径与圈 (Paths and Cycles)

魏恒峰

hfwei@hnu.edu.cn

2026 年 04 月 23 日

Definitions Theorems Proofs

Seven Bridges of Königsberg (柯尼斯堡七桥问题)



Seven Bridges of Königsberg (柯尼斯堡七桥问题)



"to devise a walk through the city
that would cross each of those bridges once and only once"


KONINGSBERGA











"to devise a walk through the graph
that would cross each of those edges once and only once"

Definition (Graph (图))

An (undirected simple) graph is a pair $G = (V, E)$ where

- V is a set of vertices (顶点);
- $E \subseteq \{\{x, y\} \mid x, y \in V \wedge x \neq y\}$ is a set of edges





Undirected Multigraph



Undirected Multigraph Permitting Loops

Definition (Walk (道路))

Given a graph G, a (finite) walk in G is a sequence of edges of the form

$$\{v_0, v_1\}, \{v_1, v_2\}, \ldots, \{v_{m-1}, v_m\}.$$

$$(v_0 \to v_1 \to v_2 \to \cdots \to v_m)$$

Definition (Walk (道路))

Given a graph G, a (finite) walk in G is a sequence of edges of the form

$$\{v_0, v_1\}, \{v_1, v_2\}, \ldots, \{v_{m-1}, v_m\}.$$

$$(v_0 \to v_1 \to v_2 \to \cdots \to v_m)$$

It is a walk from the initial vertex v_0 to the final vertex v_m.

Definition (Walk (道路))

Given a graph G, a (finite) walk in G is a sequence of edges of the form

$$\{v_0, v_1\}, \{v_1, v_2\}, \ldots, \{v_{m-1}, v_m\}.$$

$$(v_0 \to v_1 \to v_2 \to \cdots \to v_m)$$

It is a walk from the initial vertex v_0 to the final vertex v_m.

$$D \to E \to B \to A \to E \to F$$



Definition (Walk (道路))

Given a graph G, a (finite) walk in G is a sequence of edges of the form

$$\{v_0, v_1\}, \{v_1, v_2\}, \ldots, \{v_{m-1}, v_m\}.$$

$$(v_0 \to v_1 \to v_2 \to \cdots \to v_m)$$

It is a walk from the initial vertex v_0 to the final vertex v_m.

$$D \to E \to B \to A \to E \to F$$



$$D \to E \to B \to E \to F$$

Definition (Trail (迹))

A trail is a walk in which all the edges are distinct.

Definition (Trail (迹))

A trail is a walk in which all the edges are distinct.

$$D \to E \to B \to A \to E \to F$$



$$D \to E \to B \to E \to F$$

Definition (Path (路径))

A path is a trial in which all vertices are distinct.

Definition (Path (路径))

A path is a trial in which all vertices are distinct.

$$D \to E \to B \to A \to E \to F$$



$$D \to E \to F$$

Definition (Closed Walk/Trail/Path)

A walk, trail, or path is closed if $v_0 = v_m$.

Definition (Closed Walk/Trail/Path)

A walk, trail, or path is closed if $v_0 = v_m$.



Definition (Closed Walk/Trail/Path)

A walk, trail, or path is closed if $v_0 = v_m$.



Definition (Cycle)

A cycle is a closed path with at least one edge.

"to devise a walk through the graph
that would cross each of those edges once and only once"



"to devise a walk through the graph
that would cross each of those edges once and only once"



to find a trail that contains all edges of the graph

$$v_0 \to v_1 \to \cdots \to v_i \to \cdots \to v_m$$



$$v_0 \to v_1 \to \cdots \to v_i \to \cdots \to v_m$$



$$v_i \notin \{v_0, v_m\} \implies \deg(v_i) \text{ is even}$$

Lemma (Necessary Condition for Eulerian Trails)

If a graph has Eulerian trails, then zero or two vertices have an odd degree.

Lemma (Necessary Condition for Eulerian Trails)

If a graph has Eulerian trails, then zero or two vertices have an odd degree.



4 vertices of odd degree $\Longrightarrow$ has no Eulerian trails

Theorem (Euler's Theorem)

A connected graph has Eulerian trails iff it contains zero or two vertices that have an odd degree.

Theorem (Euler's Theorem)

A connected graph has Eulerian trails iff it contains zero or two vertices that have an odd degree.

Euler stated but did *not* prove the "$\Longleftarrow$ (*if*)" direction.

Theorem (Euler's Theorem)

A connected graph has Eulerian trails iff it contains zero or two vertices that have an odd degree.

Euler stated but did *not* prove the "$\Longleftarrow$ (*if*)" direction.

Carl Hierholzer (1840 $\sim$ 1871) gave the first complete proof in 1873.

Theorem (Euler's Theorem)

A connected graph has Eulerian trails iff it contains zero or two vertices that have an odd degree.

Euler stated but did *not* prove the "$\Longleftarrow$ (*if*)" direction.

Carl Hierholzer (1840 $\sim$ 1871) gave the first complete proof in 1873.

Pierre-Henry Fleury gave another proof in 1883.

Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

(Such graphs are called Eulerian graphs.)

Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

(Such graphs are called Eulerian graphs.)



cycle + cycle + cycle

Lemma

If every vertex of a graph G has degree ≥ 2, then G contains a cycle.

Lemma

If every vertex of a graph G has degree ≥ 2, then G contains a cycle.

By Extremality.

Lemma

If every vertex of a graph G has degree ≥ 2, then G contains a cycle.

By Extremality.

Let $P : u \to \ldots$ be a *maximal* path in G.



Lemma

If every vertex of a graph G has degree ≥ 2, then G contains a cycle.

By Extremality.

Let $P : u \to \ldots$ be a *maximal* path in G.



$\deg(u) \geq 2$

Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

By induction on the number of edges m.

Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

By induction on the number of edges m.



$$G' = G - C$$

Theorem

Every Eulerian graph decomposes into cycles.





Theorem (Euler's Theorem (Carl Hierholzer))

A connected graph has Eulerian circuits/tours iff every vertex has even degree.

TONCAS

"The Obvious Necessary Conditions are Also Sufficient"

Dodecahedron: 12 faces, 20 vertices, and 30 edges



Dodecahedron: 12 faces, 20 vertices, and 30 edges



Is there a cycle that visits each *vertex* exactly once?



Is there a cycle that visits each *vertex* exactly once?

Definition (Hamiltonian Path)

A Hamiltonian path is a path that visits each vertex exactly once.



Definition (Hamiltonian Cycle)

A Hamiltonian cycle is a Hamiltonian path that is a cycle.

Definition (Hamiltonian Graph)

A graph is a Hamiltonian graph if it has a Hamiltonian cycle.

Definition (Hamiltonian Graph)

A graph is a Hamiltonian graph if it has a Hamiltonian cycle.

Definition (Semi-Hamiltonian Graph)

A non-Hamiltonian graph is semi-Hamiltonian if it has a Hamiltonian path.

William Rowan Hamilton
(1805 ~ 1865)

(October 16, 1843)

$$i^2 = j^2 = k^2 = ijk = -1$$





What is “THE” theorem for finding a Hamiltonian path/cycle or determining its existence?

What is “THE” theorem for finding a Hamiltonian path/cycle or determining its existence?

I do not know.

What is “THE” theorem for finding a Hamiltonian path/cycle or determining its existence?

I do not know.

Nobody knows now.

What is “THE” theorem for finding a Hamiltonian path/cycle or determining its existence?

I do not know.

Nobody knows now.

We will probably never know it.

What is “THE” theorem for finding a Hamiltonian path/cycle or determining its existence?

I do not know.

Nobody knows now.

We will probably never know it.

I wish I am wrong.

Theorem

The Hamiltonian Path/Cycle problem is NP-complete.

Typical (Positive/Negative) Graph Examples

Sufficient Conditions

Necessary Conditions

EXAMPLES

- Every cycle is Hamiltonian



C_6

- A complete graph (完全图) with $|V| > 2$ is Hamiltonian.

- A complete graph (完全图) with $|V| > 2$ is Hamiltonian.



K_1 K_2 K_3 K_5

- A complete bipartite graph $K_{m,n}$ is Hamiltonian iff $m = n \geq 2$.

Definition (Bipartite Graph (Bigraph; 二部图))

A bipartite graph $G = (U, V, E)$ is a graph whose vertices can be divided into two disjoint sets U and V such that every edge connects a vertex in U to one in V.



Definition (Bipartite Graph (Bigraph; 二部图))

A bipartite graph $G = (U, V, E)$ is a graph whose vertices can be divided into two disjoint sets U and V such that every edge connects a vertex in U to one in V.





Definition (Complete Bipartite Graph (Biclique; 完全二部图))

A complete bipartite graph $G = (U, V, E)$ is bipartite graph where every vertex of U is connected to every vertex of V.

$$K_{m,n} : m = |U|, n = |V|$$

Definition (Complete Bipartite Graph (Biclique; 完全二部图))

A complete bipartite graph $G = (U, V, E)$ is bipartite graph where every vertex of U is connected to every vertex of V.

$$K_{m,n} : m = |U|, n = |V|$$



$K_{1,7}$ (star)



$K_{3,3}$ (utility graph)



$K_{4,7}$

- A complete bipartite graph $K_{m,n}$ is Hamiltonian iff $m = n \geq 2$.

- A complete bipartite graph $K_{m,n}$ is Hamiltonian iff $m = n \geq 2$.





- Every platonic solid (正多面体), considered as a graph, is Hamiltonian.

- Every platonic solid (正多面体), considered as a graph, is Hamiltonian.

Tetrahedron



Cube



Octahedron



Dodecahedron



Icosahedron



Theorem

- *Petersen graph is not Hamiltonian.*

Julius Petersen (1839 ~ 1910)

Theorem

- *Petersen graph is not Hamiltonian.*

Julius Petersen (1839 ~ 1910)



Theorem

- *Petersen graph is not Hamiltonian.*

Julius Petersen (1839 ~ 1910)





Theorem

- *Petersen graph is not Hamiltonian.*

Julius Petersen (1839 ~ 1910)







Theorem

- *Petersen graph is not Hamiltonian.*

Julius Petersen (1839 ~ 1910)









Theorem

The Petersen graph is non-Hamiltonian.



Theorem

The Petersen graph is non-Hamiltonian.



$$\forall v \in V \;\; \deg(v) = 3 \qquad \text{3-regular graph}$$

The Petersen graph has *no* cycles of length < 5.

By Contradiction.

By Contradiction.

Suppose that it has a Hamiltonian cycle C.



By Contradiction.

Suppose that it has a Hamiltonian cycle C.



v_0 is adjacent to v_4, v_5, or v_6.

Case I: v_0 is adjacent to v_4



Case I: v_0 is adjacent to v_4



v_5 is adjacent to v_1 or v_9

Case I: v_0 is adjacent to v_4



v_5 is adjacent to v_1 or v_9

cycle of length 4!

CASE I: v_0 is adjacent to v_4



v_5 is adjacent to v_1 or v_9

cycle of length 4!

CASE II: v_0 is adjacent to v_5



CASE I: v_0 is adjacent to v_4



v_5 is adjacent to v_1 or v_9

cycle of length 4!

CASE II: v_0 is adjacent to v_5



v_1 is adjacent to v_7

CASE I: v_0 is adjacent to v_4



v_5 is adjacent to v_1 or v_9

cycle of length 4!

CASE II: v_0 is adjacent to v_5



v_1 is adjacent to v_7

back to CASE I!

"If G has enough edges, then G is Hamiltonian."

Theorem (Ore's Theorem, 1960)

Let G be a simple graph with $n \geq 3$ vertices. If

$$\deg(u) + \deg(v) \geq n$$

for each pair of non-adjacent vertices u and v, then G is Hamiltonian.

Theorem (Ore's Theorem, 1960)

Let G be a simple graph with $n \geq 3$ vertices. If

$$\deg(u) + \deg(v) \geq n$$

for each pair of non-adjacent vertices u and v, then G is Hamiltonian.



By Contradiction.

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

We need to derive a contradiction.

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

We need to derive a contradiction.

Adding edges cannot violate the Ore's Condition.

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

We need to derive a contradiction.

Adding edges cannot violate the Ore's Condition.

By Extremality.

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

We need to derive a contradiction.

Adding edges cannot violate the Ore's Condition.

By Extremality.

We can obtain *maximal* non-Hamiltonian graphs:

By Contradiction.

Suppose that G meets the Ore's Condition, but G is not Hamiltonian.

We need to derive a contradiction.

Adding edges cannot violate the Ore's Condition.

By Extremality.

We can obtain *maximal* non-Hamiltonian graphs:

adding any edge gives a Hamiltonian graph.

By its "maximality", G contains a Hamiltonian path

(G is a semi-Hamiltonian graph)

$$v_1 \to v_2 \to \cdots \to v_n$$

By its "maximality", G contains a Hamiltonian path

(G is a semi-Hamiltonian graph)

$$v_1 \to v_2 \to \cdots \to v_n$$

v_1 and v_n are non-adjacent

By its "maximality", G contains a Hamiltonian path

(G is a semi-Hamiltonian graph)

$$v_1 \to v_2 \to \cdots \to v_n$$

v_1 and v_n are non-adjacent

$$\deg(v_1) + \deg(v_2) \geq n$$

By its "maximality", G contains a Hamiltonian path

(G is a semi-Hamiltonian graph)

$$v_1 \to v_2 \to \cdots \to v_n$$

v_1 and v_n are non-adjacent

$$\deg(v_1) + \deg(v_2) \geq n$$



There must be some vertex v_i adjacent to v_1

such that v_{i-1} is adjacent to v_n.

Theorem (Dirac's Theorem (1952; Gabriel Andrew Dirac))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\forall v \in V.\ \deg(v) \geq n/2.$$

Theorem (Dirac's Theorem (1952; Gabriel Andrew Dirac))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\forall v \in V.\ \deg(v) \geq n/2.$$

$$\delta(G) \triangleq \min_{v \in V} \deg(v) \qquad \Delta(G) \triangleq \max_{v \in V} \deg(v)$$

$$\delta(G) \geq n/2$$

Theorem (Dirac's Theorem (1952; Gabriel Andrew Dirac))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\forall v \in V.\ \deg(v) \geq n/2.$$

$$\delta(G) \triangleq \min_{v \in V} \deg(v) \qquad \Delta(G) \triangleq \max_{v \in V} \deg(v)$$

$$\delta(G) \geq n/2$$

Family [edit | edit source]

He was born Balázs Gábor in Budapest, to Richárd Balázs, a military officer and businessman, and Margit "Manci" Wigner (sister of Eugene Wigner).[5] When his mother married Paul Dirac in 1937, he and his sister resettled in England and were formally adopted, changing their family name to Dirac.[6]

Theorem (Dirac's Theorem (1952))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\delta(G) \geq n/2$$

Theorem (Dirac's Theorem (1952))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\delta(G) \geq n/2$$

$$\delta(G) = \lfloor (n-1)/2 \rfloor$$

Theorem (Dirac's Theorem (1952))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\delta(G) \geq n/2$$

$$\delta(G) = \lfloor (n-1)/2 \rfloor$$

Counterexample: $K_{\lfloor (n+1)/2 \rfloor}$ and $K_{\lceil (n+1)/2 \rceil}$ sharing a vertex

Theorem (Dirac's Theorem (1952))

A simple graph $G = (V, E)$ with $n \geq 3$ vertices is Hamiltonian

$$\delta(G) \geq n/2$$

$$\delta(G) = \lfloor (n-1)/2 \rfloor$$

Counterexample: $K_{\lfloor (n+1)/2 \rfloor}$ and $K_{\lceil (n+1)/2 \rceil}$ sharing a vertex

$$n = 6$$

"If G is Hamiltonian, then G must be somewhat connected."

"If G is Hamiltonian, then G must be somewhat connected."

"If G is not so connected, then G is non-Hamiltonian."

Theorem

If $G = (V, E)$ is Hamiltonian, then for each nonempty set $S \subset V$, the graph $G - S$ has $\leq |S|$ components.

Definition (Components (连通分支))

A component of an undirected graph is an subgraph in which any two vertices are connected to each other by paths, and which is connected to no additional vertices in the rest of the graph.



Definition (Components (连通分支))

A component of an undirected graph is an subgraph in which any two vertices are connected to each other by paths, and which is connected to no additional vertices in the rest of the graph.



$$c(G) = 3$$

Theorem

If $G = (V, E)$ is Hamiltonian, then

$$\forall S \subset V \; c(G - S) \leq |S|.$$

Theorem

If $G = (V, E)$ is Hamiltonian, then

$$\forall S \subset V \; c(G - S) \leq |S|.$$

$$S \neq V$$

Theorem

If $G = (V, E)$ is Hamiltonian, then

$$\forall S \subset V \; c(G - S) \leq |S|.$$

$$S \neq V$$

$$K_{\lfloor (n+1)/2 \rfloor} \text{ and } K_{\lceil (n+1)/2 \rceil} \text{ sharing a vertex}$$

- A complete bipartite graph $K_{m,n}$ is Hamiltonian iff $m = n \geq 2$.

- A complete bipartite graph $K_{m,n}$ is Hamiltonian iff $m = n \geq 2$.



A non-Hamiltonian bipartite graph with $m = n = 10$

Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \ c(G - S) \leq |S|.$$

Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \ c(G - S) \leq |S|.$$



Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \; c(G - S) \leq |S|.$$



"When a Hamiltonian cycle leaves a component of $G - S$, it can go only to a distinct vertex in S."

Theorem

If $G = (V, E)$ *is Hamiltonian,*

$$\forall S \subset V \; c(G - S) \leq |S|.$$

Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \; c(G - S) \leq |S|.$$

The condition is *not* sufficient.

Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \; c(G - S) \leq |S|.$$

The condition is *not* sufficient.



Theorem

If $G = (V, E)$ is Hamiltonian,

$$\forall S \subset V \; c(G - S) \leq |S|.$$

The condition is *not* sufficient.



All edges incident to vertices of degree 2 must be used.

Chessboard Problem ("马踏棋盘" 问题)

Is it possible for a "knight" to visit every field of a 4×4 or 5×5 chessboard exactly once and return to the starting point?





5×5



5×5



$$G = (U, V, E) : |U| = 12, |V| = 13$$

4×4

4×4

Removing the middle 4 squares leaves ≥ 5 components.

Chessboard Problem





$4 \times n$

Definition (Travelling Salesman/Salesperson Problem (TSP; 旅行商问题))

Given a list of cities and the distances between each pair of cities, what is the shortest possible route that visits each city exactly once and returns to the origin city?




William J. Cook
In Pursuit of the Traveling Salesman
Mathematics at the Limits of Computation

Thank You!



hfwei@hnu.edu.cn