


# Asymptotics, Recurrences, and Divide and Conquer

Hengfeng Wei

hfwei@nju.edu.cn

April 17, 2018



Problem  $P$

Algorithm  $A$

Problem  $P$       Algorithm  $A$

Inputs:  $\mathcal{X}_n$  of size  $n$

Problem  $P$       Algorithm  $A$

Inputs:  $\mathcal{X}_n$  of size  $n$

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

## Problem $P$      Algorithm $A$

Inputs:  $\mathcal{X}_n$  of size  $n$

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

## Problem $P$      Algorithm $A$

Inputs:  $\mathcal{X}_n$  of size  $n$

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

$$A(n) = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$

## Problem $P$      Algorithm $A$

Inputs:  $\mathcal{X}_n$  of size  $n$

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

$$A(n) = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)]$$



## Average-case Time Complexity (Problem 1.8)

Input :  $r \in [1, n]$ ,  $r \in \mathbb{Z}^+$

$$P\{r = i\} = \begin{cases} \frac{1}{n}, & 1 \leq i \leq \frac{n}{4} \\ \frac{2}{n}, & \frac{n}{4} < i \leq \frac{n}{2} \\ \frac{1}{2n}, & \frac{n}{2} < i \leq n \end{cases}$$

## Average-case Time Complexity (Problem 1.8)

Input :  $r \in [1, n]$ ,  $r \in \mathbb{Z}^+$

$$P\{r = i\} = \begin{cases} \frac{1}{n}, & 1 \leq i \leq \frac{n}{4} \\ \frac{2}{n}, & \frac{n}{4} < i \leq \frac{n}{2} \\ \frac{1}{2n}, & \frac{n}{2} < i \leq n \end{cases}$$

$$\begin{aligned} A &= \sum_{X \in \mathcal{X}} T(X) \cdot P(X) \\ &= T(1)P(1) + T(2)P(2) + \cdots + T(n)P(n) \\ &= \frac{n}{4} \times 10 \times \frac{1}{n} + \frac{n}{4} \times 20 \times \frac{2}{n} + \frac{n}{4} \times 30 \times \frac{1}{2n} + \frac{n}{4} \times n \times \frac{1}{2n} \\ &= \frac{1}{8}n + \frac{65}{4} \end{aligned}$$

# Average-case Analysis of Quicksort

$$A(n) = n - 1 + \frac{1}{n} \sum_{i=0}^{n-1} (A(i) + A(n - i - 1))$$

$$A(n) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)] = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$

# Average-case Analysis of Quicksort

$$A(n) = n - 1 + \frac{1}{n} \sum_{i=0}^{i=n-1} (A(i) + A(n - i - 1))$$

$$A(n) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)] = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$

$$\begin{aligned} A(n) &= \mathbb{E}[T(X)] \\ &= \mathbb{E}[\mathbb{E}[T(X)|I]] \\ &= \sum_{i=0}^{i=n-1} P(I = i) \mathbb{E}[T(X) \mid I = i] \\ &= \sum_{i=0}^{i=n-1} \frac{1}{n} [n - 1 + A(i) + A(n - i - 1)] \end{aligned}$$

# Mathematical Induction



## Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

---

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```

---

## Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

---

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```

---

Loop invariant (after the  $k$ -th loop):

## Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

---

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```


---

Loop invariant (after the  $k$ -th loop):

$$\mathcal{I} : p = \sum_{i=n-k}^{i=n} a_i x^{k-(n-i)}$$



$$\mathcal{I}: p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$



$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$



When you are in an exam:

20% : Finding  $\mathcal{I}$

80% : Proving  $\mathcal{I}$  by PMI

$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$

Proof.

Prove by mathematical induction on non-negative integer  $k$ ,  
the number of loops.

$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$

Proof.

Prove by mathematical induction on non-negative integer  $k$ ,  
the number of loops.

Basis:

$$k = 0 : p = a_n = \mathcal{I}_0$$

Inductive Hypothesis:

Inductive Step:



## Integer Multiplication (Problem 1.6)

---

```
1: procedure INT-MULT( $y, z$ )  
2:   if  $z = 0$  then  
3:     return 0  
4:   return INT-MULT( $cy, \lfloor \frac{z}{c} \rfloor$ ) +  $y(z \bmod c)$ 
```

---

## Integer Multiplication (Problem 1.6)

---

```
1: procedure INT-MULT( $y, z$ )  
2:   if  $z = 0$  then  
3:     return 0  
4:   return INT-MULT( $cy, \lfloor \frac{z}{c} \rfloor$ ) +  $y(z \bmod c)$ 
```

---

Proof.

Prove by mathematical induction on non-negative integer  $z$ .



2-tree; full binary tree (Problem 2.5)

$$n_0 = n_2 + 1$$

Proof.

2-tree; full binary tree (Problem 2.5)

$$n_0 = n_2 + 1$$

Proof.

Prove by mathematical induction on *the structure of binary tree*.

