

# Minimum Spanning Tree (MST)

Hengfeng Wei

hfwei@nju.edu.cn

June 19, 2018

## Cut Property

$$G = (V, E, w)$$

## Cut Property (Strong)

- ▶  $X$  is some part of an MST  $T$  of  $G$
- ▶ Any **cut**  $(S, V \setminus S)$  s.t.  $X$  does **not** cross  $(S, V \setminus S)$   $\hat{=}$
- ▶ Let  $e$  be **a** lightest edge across  $(S, V \setminus S)$

## Cut Property (Strong)

- ▶  $X$  is some part of an MST  $T$  of  $G$
- ▶ Any **cut**  $(S, V \setminus S)$  s.t.  $X$  does **not** cross  $(S, V \setminus S)$
- ▶ Let  $e$  be **a** lightest edge across  $(S, V \setminus S)$

Then  $X \cup \{e\}$  is some part of an MST  $T'$  of  $G$ .

## Cut Property (Strong)


- ▶  $X$  is some part of an MST  $T$  of  $G$
- ▶ Any **cut**  $(S, V \setminus S)$  s.t.  $X$  does **not** cross  $(S, V \setminus S)$
- ▶ Let  $e$  be **a** lightest edge across  $(S, V \setminus S)$

Then  $X \cup \{e\}$  is some part of an MST  $T'$  of  $G$ .


Correctness of Prim's and Kruskal's algorithms.

## By Exchange Argument.

## By Exchange Argument.



## By Exchange Argument.



$$T + \{e\} - \{e'\}$$



## Cut Property (Weak)

A cut  $(S, V \setminus S)$

Let  $e = (u, v)$  be **a** minimum-weight edge across  $(S, V \setminus S)$

## Cut Property (Weak)

A cut  $(S, V \setminus S)$

Let  $e = (u, v)$  be *a* minimum-weight edge across  $(S, V \setminus S)$


Then  $e$  must be in *some* MST of  $G$ .

## Cut Property (Weak)

A cut  $(S, V \setminus S)$

Let  $e = (u, v)$  be **a** minimum-weight edge across  $(S, V \setminus S)$

Then  $e$  must be in **some** MST of  $G$ .




## Cut Property (Weak)

A cut  $(S, V \setminus S)$

Let  $e = (u, v)$  be **a** minimum-weight edge across  $(S, V \setminus S)$

Then  $e$  must be in **some** MST of  $G$ .



“a”  $\rightarrow$  “the”  $\Rightarrow$  “some”  $\rightarrow$  “any”

## Converse of Cut Property (Weak)

$$e = (u, v) \in \exists \text{ MST } T \text{ of } G$$

$\implies$


$e$  is a lightest edge across some cut  $(S, V \setminus S)$

## Converse of Cut Property (Weak)

$$e = (u, v) \in \exists \text{ MST } T \text{ of } G$$

$\implies$

$e$  is a lightest edge across some cut  $(S, V \setminus S)$



$$T' = \underbrace{T - \{e\}}_{\text{to find } (S, V \setminus S)} + \underbrace{\{e'\}}_{\exists?}$$

## Application of Cut Property [Problem: 10.15 (3)]

$e \in G$  is a lightest edge  $\implies e \in \exists$  MST of  $G$

## Application of Cut Property [Problem: 10.15 (3)]

$e \in G$  is a lightest edge  $\implies e \in \exists$  MST of  $G$

$$(S = \{u\}, V \setminus S)$$



## Application of Cut Property [Problem: 10.15 (4)]

$e \in G$  is the unique lightest edge  $\Rightarrow e \in \forall \text{ MST}$

## Application of Cut Property [Problem: 10.15 (4)]

$e \in G$  is the unique lightest edge  $\Rightarrow e \in \forall \text{ MST}$

By contradiction.

$$e \notin T : T' = T + \{e\} - \{e'\} \implies w(T') < w(T)$$

Wrong divide-and-conquer algorithm for MST [Problem: 10.21]


$$(V_1, V_2) : ||V_1| - |V_2|| \leq 1$$

$$T_1 + T_2 + \{e\} : e \text{ is a lightest edge across } (V_1, V_2)$$

## Wrong divide-and-conquer algorithm for MST [Problem: 10.21]

$$(V_1, V_2) : ||V_1| - |V_2|| \leq 1$$

$$T_1 + T_2 + \{e\} : e \text{ is a lightest edge across } (V_1, V_2)$$




## Cycle Property

$$G = (V, E, w)$$

## Cycle property [Problem: 10.19(b)]

- ▶ Let  $C$  be any cycle in  $G$
- ▶ Let  $e = (u, v)$  be **a** maximum-weight edge in  $C$


Then  $\exists$  MST  $T$  of  $G : e \notin T$ .



## Cycle property [Problem: 10.19(b)]

- ▶ Let  $C$  be any cycle in  $G$
- ▶ Let  $e = (u, v)$  be **a** maximum-weight edge in  $C$

Then  $\exists$  MST  $T$  of  $G : e \notin T$ .



$$T' = \underbrace{T - \{e\}}_{e \in T} + \{e'\}$$

Anti-Kruskal algorithm [Problem: 10.19(c)]

Reverse-delete algorithm (wiki)

$$O\left(m \log n (\log \log n)^3\right)$$



Anti-Kruskal algorithm [Problem: 10.19(c)]

Reverse-delete algorithm (wiki)

$$O\left(m \log n (\log \log n)^3\right)$$

Proof.

**Invariant:** If  $F$  is the set of edges remained at the end of the while loop, then there is some MST that are a subset of  $F$ . □

Anti-Kruskal algorithm [Problem: 10.19(c)]

Reverse-delete algorithm (wiki)

$$O(m \log n (\log \log n)^3)$$

Proof.

**Invariant:** If  $F$  is the set of edges remained at the end of the while loop, then there is some MST that are a subset of  $F$ . □

*“On the Shortest Spanning Subtree of a Graph  
and the Traveling Salesman Problem”* — Kruskal, 1956.

## Application of Cycle Property [Problem: 10.15 (1)]

$G = (V, E), |E| > |V| - 1, e$  unique maximum-weighted edge

$\Rightarrow$  ?

$e \notin$  any MST

## Application of Cycle Property [Problem: 10.15 (1)]

$G = (V, E), |E| > |V| - 1, e$  unique maximum-weighted edge

$\Rightarrow$  ?

$e \notin$  any MST

Bridge

## Application of Cycle Property [Problem: 10.15 (2)]

$C \subseteq G, e \in C, e$  is the unique maximum-weighted edge of  $G$

$\implies$

$e \notin \text{any MST of } G$

## Application of Cycle Property [Problem: 10.15 (2)]

$C \subseteq G, e \in C, e$  is the unique maximum-weighted edge of  $G$

$\implies$

$e \notin \text{any MST of } G$

By contradiction.


$$T' = T - \{e\} + \{e'\}$$

## Application of Cycle Property [Problem: 10.15 (5)]

$C \subseteq G, e \in C, e$  is the unique lightest edge of  $C \implies e \in \forall \text{ MST}$

## Application of Cycle Property [Problem: 10.15 (5)]

$C \subseteq G, e \in C, e$  is the unique lightest edge of  $C \implies e \in \forall \text{ MST}$





## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

By contradiction.

## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

By contradiction.

$\exists$  MSTs  $T_1 \neq T_2$

## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

By contradiction.

$\exists$  MSTs  $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

By contradiction.

$\exists$  MSTs  $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

$$e = \min \Delta E$$

## Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights  $\implies$  unique MST.

By contradiction.


$\exists$  MSTs  $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$


$$e = \min \Delta E$$

$$e \in T_1 \setminus T_2 \text{ (w.l.o.g.)}$$

$$e \in T_1 \setminus T_2$$




$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$




$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \Rightarrow C$$

$$\exists (e' \in C) \notin T_1 \Rightarrow e' \in T_2 \setminus T_1 \Rightarrow e' \in \Delta E$$


$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$

$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists(e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$


$$T' = T_2 + \{e\} - \{e'\} \implies w(T') < w(T_2)$$

## Condition for Uniqueness of MST [Problem: 10.18 (2)]

Unique MST  $\not\Rightarrow$  Equal weights.

## Condition for Uniqueness of MST [Problem: 10.18 (2)]

Unique MST  $\not\Rightarrow$  Equal weights.




## Unique MST [Problem: 10.21(3)]

Unique MST  $\nRightarrow$  Minimum-weight edge across any cut is unique.


## Unique MST [Problem: 10.21(3)]

Unique MST  $\not\Rightarrow$  Minimum-weight edge across any cut is unique.



## Unique MST [Problem: 10.21(3)]

Unique MST  $\not\Rightarrow$  Minimum-weight edge across any cut is unique.



## Theorem

*Minimum-weight edge across any cut is unique  $\implies$  Unique MST.*




## Unique MST [Problem: 10.21 (3)]

Unique MST  $\not\Rightarrow$  Maximum-weight edge in any cycle is unique.


## Unique MST [Problem: 10.21 (3)]

Unique MST  $\not\Rightarrow$  Maximum-weight edge in any cycle is unique.



## Unique MST [Problem: 10.21 (3)]

Unique MST  $\not\Rightarrow$  Maximum-weight edge in any cycle is unique.




## Theorem (Conjecture)

*Maximum-weight edge in any cycle is unique  $\implies$  Unique MST.*


## Unique MST [Problem: 10.21 (3)]

Unique MST  $\Rightarrow$  Maximum-weight edge in any cycle is unique.



## Theorem (Conjecture)

Maximum-weight edge in any cycle is unique  $\Rightarrow$  Unique MST.



## Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

## Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

Ties breaking in Prim's and Kruskal's algorithms

## Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

Ties breaking in Prim's and Kruskal's algorithms

Proof.

Cut property and Cycle property.



## Adding a Vertex $v$ to MST $T$ [Problem: 10.7]

$$G' = (V', E') : V' = V + \{v\}, E' = E + E_v$$

To find an MST  $T'$  of  $G'$ .

$$O((m+n) \log n) \quad (\text{recompute on } G')$$



## Adding a Vertex $v$ to MST $T$ [Problem: 10.7]

$$G' = (V', E') : V' = V + \{v\}, E' = E + E_v$$

To find an MST  $T'$  of  $G'$ .

$$O((m + n) \log n) \quad (\text{recompute on } G')$$

### Theorem

*There exists an MST of  $G'$  that includes no edges in  $G \setminus T$ .*

## Adding a Vertex $v$ to MST $T$ [Problem: 10.7]

$$G' = (V', E') : V' = V + \{v\}, E' = E + E_v$$

To find an MST  $T'$  of  $G'$ .

$$O((m+n) \log n) \quad (\text{recompute on } G')$$

## Theorem

*There exists an MST of  $G'$  that includes no edges in  $G \setminus T$ .*

$$O(n \log n) \quad (\text{recompute on } G'' = (V + \{v\}, T + E_v))$$

## Adding a Vertex $v$ to MST $T$ [Problem: 10.7]

$$G' = (V', E') : V' = V + \{v\}, E' = E + E_v$$

To find an MST  $T'$  of  $G'$ .

$$O((m+n) \log n) \quad (\text{recompute on } G')$$

## Theorem

*There exists an MST of  $G'$  that includes no edges in  $G \setminus T$ .*

$$O(n \log n) \quad (\text{recompute on } G'' = (V + \{v\}, T + E_v))$$

$$O(n)$$

## Adding a Vertex $v$ to MST $T$ [Problem: 10.7]

$$G' = (V', E') : V' = V + \{v\}, E' = E + E_v$$

To find an MST  $T'$  of  $G'$ .

$$O((m+n) \log n) \quad (\text{recompute on } G')$$

## Theorem

*There exists an MST of  $G'$  that includes no edges in  $G \setminus T$ .*

$$O(n \log n) \quad (\text{recompute on } G'' = (V + \{v\}, T + E_v))$$

$$O(n)$$

“On Finding and Updating Spanning Trees and Shortest Paths”, 1975

“Algorithms for Updating Minimum Spanning Trees”, 1978

## Feedback Edge Set: [Problem: 10.8]

1. Maximum spanning tree
  2. (Minimum) feedback edge set:
    - ▶ a set of edges which, when removed from the graph, leave an acyclic graph  $G'$
    - ▶ assuming  $G$  is connected  $\Rightarrow G'$  is connected
    - ▶ feedback *arc* set: “cycle”  $\Rightarrow$  circular dependency
- 
- ▶  $G'$  is connected + acyclic  $\Rightarrow G'$  is an ST
  - ▶  $\text{FES} \Leftrightarrow G \setminus \text{Max-ST}$

## Edge Weights


- ▶ [Problem: 10.15 (7)]: negative edges for Prim algorithm
- ▶ [Problem: 10.16]:  $w'(e) = (w(e))^2$

## MST with Specified Leaves: [Problem: 10.11]

- ▶  $G = (V, E), U \subset V$
- ▶ finding an MST with  $U$  as leaves

## MST with Specified Leaves: [Problem: 10.11]

- ▶  $G = (V, E), U \subset V$
- ▶ finding an MST with  $U$  as leaves




MST  $T'$  of  $G' = G \setminus U$



## MST with Specified Leaves: [Problem: 10.11]

- ▶  $G = (V, E), U \subset V$
- ▶ finding an MST with  $U$  as leaves



MST  $T'$  of  $G' = G \setminus U$

Attach  $\forall u \in U$  to  $T'$  (with lightest edge)

## MST with Specified Edges: [Problem: 10.13]

$$G = (V, E), \quad S \subset E \text{ (no cycle in } S\text{)}$$

To find an MST with  $S$  as edges.

## MST with Specified Edges: [Problem: 10.13]


$$G = (V, E), \quad S \subset E \text{ (no cycle in } S\text{)}$$

To find an MST with  $S$  as edges.

$G \rightarrow G'$  : contract each component of  $S$  to a vertex

## MST vs. Shortest Paths [Problem: 10.15 (6)]

✗ The shortest path between two nodes is necessarily part of some MST.



## Sharing edges [Problem: 3.6.5]

- ▶  $G = (V, E), w(e) > 0$
- ▶ Given  $s$ : all sssp trees from  $s$  must share some edge with **all** (some) MSTs of  $G$

✓  $w > 0$ ; Vertex  $s$ ; shortest-path tree of  $s$  and some MST share a common edge [Problem: 10.9]

## Solution

$E'$ : lightest edges leaving  $s$

- ▶ any MST  $T$  of  $G$ :  $T \cap E' \neq \emptyset$
- ▶  $E' \subset \forall$  sssp trees

