

Journal

Numerische Mathematik

in: Numerische Mathematik | Journal

476 page(s)

Terms and Conditions

The Göttingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library. Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept there Terms and Conditions. Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact:

Niedersächsische Staats- und Universitätsbibliothek
Digitalisierungszentrum
37070 Goettingen
Germany
Email: gdz@sub.uni-goettingen.de

Purchase a CD-ROM

The Goettingen State and University Library offers CD-ROMs containing whole volumes / monographs in PDF for Adobe Acrobat. The PDF-version contains the table of contents as bookmarks, which allows easy navigation in the document. For availability and pricing, please contact:

Niedersaechisische Staats- und Universitaetsbibliothek Goettingen - Digitalisierungszentrum
37070 Goettingen, Germany, Email: gdz@sub.uni-goettingen.de

Gaussian Elimination is not Optimal

VOLKER STRASSEN*

Received December 12, 1968

1. Below we will give an algorithm which computes the coefficients of the product of two square matrices A and B of order n from the coefficients of A and B with less than $4.7 \cdot n^{\log 7}$ arithmetical operations (all logarithms in this paper are for base 2, thus $\log 7 \approx 2.8$; the usual method requires approximately $2n^3$ arithmetical operations). The algorithm induces algorithms for inverting a matrix of order n , solving a system of n linear equations in n unknowns, computing a determinant of order n etc. all requiring less than const $n^{\log 7}$ arithmetical operations.

This fact should be compared with the result of KLYUYEV and KOKOVKIN-SHCHERBAK [1] that Gaussian elimination for solving a system of linear equations is optimal if one restricts oneself to operations upon rows and columns as a whole. We also note that WINOGRAD [2] modifies the usual algorithms for matrix multiplication and inversion and for solving systems of linear equations, trading roughly half of the multiplications for additions and subtractions.

It is a pleasure to thank D. BRILLINGER for inspiring discussions about the present subject and ST. COOK and B. PARLETT for encouraging me to write this paper.

2. We define algorithms $\alpha_{m,k}$ which multiply matrices of order $m2^k$, by induction on k : $\alpha_{m,0}$ is the usual algorithm for matrix multiplication (requiring m^3 multiplications and $m^2(m-1)$ additions). $\alpha_{m,k}$ already being known, define $\alpha_{m,k+1}$ as follows:

If A, B are matrices of order $m2^{k+1}$ to be multiplied, write

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad AB = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

where the A_{ik}, B_{ik}, C_{ik} are matrices of order $m2^k$. Then compute

$$\begin{aligned} \text{I} &= (A_{11} + A_{22})(B_{11} + B_{22}), \\ \text{II} &= (A_{21} + A_{22})B_{11}, \\ \text{III} &= A_{11}(B_{12} - B_{22}), \\ \text{IV} &= A_{22}(-B_{11} + B_{21}), \\ \text{V} &= (A_{11} + A_{12})B_{22}, \\ \text{VI} &= (-A_{11} + A_{21})(B_{11} + B_{12}), \\ \text{VII} &= (A_{12} - A_{22})(B_{21} + B_{22}), \end{aligned}$$

* The results have been found while the author was at the Department of Statistics of the University of California, Berkeley. The author wishes to thank the National Science Foundation for their support (NSF GP-7454).

$$\begin{aligned}
C_{11} &= I + IV - V + VII, \\
C_{21} &= II + IV, \\
C_{12} &= III + V, \\
C_{22} &= I + III - II + VI,
\end{aligned}$$

using $\alpha_{m,k}$ for multiplication and the usual algorithm for addition and subtraction of matrices of order $m2^k$.

By induction on k one easily sees

Fact 1. $\alpha_{m,k}$ computes the product of two matrices of order $m2^k$ with $m^3 7^k$ multiplications and $(5 + m)m^2 7^k - 6(m2^k)^2$ additions and subtractions of numbers.

Thus one may multiply two matrices of order 2^k with 7^k numbermultiplications and less than $6 \cdot 7^k$ additions and subtractions.

Fact 2. The product of two matrices of order n may be computed with $< 4.7 \cdot n^{\log 7}$ arithmetical operations.

Proof. Put

$$\begin{aligned}
k &= [\log n - 4], \\
m &= [n2^{-k}] + 1,
\end{aligned}$$

then

$$n \leq m2^k.$$

Imbedding matrices of order n into matrices of order $m2^k$ reduces our task to that of estimating the number of operations of $\alpha_{m,k}$. By Fact 1 this number is

$$\begin{aligned}
&(5 + 2m)m^2 7^k - 6(m2^k)^2 \\
&< (5 + 2(n2^{-k} + 1))(n2^{-k} + 1)^2 7^k \\
&< 2n^3 (7/8)^k + 12.03 n^2 (7/4)^k
\end{aligned}$$

(here we have used $16 \cdot 2^k \leq n$)

$$\begin{aligned}
&= (2(8/7)^{\log n - k} + 12.03(4/7)^{\log n - k}) n^{\log 7} \\
&\leq \max_{4 \leq t \leq 5} (2(8/7)^t + 12.03(4/7)^t) n^{\log 7} \\
&\leq 4.7 \cdot n^{\log 7}
\end{aligned}$$

by a convexity argument.

We now turn to matrix inversion. To apply the algorithms below it is necessary to assume not only that the matrix is invertible but that all occurring divisions make sense (a similar assumption is of course necessary for Gaussian elimination).

We define algorithms $\beta_{m,k}$ which invert matrices of order $m2^k$, by induction on k : $\beta_{m,0}$ is the usual Gaussian elimination algorithm. $\beta_{m,k}$ already being known, define $\beta_{m,k+1}$ as follows:

If A is a matrix of order $m2^{k+1}$ to be inverted, write

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad A^{-1} = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

where the A_{ik} , C_{ik} are matrices of order $m2^k$. Then compute

$$\begin{aligned}
 \text{I} &= A_{11}^{-1}, \\
 \text{II} &= A_{21} \text{I}, \\
 \text{III} &= \text{I} A_{12}, \\
 \text{IV} &= A_{21} \text{III}, \\
 \text{V} &= \text{IV} - A_{22}, \\
 \text{VI} &= \text{V}^{-1}, \\
 C_{12} &= \text{III} \cdot \text{VI}, \\
 C_{21} &= \text{VI} \cdot \text{II}, \\
 \text{VII} &= \text{III} \cdot C_{21}, \\
 C_{11} &= \text{I} - \text{VII}, \\
 C_{22} &= -\text{VI}
 \end{aligned}$$

using $\alpha_{m,k}$ for multiplication, $\beta_{m,k}$ for inversion and the usual algorithm for addition or subtraction of two matrices of order $m2^k$.

By induction on k one easily sees

Fact 3. $\beta_{m,k}$ computes the inverse of a matrix of order $m2^k$ with $m2^k$ divisions, $\leq \frac{9}{5} m^3 7^k - m2^k$ multiplications and $\leq \frac{9}{5} (5+m) m^2 7^k - 7(m2^k)^2$ additions and subtractions of numbers. The next Fact follows in the same way as Fact 2.

Fact 4. The inverse of a matrix of order n may be computed with $< 5.64 \cdot n^{\log 7}$ arithmetical operations.

Similar results hold for solving a system of linear equations or computing a determinant (use $\text{Det } A = (\text{Det } A_{11}) \text{Det } (A_{22} - A_{21} A_{11}^{-1} A_{12})$).

References

1. KLYUYEV, V. V., and N. I. KOKOVKIN-SHCHERBAK: On the minimization of the number of arithmetic operations for the solution of linear algebraic systems of equations. Translation by G. I. TEE: Technical Report CS 24, June 14, 1965, Computer Science Dept., Stanford University.
2. WINOGRAD, S.: A new algorithm for inner product. IBM Research Report RC-1943, Nov. 21, 1967.

Prof. VOLKER STRASSEN
Seminar für angewandte Mathematik
der Universität
8032 Zürich, Freie Str. 36