


Asymptotics, Recurrences, and Divide and Conquer

Hengfeng Wei

hfwei@nju.edu.cn

April 17, 2018





Problem P

Algorithm A

Problem P Algorithm A

Inputs: \mathcal{X}_n of size n

Problem P Algorithm A

Inputs: \mathcal{X}_n of size n

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

Problem P Algorithm A

Inputs: \mathcal{X}_n of size n

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

Problem P Algorithm A

Inputs: \mathcal{X}_n of size n

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

$$A(n) = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$

Problem P Algorithm A

Inputs: \mathcal{X}_n of size n

$$W(n) = \max_{X \in \mathcal{X}_n} T(X)$$

$$B(n) = \min_{X \in \mathcal{X}_n} T(X)$$

$$A(n) = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)]$$

Average-case Time Complexity (Problem 1.8)

Input : $r \in [1, n]$, $r \in \mathbb{Z}^+$

$$P\{r = i\} = \begin{cases} \frac{1}{n}, & 1 \leq i \leq \frac{n}{4} \\ \frac{2}{n}, & \frac{n}{4} < i \leq \frac{n}{2} \\ \frac{1}{2n}, & \frac{n}{2} < i \leq n \end{cases}$$

Average-case Time Complexity (Problem 1.8)

Input : $r \in [1, n]$, $r \in \mathbb{Z}^+$

$$P\{r = i\} = \begin{cases} \frac{1}{n}, & 1 \leq i \leq \frac{n}{4} \\ \frac{2}{n}, & \frac{n}{4} < i \leq \frac{n}{2} \\ \frac{1}{2n}, & \frac{n}{2} < i \leq n \end{cases}$$

$$\begin{aligned} A &= \sum_{X \in \mathcal{X}} T(X) \cdot P(X) \\ &= T(1)P(1) + T(2)P(2) + \cdots + T(n)P(n) \\ &= \frac{n}{4} \times 10 \times \frac{1}{n} + \frac{n}{4} \times 20 \times \frac{2}{n} + \frac{n}{4} \times 30 \times \frac{1}{2n} + \frac{n}{4} \times n \times \frac{1}{2n} \\ &= \frac{1}{8}n + \frac{65}{4} \end{aligned}$$

Average-case Analysis of Quicksort

$$A(n) = n - 1 + \frac{1}{n} \sum_{i=0}^{i=n-1} (A(i) + A(n - i - 1))$$

$$A(n) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)] = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$

Average-case Analysis of Quicksort

$$A(n) = n - 1 + \frac{1}{n} \sum_{i=0}^{i=n-1} (A(i) + A(n - i - 1))$$

$$A(n) = \mathbb{E}_{X \in \mathcal{X}_n} [T(X)] = \sum_{X \in \mathcal{X}_n} T(X) \cdot P(X)$$


$$\begin{aligned} A(n) &= \mathbb{E}[T(X)] \\ &= \mathbb{E}[\mathbb{E}[T(X)|I]] \\ &= \sum_{i=0}^{i=n-1} P(I = i) \mathbb{E}[T(X) \mid I = i] \\ &= \sum_{i=0}^{i=n-1} \frac{1}{n} [n - 1 + A(i) + A(n - i - 1)] \end{aligned}$$

3-element Sorting (Problem 1.1)

- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.


3-element Sorting (Problem 1.1)

- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) =$$

3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3$$

3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) =$$

3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2$$

3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2 \quad A(3) =$$

3-element Sorting (Problem 1.1)


- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{1}{6}(3 + 3 + 2 + 3 + 3 + 2) = \frac{8}{3}$$

3-element Sorting (Problem 1.1)

- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.




$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{1}{6}(3 + 3 + 2 + 3 + 3 + 2) = \frac{8}{3}$$

$$LB(3) =$$

3-element Sorting (Problem 1.1)

- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.




$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{1}{6}(3 + 3 + 2 + 3 + 3 + 2) = \frac{8}{3}$$

$$LB(3) = 3$$

3-element Sorting (Problem 1.1)

- (1) Design an algorithm for **sorting** 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{1}{6}(3 + 3 + 2 + 3 + 3 + 2) = \frac{8}{3}$$


$$LB(3) = 3 \quad (LB(3) \geq \log 3!)$$

3-element Median Selection (Problem 1.2)

- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.


3-element Median Selection (Problem 1.2)

- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



3-element Median Selection (Problem 1.2)


- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{8}{3}$$

3-element Median Selection (Problem 1.2)

- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.




$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{8}{3}$$

$$LB(3) =$$

3-element Median Selection (Problem 1.2)

- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.




$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{8}{3}$$

$$LB(3) = 3$$

3-element Median Selection (Problem 1.2)

- (1) Design an algorithm for **selecting the median** of 3 distinct elements.
- (2) Worst-case and average-case time complexity.
- (3) Worst-case lower bound.



$$W(3) = 3 \quad B(3) = 2 \quad A(3) = \frac{8}{3}$$
$$LB(3) = 3 \quad (LB(3) \geq \frac{3n}{2} - \frac{3}{2})$$



$$LB = 2$$



LB = 2

```
1: procedure MEDIAN( $a, b, c$ )
2:   if  $(a - b)(a - c) < 0$  then
3:     return  $a$ 
4:   if  $(b - a)(b - c) < 0$  then
5:     return  $b$ 
6:   return  $c$ 
```



LB = 2

```
1: procedure MEDIAN( $a, b, c$ )
2:   if  $(a - b)(a - c) < 0$  then
3:     return  $a$ 
4:   if  $(b - a)(b - c) < 0$  then
5:     return  $b$ 
6:   return  $c$ 
```

Not comparison-based!

Exercise

$$n = 5$$

Exercise

$$n = 5$$

Reference

“The Art of Computer Programming, Vol 3: Sorting and Searching (Section 5.3.1)” by Donald E. Knuth

$$S(21) = 66$$

Mathematical Induction



Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```

Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```

Loop invariant (after the k -th loop):

Horner's rule (Problem 1.5)

$$P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n$$

```
1: procedure HORNER( $A[0 \dots n], x$ )                                ▷  $A : \{a_0 \dots a_n\}$ 
2:    $p \leftarrow A[n]$ 
3:   for  $i \leftarrow n - 1$  downto 0 do
4:      $p \leftarrow px + A[i]$ 
5:   return  $p$ 
```

Loop invariant (after the k -th loop):

$$\mathcal{I} : p = \sum_{i=n-k}^{i=n} a_i x^{k-(n-i)}$$

$$\mathcal{I}: p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$



$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$



When you are in an exam:

20% : Finding \mathcal{I}

80% : Proving \mathcal{I} by PMI

$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$

Proof.

Prove by mathematical induction on non-negative integer k ,
the number of loops.

$$\mathcal{I} : p = \sum_{i=n}^{i=n-k} a_i x^{k-(n-i)}$$

Proof.

Prove by mathematical induction on non-negative integer k ,
the number of loops.

Basis:

$$k = 0 : p = a_n = \mathcal{I}_0$$

Inductive Hypothesis:

Inductive Step:



Integer Multiplication (Problem 1.6)

```
1: procedure INT-MULT( $y, z$ )  
2:   if  $z = 0$  then  
3:     return 0  
4:   return INT-MULT( $cy, \lfloor \frac{z}{c} \rfloor$ ) +  $y(z \bmod c)$ 
```

Integer Multiplication (Problem 1.6)


```
1: procedure INT-MULT( $y, z$ )
2:   if  $z = 0$  then
3:     return 0
4:   return INT-MULT( $cy, \lfloor \frac{z}{c} \rfloor$ ) +  $y(z \bmod c)$ 
```

Proof.

Prove by mathematical induction on non-negative integer z .




2-tree; full binary tree (Problem 2.5)



$$n_0 = n_2 + 1$$

Proof.

2-tree; full binary tree (Problem 2.5)




$$n_0 = n_2 + 1$$

Proof.


Prove by mathematical induction on *the structure of binary tree*.



Asymptotics



Asymptotics



$Q: \theta(f)?$

$$O(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\Omega(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$O(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\Omega(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$\Theta(g(n)) = \{f(n) \mid \exists c_1 > 0, \exists c_2 > 0, \exists n_0, \forall n \geq n_0 : \\ 0 \leq c_1 g(n) \leq f(n) \leq c_2 g(n)\}$$

$$O(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\Omega(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$\Theta(g(n)) = \{f(n) \mid \exists c_1 > 0, \exists c_2 > 0, \exists n_0, \forall n \geq n_0 : \\ 0 \leq c_1 g(n) \leq f(n) \leq c_2 g(n)\}$$

$$o(g(n)) = \{f(n) \mid \forall c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\omega(g(n)) = \{f(n) \mid \forall c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$O(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\Omega(g(n)) = \{f(n) \mid \exists c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$\Theta(g(n)) = \{f(n) \mid \exists c_1 > 0, \exists c_2 > 0, \exists n_0, \forall n \geq n_0 : \\ 0 \leq c_1 g(n) \leq f(n) \leq c_2 g(n)\}$$

$$o(g(n)) = \{f(n) \mid \forall c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq f(n) \leq cg(n)\}$$

$$\omega(g(n)) = \{f(n) \mid \forall c > 0, \exists n_0, \forall n \geq n_0 : 0 \leq cg(n) \leq f(n)\}$$

$$f(n) \sim g(n) \iff \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$

Asymptotics (Problem 2.6 (4))

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n))$$

Asymptotics (Problem 2.6 (4))

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n))$$

Asymptotics (Problem 2.6 (6))

$$\Theta(g(n)) \cap o(g(n)) = \emptyset$$

Asymptotics (Problem 2.6 (4))

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n))$$

Asymptotics (Problem 2.6 (6))

$$\Theta(g(n)) \cap o(g(n)) = \emptyset$$

$$Q: f(n) = O(g(n)) \vee g(n) = \Omega(f(n))?$$

Asymptotics (Problem 2.6 (4))

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n))$$

Asymptotics (Problem 2.6 (6))

$$\Theta(g(n)) \cap o(g(n)) = \emptyset$$

$$Q: f(n) = O(g(n)) \vee g(n) = \Omega(f(n))?$$

$$f(n) = n, \quad g(n) = n^{1+\sin n}$$

Asymptotics (Problem 2.6 (4))

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n))$$

Asymptotics (Problem 2.6 (6))

$$\Theta(g(n)) \cap o(g(n)) = \emptyset$$

$Q : f(n) = O(g(n)) \vee g(n) = \Omega(f(n))?$

$$f(n) = n, \quad g(n) = n^{1+\sin n}$$

Reference:

“Big Omicron and Big Omega and Big Theta” by Donald E. Knuth, 1976.

Asymptotics (Problem 2.7 (2))

$$(\log n)^2 \text{ vs. } \sqrt{n}$$

Asymptotics (Problem 2.7 (2))

$$(\log n)^2 \text{ vs. } \sqrt{n}$$

$$(\log n)^{c_1} = O(n^{c_2}) \quad c_1, c_2 > 0$$

Asymptotics (Problem 2.10)

$$\log(n!) = \Theta(n \log n)$$

Asymptotics (Problem 2.10)

$$\log(n!) = \Theta(n \log n)$$

Stirling Formula (by *James Stirling*):

$$n! = \Theta\left(\sqrt{2\pi n} \left(\frac{n}{e}\right)^n\right)$$



Asymptotics (Problem 2.10)

$$\log(n!) = \Theta(n \log n)$$

Stirling Formula (by *James Stirling*):

$$n! = \Theta\left(\sqrt{2\pi n} \left(\frac{n}{e}\right)^n\right)$$



$$\log(n!) = \log 1 + \log 2 + \cdots + \log n$$

Asymptotics (Problem 2.10)

$$\log(n!) = \Theta(n \log n)$$

Stirling Formula (by *James Stirling*):

$$n! = \Theta\left(\sqrt{2\pi n} \left(\frac{n}{e}\right)^n\right)$$



$$\log(n!) = \log 1 + \log 2 + \cdots + \log n$$

$$\log(n!) \leq n \log n$$

Asymptotics (Problem 2.10)

$$\log(n!) = \Theta(n \log n)$$

Stirling Formula (by *James Stirling*):

$$n! = \Theta\left(\sqrt{2\pi n} \left(\frac{n}{e}\right)^n\right)$$



$$\log(n!) = \log 1 + \log 2 + \cdots + \log n$$

$$\log(n!) \leq n \log n \qquad \log(n!) \geq \frac{n}{2} \log \frac{n}{2}$$

Summation (Problem 2.20)

```
1: procedure CONUNDRUM( $n$ )
2:    $r \leftarrow 0$ 
3:   for  $i \leftarrow 1$  to  $n$  do
4:     for  $j \leftarrow i + 1$  to  $n$  do
5:       for  $k \leftarrow i + j - 1$  to  $n$  do
6:          $r \leftarrow r + 1$ 
7:   return  $r$ 
```

Summation (Problem 2.20)

```
1: procedure CONUNDRUM( $n$ )
2:    $r \leftarrow 0$ 
3:   for  $i \leftarrow 1$  to  $n$  do
4:     for  $j \leftarrow i + 1$  to  $n$  do
5:       for  $k \leftarrow i + j - 1$  to  $n$  do
6:          $r \leftarrow r + 1$ 
7:   return  $r$ 
```


$$\sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=i+j-1}^n 1 =$$

Summation (Problem 2.20)

```
1: procedure CONUNDRUM( $n$ )
2:    $r \leftarrow 0$ 
3:   for  $i \leftarrow 1$  to  $n$  do
4:     for  $j \leftarrow i + 1$  to  $n$  do
5:       for  $k \leftarrow i + j - 1$  to  $n$  do
6:          $r \leftarrow r + 1$ 
7:   return  $r$ 
```

$$\sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=i+j-1}^n 1 = \frac{n^2 - n}{2} = \Theta(n^2)$$

Recurrences



$$T(n) = aT(n/b) + f(n) \quad (a > 0, b > 1)$$

$$T(n) = aT(n/b) + f(n) \quad (a > 0, b > 1)$$

Assume that $T(n)$ is constant for sufficiently small n .

$$T(n) = aT(n/b) + f(n) \quad (a > 0, b > 1)$$

Assume that $T(n)$ is constant for sufficiently small n .

$$a^{\log_b n} f(c) = \Theta \left(n^{\log_b a} \begin{array}{l} f(n) \\ af(\frac{n}{b}) \\ a^2 f(\frac{n}{b^2}) \\ \vdots \end{array} \right)$$

$$T(n) = aT(n/b) + f(n) \quad (a > 0, b > 1)$$

Assume that $T(n)$ is constant for sufficiently small n .

$$\left. \begin{array}{c} f(n) \\ af(\frac{n}{b}) \\ a^2 f(\frac{n}{b^2}) \\ \vdots \\ a^{\log_b n} f(c) = \Theta(n^{\log_b a}) \end{array} \right\} \sum f(n) \stackrel{\text{vs.}}{=} n^E \left\{ \begin{array}{ll} n^{\log_b a} & f(n) = O(n^{E-\epsilon}) \\ n^{\log_b a} \log n & f(n) = \Theta(n^E) \\ f(n) & f(n) = \Omega(n^{E+\epsilon}) \end{array} \right.$$

Solving Recurrences (Problem 2.15)

- (1) $\Theta(n^{\log_3 2})$
- (2) $\Theta(\log^2 n)$
- (3) $\Theta(n)$
- (4) $\Theta(n \log n)$
- (5) $\Theta(n \log^2 n)$
- (6) $\Theta(n^2)$
- (7) $\Theta(n^{\frac{3}{2}} \log n)$
- (8) $\Theta(n)$
- (9) $\Theta(n^{c+1})$
- (10) $\Theta(c^{n+1})$
- (11) \dots

$$T(n) = T(n/2) + \log n$$

$$T(n) = 2T(n/2) + n \log n$$

Solving Recurrences (Problem 2.15)

- (1) $\Theta(n^{\log_3 2})$
- (2) $\Theta(\log^2 n)$
- (3) $\Theta(n)$
- (4) $\Theta(n \log n)$
- (5) $\Theta(n \log^2 n)$
- (6) $\Theta(n^2)$
- (7) $\Theta(n^{\frac{3}{2}} \log n)$
- (8) $\Theta(n)$
- (9) $\Theta(n^{c+1})$
- (10) $\Theta(c^{n+1})$
- (11) \dots

$$T(n) = T(n/2) + \log n$$

$$T(n) = 2T(n/2) + n \log n$$

Reference:

$$f(n) = \Theta(n^{\log_b a} \log^k n) \implies \Theta(n^{\log_b a} \log^{k+1} n)$$

Solving Recurrences (Problem 2.15)

- (1) $\Theta(n^{\log_3 2})$
- (2) $\Theta(\log^2 n)$
- (3) $\Theta(n)$
- (4) $\Theta(n \log n)$
- (5) $\Theta(n \log^2 n)$
- (6) $\Theta(n^2)$
- (7) $\Theta(n^{\frac{3}{2}} \log n)$
- (8) $\Theta(n)$
- (9) $\Theta(n^{c+1})$
- (10) $\Theta(c^{n+1})$
- (11) \dots

$$T(n) = T(n/2) + \log n$$

$$T(n) = 2T(n/2) + n \log n$$

Reference:

$$f(n) = \Theta(n^{\log_b a} \log^k n) \implies \Theta(n^{\log_b a} \log^{k+1} n)$$

$$T(n) = T(n-1) + n^c \quad c \geq 1$$

$$T(n) = T(n-1) + c^n \quad c > 1$$

Solving Recurrences (Problem 2.15 (11))

$$T(n) = T(n/2) + T(n/4) + T(n/8)$$

Solving Recurrences (Problem 2.15 (11))

$$T(n) = T(n/2) + T(n/4) + T(n/8) + n$$

Solving Recurrences (Problem 2.15 (11))

$$T(n) = T(n/2) + T(n/4) + T(n/8) + n$$

By recursion-tree.

$$T(n) = \Theta(n)$$

Solving Recurrences (Problem 2.15 (11))

$$T(n) = T(n/2) + T(n/4) + T(n/8) + n$$

By recursion-tree.

$$T(n) = \Theta(n)$$

Exercise: Prove it by mathematical induction.

Solving Recurrences (Problem 2.15 (11))

$$T(n) = T(n/2) + T(n/4) + T(n/8) + n$$

By recursion-tree.

$$T(n) = \Theta(n)$$

Exercise: Prove it by mathematical induction.

Reference:

“On the Solution of Linear Recurrence Equations” by Akra & Bazzi, 1996.

$$T(n) = \sum_{i=1}^k a_i T(n/b_i) + f(n)$$

Solving Recurrences (Problem 2.17)

$$\begin{aligned}T(n) &= \sqrt{n} \, T(\sqrt{n}) + n \\&= n^{\frac{1}{2}} \, T\left(n^{\frac{1}{2}}\right) + n \\&= n^{\frac{1}{2}} \left(n^{\frac{1}{2^2}} \, T\left(n^{\frac{1}{2^2}}\right) + n^{\frac{1}{2}} \right) + n \\&= n^{\frac{1}{2} + \frac{1}{2^2}} \, T\left(n^{\frac{1}{2^2}}\right) + 2n \\&= n^{\frac{1}{2} + \frac{1}{2^2}} \left(n^{\frac{1}{2^3}} \, T\left(n^{\frac{1}{2^3}}\right) + n^{\frac{1}{2^2}} \right) + 2n \\&= n^{\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3}} \, T\left(n^{\frac{1}{2^3}}\right) + 3n \\&= \dots \\&= n^{\sum_{i=1}^k \frac{1}{2^i}} \, T\left(n^{\frac{1}{2^k}}\right) + kn\end{aligned}$$

$$n^{\frac{1}{2^k}} = 2$$

$$n^{\frac{1}{2^k}} = 2 \implies k = \log \log n$$

$$n^{\frac{1}{2^k}} = \textcolor{red}{2} \implies k = \log \log n$$

$$\begin{aligned} T(n) &= n \sum_{i=1}^k \frac{1}{2^i} T\left(n^{\frac{1}{2^k}}\right) + kn \\ &= n \sum_{i=1}^{\log \log n} \frac{1}{2^i} T(2) + n \log \log n \end{aligned}$$

$$n^{\frac{1}{2^k}} = 2 \implies k = \log \log n$$

$$\begin{aligned} T(n) &= n \sum_{i=1}^k \frac{1}{2^i} T\left(n^{\frac{1}{2^i}}\right) + kn \\ &= n \sum_{i=1}^{\log \log n} \frac{1}{2^i} T(2) + n \log \log n \end{aligned}$$

$$\sum_{i=1}^{\log_2 \log_2(n)} \frac{1}{2^i} < 1 \implies T(n) = \Theta(n \log \log n)$$

$$n^{\frac{1}{2^k}} = 2 \implies k = \log \log n$$

$$\begin{aligned} T(n) &= n \sum_{i=1}^k \frac{1}{2^i} T\left(n^{\frac{1}{2^i}}\right) + kn \\ &= n \sum_{i=1}^{\log \log n} \frac{1}{2^i} T(2) + n \log \log n \end{aligned}$$

$$\sum_{i=1}^{\log_2 \log_2(n)} \frac{1}{2^i} < 1 \implies T(n) = \Theta(n \log \log n)$$

Exercise: Prove it by mathematical induction.

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$\frac{T(2^m)}{2^m} = \frac{T(2^{m/2})}{2^{m/2}} + 1$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$\frac{T(2^m)}{2^m} = \frac{T(2^{m/2})}{2^{m/2}} + 1$$

$$S(m) \leftrightarrow \frac{T(2^m)}{2^m}$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$\frac{T(2^m)}{2^m} = \frac{T(2^{m/2})}{2^{m/2}} + 1$$

$$S(m) \leftrightarrow \frac{T(2^m)}{2^m}$$

$$S(m) = S(m/2) + 1 =$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$\frac{T(2^m)}{2^m} = \frac{T(2^{m/2})}{2^{m/2}} + 1$$

$$S(m) \leftrightarrow \frac{T(2^m)}{2^m}$$

$$S(m) = S(m/2) + 1 = \Theta(\log m)$$

$$T(n) = \sqrt{n}T(\sqrt{n}) + n$$

$$\frac{T(n)}{n} = \frac{T(\sqrt{n})}{\sqrt{n}} + 1$$

$$n \leftrightarrow 2^m$$

$$\frac{T(2^m)}{2^m} = \frac{T(2^{m/2})}{2^{m/2}} + 1$$

$$S(m) \leftrightarrow \frac{T(2^m)}{2^m}$$

$$S(m) = S(m/2) + 1 = \Theta(\log m)$$

$$T(n) = n \log \log n$$