

(十二) 图论: 匹配与网络流 (Matching and Network Flow)

魏恒峰

hfwei@nju.edu.cn

2021 年 05 月 27 日






3 Theorems + 1 Algorithm

Definition (Network (网络))


A **network** is a **digraph** with

- ▶ a distinguished **source vertex** s ,
- ▶ a distinguished **sink vertex** t ,
- ▶ a **capacity** $c(e) \geq 0$ on each edge e




Definition (Flow (流))

A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



Definition (Flow (流))


A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



$$f^+(v) = \sum_{vw \in E} f(vw)$$

Definition (Flow (流))

A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



$$f^+(v) = \sum_{vw \in E} f(vw) \quad f^-(v) = \sum_{uv \in E} f(uv)$$

Definition (Feasible)


A flow f is **feasible** if it satisfies

Capacity Constraints:

$$\forall e \in E(G). 0 \leq f(e) \leq c(e)$$

Conservation Constraints:


$$\forall v \in V(G) - \{s, t\}. f^+(v) = f^-(v)$$



Definition (Value (值))

The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$



Definition (Value (值))

The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$



Definition (Value (值))



The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$





Definition (Maximum Flow (最大流))

A **maximum flow** is a **feasible flow** of maximum **value**.





$$s \rightarrow x \rightarrow v \rightarrow t$$



$$s \rightarrow x \rightarrow v \rightarrow t$$


Definition (f -augmenting Paths (增广路径))


$$\min_{e \in E(P)} \epsilon(e)$$

Definition (Source/Sink Cut (割))

In a network, a **source/sink cut** $[S, T]$ consists of the edges **from** a **source set** S **to** a **sink set** T , where


$$V = S \uplus T \wedge s \in S \wedge t \in T$$





Definition (Capacity of Cut (割的容量))

$$\text{cap}(S, T) = \sum_{u \in S, v \in T, uv \in E} c(uv)$$




Definition (Minimum Cut (最小割))

A **minimum cut** is a **cut** of minimum value.

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.


$$\text{val}(f) \leq \text{cap}(S, T).$$



Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S)$$

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S)$$

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.


$$\text{val}(f) \leq \text{cap}(S, T).$$



$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S) \leq \text{cap}(S, T)$$

Lemma

$$\max_f \text{val}(f) \leq \min_{[S,T]} \text{cap}(S, T)$$



What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?


$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^{-1}(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^{-1}(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$



Theorem (Max-flow Min-cut Theorem (Ford and Fulkerson; 1956))

$$\max_f \text{val}(f) = \min_{[S,T]} \text{cap}(S,T)$$


(Strong Duality)



L. R. Ford Jr. (1927 ~ 2017)



D. R. Fulkerson (1924 ~ 1976)



Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

We construct a cut $[S, T]$ with $\text{val}(f) = \text{cap}(S, T)$.

Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

We construct a cut $[S, T]$ with $\text{val}(f) = \text{cap}(S, T)$.

$$S \triangleq$$

Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

We construct a cut $[S, T]$ with $\text{val}(f) = \text{cap}(S, T)$.

$$S \triangleq$$

$$T \triangleq V - S$$

Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

We construct a cut $[S, T]$ with $\text{val}(f) = \text{cap}(S, T)$.

$$S \triangleq$$

$$T \triangleq V - S$$

$$f^{-1}(S) = 0 \wedge f^{+}(S) = \text{cap}(S, T)$$

Theorem

A feasible flow f is maximum iff there are no f -augmenting paths.

We construct a cut $[S, T]$ with $\text{val}(f) = \text{cap}(S, T)$.

$$S \triangleq$$

$$T \triangleq V - S$$

$$f^{-1}(S) = 0 \wedge f^{+}(S) = \text{cap}(S, T)$$


$$\text{val}(f) = f^{+}(S) - f^{-}(S) = f^{+}(S) = \text{cap}(S, T)$$

The Ford-Fulkerson Method

Repeatedly finding f -augmenting paths until no more ones exist.


The Ford-Fulkerson Method

Repeatedly finding f -augmenting paths until no more ones exist.



The Ford-Fulkerson Method

Repeatedly finding f -augmenting paths until no more ones exist.




The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find f -augmenting paths.


The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find f -augmenting paths.



The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find f -augmenting paths.



Theorem (Hall Theorem; 1935)


There is an *X -perfect matching* of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

Theorem (Hall Theorem; 1935)

There is an *X*-perfect matching of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$




$$\forall x \in X. c(s, x) = 1 \quad \forall y \in Y. c(y, t) = 1 \quad \forall x \in X, y \in Y. c(x, y) = \infty$$

Theorem (Hall Theorem; 1935)

There is an X -perfect matching of G iff


$$\forall W \subseteq X. |W| \leq |N_G(W)|$$




$$\forall x \in X. c(s, x) = 1 \quad \forall y \in Y. c(y, t) = 1 \quad \forall x \in X, y \in Y. c(x, y) = \infty$$

We need to show that $\max_f \text{val}(f) = |X|$.

We need to show that $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$.




We need to show that $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$.



$$\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \leq |X|$$


We need to show that $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$.




$$\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \leq |X|$$

Therefore, we need to show that $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \geq |X|$.

Let $[S, \bar{S}]$ be a minimum cut. We need to show that $\text{cap}(S, \bar{S}) = |X|$.




Let $[S, \bar{S}]$ be a minimum cut. We need to show that $\text{cap}(S, \bar{S}) = |X|$.





$$N(S \cap X) \subseteq (S \cap Y)$$

Let $[S, \bar{S}]$ be a minimum cut. We need to show that $\text{cap}(S, \bar{S}) = |X|$.




$$N(S \cap X) \subseteq (S \cap Y)$$







$$\text{cap}(S, \bar{S}) = \sum_{u \in S, v \in \bar{S}} c(u, v)$$



$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(s, v) + \sum_{u \in S \cap Y} c(u, t)
 \end{aligned}$$



$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(\textcolor{red}{s}, v) + \sum_{u \in S \cap Y} c(u, \textcolor{red}{t}) \\
 &= |X| - |S \cap X| + |S \cap Y| \\
 &\geq |X| - |S \cap X| + \textcolor{red}{|N(S \cap X)|}
 \end{aligned}$$



$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(\textcolor{red}{s}, v) + \sum_{u \in S \cap Y} c(u, \textcolor{red}{t}) \\
 &= |X| - |S \cap X| + |S \cap Y| \\
 &\geq |X| - |S \cap X| + \textcolor{red}{|N(S \cap X)|} \\
 &\geq |X|
 \end{aligned}$$

Theorem (König (1931), Egerváry (1931))


If G is a bipartite graph, then the maximum size of a matching in G equals the minimum size of a vertex cover of G

Theorem (König (1931), Egerváry (1931))

If G is a bipartite graph, then the maximum size of a matching in G equals the minimum size of a vertex cover of G



Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn