

(十三) 群论: 群的基本概念 (What are Groups?)

魏恒峰

hfwei@nju.edu.cn

2021 年 06 月 03 日



Definition (Group (群))

A **group** $(G, *)$ is a **set** G together with a **binary operation** $*$ such that the following four **group axioms** are satisfied:

Definition (Group (群))

A **group** $(G, *)$ is a **set** G together with a **binary operation** $*$ such that the following four **group axioms** are satisfied:

Closure (封闭):

$$\forall a, b \in G. a * b \in G$$

Definition (Group (群))

A **group** $(G, *)$ is a **set** G together with a **binary operation** $*$ such that the following four **group axioms** are satisfied:

Closure (封闭):

$$\forall a, b \in G. a * b \in G$$

Associativity (结合律):

$$\forall a, b, c \in G. (a * b) * c = a * (b * c)$$

Definition (Group (群))

A **group** $(G, *)$ is a **set** G together with a **binary operation** $*$ such that the following four **group axioms** are satisfied:

Closure (封闭):

$$\forall a, b \in G. a * b \in G$$

Associativity (结合律):

$$\forall a, b, c \in G. (a * b) * c = a * (b * c)$$

Identity (单位元):

$$\exists e \in G. \forall a \in G. e * a = a * e = a$$

Definition (Group (群))

A **group** $(G, *)$ is a **set** G together with a **binary operation** $*$ such that the following four **group axioms** are satisfied:

Closure (封闭):

$$\forall a, b \in G. a * b \in G$$

Associativity (结合律):

$$\forall a, b, c \in G. (a * b) * c = a * (b * c)$$

Identity (单位元):

$$\exists e \in G. \forall a \in G. e * a = a * e = a$$

Inverse (逆元): Let e be **the** identity of G .

$$\forall a \in G. \exists b \in G. a * b = b * a = e$$

The inverse of a is denoted a^{-1} .

Definition (Commutative Group (交换群); Abelian Group (阿贝尔群))

Let $(G, *)$ be a group. If $*$ is commutative,

$$\forall a, b \in G. a * b = b * a,$$

then $(G, *)$ is a commutative group.

$$(\mathbb{Z}, +)$$

$$(\mathbb{Z}, +)$$

$$(\mathbb{Q} \setminus \{0\}, \times)$$

$$(\mathbb{Z}, +)$$

$$(\mathbb{Q} \setminus \{0\}, \times)$$


$$(1, -1, \mathbf{i}, -\mathbf{i})$$

Group of n -th Roots of Unity (n 次单位根群)

$$\begin{aligned} U_n &= \{z \in \mathbb{C} \mid z^n = 1\} \\ &= \left\{ \cos \frac{2k\pi}{n} + \mathbf{i} \sin \frac{2k\pi}{n} \mid k = 0, 1, \dots, n-1 \right\} \end{aligned}$$

Group of n -th Roots of Unity (n 次单位根群)

$$\begin{aligned}U_n &= \{z \in \mathbb{C} \mid z^n = 1\} \\&= \left\{ \cos \frac{2k\pi}{n} + \mathbf{i} \sin \frac{2k\pi}{n} \mid k = 0, 1, \dots, n-1 \right\}\end{aligned}$$



Quaternion Group (四元数群)

$$(1, i, j, k, -1, -i, -j, -k)$$

x	e	\bar{e}	i	\bar{i}	j	\bar{j}	k	\bar{k}
e	e	\bar{e}	i	\bar{i}	j	\bar{j}	k	\bar{k}
\bar{e}	\bar{e}	e	\bar{i}	i	\bar{j}	j	\bar{k}	k
i	i	\bar{i}	\bar{e}	e	k	\bar{k}	\bar{j}	j
\bar{i}	\bar{i}	i	e	\bar{e}	\bar{k}	k	j	\bar{j}
j	j	\bar{j}	\bar{k}	k	\bar{e}	e	i	\bar{i}
\bar{j}	\bar{j}	j	k	\bar{k}	e	\bar{e}	\bar{i}	i
k	k	\bar{k}	j	\bar{j}	\bar{i}	i	\bar{e}	e
\bar{k}	\bar{k}	k	\bar{j}	j	i	\bar{i}	e	\bar{e}



Cayley Table

$$i^2 = j^2 = k^2 = 1 \quad ij = k, ji = -k, jk = i, kj = -i, ki = j, ik = -j$$

Theorem

Let G be a group.

(1) *The identity is unique.*

Theorem

Let G be a group.

- (1) *The identity is unique.*
- (2) *The inverse of each element is unique.*

Theorem

Let G be a group.

- (1) *The identity is unique.*
- (2) *The inverse of each element is unique.*
- (3) $\forall a \in G. (a^{-1})^{-1} = a.$

Theorem

Let G be a group.

- (1) *The identity is unique.*
- (2) *The inverse of each element is unique.*
- (3) $\forall a \in G. (a^{-1})^{-1} = a.$
- (4) $\forall a, b \in G. (ab)^{-1} = b^{-1}a^{-1}.$

Theorem

Let G be a group.

- (1) *The identity is unique.*
- (2) *The inverse of each element is unique.*
- (3) $\forall a \in G. (a^{-1})^{-1} = a.$
- (4) $\forall a, b \in G. (ab)^{-1} = b^{-1}a^{-1}.$
- (5) $\forall a, b, c \in G. (ab = ac \implies b = c) \wedge (ba = ca \implies b = c).$

Theorem

Let G be a group.

- (1) *The identity is unique.*
- (2) *The inverse of each element is unique.*
- (3) $\forall a \in G. (a^{-1})^{-1} = a.$
- (4) $\forall a, b \in G. (ab)^{-1} = b^{-1}a^{-1}.$
- (5) $\forall a, b, c \in G. (ab = ac \implies b = c) \wedge (ba = ca \implies b = c).$
- (6) $\forall a, b \in G. \exists! x \in G. ax = b \wedge ya = b.$

Additive Group of Integers Modulo m (模 m 剩余类加群)

$$(\mathbb{Z}_m = \{0, 1, \dots, m-1\}, +_m)$$

Additive Group of Integers Modulo m (模 m 剩余类加群)

$$(\mathbb{Z}_m = \{0, 1, \dots, m-1\}, +_m)$$

$$(\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}, \times_6)$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Bézout's Identity)

$$(a, b) = d \implies \exists u, v \in \mathbb{Z}. au + bv = d$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Bézout's Identity)

$$(a, b) = d \implies \exists u, v \in \mathbb{Z}. au + bv = d$$

$$(a, m) = 1 \implies \exists u, v \in \mathbb{Z}. au + mv = 1$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Bézout's Identity)

$$(a, b) = d \implies \exists u, v \in \mathbb{Z}. au + bv = d$$

$$(a, m) = 1 \implies \exists u, v \in \mathbb{Z}. au + mv = 1$$

$$a^{-1} = u$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Bézout's Identity)

$$(a, b) = d \implies \exists u, v \in \mathbb{Z}. au + bv = d$$

$$(a, m) = 1 \implies \exists u, v \in \mathbb{Z}. au + mv = 1$$

$$a^{-1} = u$$

$$(u, m) = 1$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Bézout's Identity)

$$(a, b) = d \implies \exists u, v \in \mathbb{Z}. au + bv = d$$

$$(a, m) = 1 \implies \exists u, v \in \mathbb{Z}. au + mv = 1$$

$$a^{-1} = u$$

$$(u, m) = 1 \quad ua = au = au + mv = 1 \pmod{m}$$

When p is a prime,

$$\mathbb{Z}_p^* \triangleq U(p) = \{1, 2, \dots, p-1\}$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$|U(m)| = \varphi(m)$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$|U(m)| = \varphi(m)$$

Definition (Euler's Totient Function (1763))

$$\varphi(m) = m \prod_{p|m \wedge p \text{ is a prime}} \left(1 - \frac{1}{p}\right)$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$|U(m)| = \varphi(m)$$

Definition (Euler's Totient Function (1763))

$$\varphi(m) = n \prod_{p|n \wedge p \text{ is a prime}} \left(1 - \frac{1}{p}\right)$$

$$\varphi(20) = 20\left(1 - \frac{1}{2}\right)\left(1 - \frac{1}{5}\right) = 8$$

Multiplicative Group of Integers Modulo m (模 m 剩余类乘法群)

$$U(m) = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$|U(m)| = \varphi(m)$$

Definition (Euler's Totient Function (1763))

$$\varphi(m) = n \prod_{p|n \wedge p \text{ is a prime}} \left(1 - \frac{1}{p}\right)$$

$$\varphi(20) = 20\left(1 - \frac{1}{2}\right)\left(1 - \frac{1}{5}\right) = 8$$

$$U(20) = \{1, 3, 7, 9, 11, 13, 17, 19\}$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

$$G = \{a_1, a_2, \dots, a_n\}$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

$$G = \{a_1, a_2, \dots, a_n\}$$

$$aG \triangleq \{aa_1, aa_2, \dots, aa_n\} = G$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

$$G = \{a_1, a_2, \dots, a_n\}$$

$$aG \triangleq \{aa_1, aa_2, \dots, aa_n\} = G$$

$$\prod_{i=1}^n (aa_i) = a_1 \dots a_n$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

$$G = \{a_1, a_2, \dots, a_n\}$$

$$aG \triangleq \{aa_1, aa_2, \dots, aa_n\} = G$$

$$\prod_{i=1}^n (aa_i) = a_1 \dots a_n$$

$$a^n a_1 \dots a_n = a_1 \dots a_n$$

Theorem

Let G be an abelian group of *order* n .

$$\forall a \in G. a^n = e.$$

$$G = \{a_1, a_2, \dots, a_n\}$$

$$aG \triangleq \{aa_1, aa_2, \dots, aa_n\} = G$$

$$\prod_{i=1}^n (aa_i) = a_1 \dots a_n$$

$$a^n a_1 \dots a_n = a_1 \dots a_n \implies a^n = e$$

Theorem (Euler Theorem (1736))

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

Theorem (Euler Theorem (1736))

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$7^{222} \pmod{10}$$

Theorem (Euler Theorem (1736))

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$7^{222} \pmod{10}$$

$$(7, 10) = 1 \quad \varphi(10) = 4$$

Theorem (Euler Theorem (1736))

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$7^{222} \pmod{10}$$

$$(7, 10) = 1 \quad \varphi(10) = 4$$

$$7^4 \equiv 1 \pmod{10}$$

Theorem (Euler Theorem (1736))

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$7^{222} \pmod{10}$$

$$(7, 10) = 1 \quad \varphi(10) = 4$$

$$7^4 \equiv 1 \pmod{10}$$

$$7^{222} \equiv 7^{4 \times 55 + 2} \equiv 7^2 \equiv 9 \pmod{10}$$

Theorem (Euler Theorem)

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

Theorem (Euler Theorem)

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$U_m = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

Theorem (Euler Theorem)

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$U_m = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$(a, m) = 1 \implies a \in U_m$$

Theorem (Euler Theorem)

Let $m \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

$$U_m = \{a \in \mathbb{Z}_m \mid (a, m) = 1\}$$

$$(a, m) = 1 \implies a \in U_m$$

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

Theorem (Fermat's Little Theorem (1640))

Let p be a prime. Then for any $a \in \mathbb{Z}^+$,

$$a^{p-1} \equiv 1 \pmod{p}$$

Theorem (Fermat's Little Theorem (1640))

Let p be a prime. Then for any $a \in \mathbb{Z}^+$,

$$a^{p-1} \equiv 1 \pmod{p}$$

$$\varphi(p) = p - 1$$

permutation

Definition (Subgroup (子群))

Let $(G, *)$ be a group and $\emptyset \neq H \subseteq G$.

If $(H, *)$ is a group, then we call H a **subgroup** of G , denoted $H \leq G$.

Definition (Subgroup (子群))

Let $(G, *)$ be a group and $\emptyset \neq H \subseteq G$.

If $(H, *)$ is a group, then we call H a **subgroup** of G , denoted $H \leq G$.

$H = G, H = \{e\}$ are two **trivial** (平凡) subgroups.

Definition (Subgroup (子群))

Let $(G, *)$ be a group and $\emptyset \neq H \subseteq G$.

If $(H, *)$ is a group, then we call H a **subgroup** of G , denoted $H \leq G$.

$H = G, H = \{e\}$ are two **trivial** (平凡) subgroups.

If $H \subset G$, then H is a **proper** subgroup (真子群).

$$(H = \{mz \mid z \in \mathbb{Z}\}, +) \leq (\mathbb{Z}, +)$$

$$(H = \{mz \mid z \in \mathbb{Z}\}, +) \leq (\mathbb{Z}, +)$$

$$H = \{1, 2, 4\} \leq G = \mathbb{Z}_7^* = \{1, 2, 3, 4, 5, 6\}$$

Theorem

Suppose that $H \leq G$.

(1) The identity of H is the same with that of G .

$$e_H = e_G$$

(2) The inversion of a in H is the same with that in G .

$$\forall a \in H. a_H^{-1} = a_G^{-1}$$

Theorem

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. ab^{-1} \in H.$$

Theorem

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

Theorem

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

$$a^{-1} = ea^{-1} \in H$$

Theorem

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

$$a^{-1} = ea^{-1} \in H$$

$$ab = a(b^{-1})^{-1} \in H$$

Theorem

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G.$$

Theorem

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G.$$

$$H_1 = 2\mathbb{Z} \leq \mathbb{Z} \quad H_2 = 3\mathbb{Z} \leq \mathbb{Z}$$

Theorem

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G.$$

$$H_1 = 2\mathbb{Z} \leq \mathbb{Z} \quad H_2 = 3\mathbb{Z} \leq \mathbb{Z}$$

$$H_1 \cap H_2 = 6\mathbb{Z} \leq \mathbb{Z}$$

Theorem

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G.$$

$$H_1 = 2\mathbb{Z} \leq \mathbb{Z} \quad H_2 = 3\mathbb{Z} \leq \mathbb{Z}$$

$$H_1 \cap H_2 = 6\mathbb{Z} \leq \mathbb{Z}$$

$$H_1 \cup H_2?$$

Center (中心)

Let G be a group. Let

$$C(G) \triangleq \{g \in G \mid gx = xg, \forall x \in G\}.$$

Then $C(G) \leq G$.

Definition (Isomorphism (同构))

Let (G, \cdot) and $(G', *)$ be two groups. Let ϕ be a **bijection** such that

$$\forall a, b \in G. \phi(a \cdot b) = \phi(a) * \phi(b).$$

Then ϕ is an **isomorphism** from G to G' .

Definition (Isomorphism (同构))

Let (G, \cdot) and $(G', *)$ be two groups. Let ϕ be a **bijection** such that

$$\forall a, b \in G. \phi(a \cdot b) = \phi(a) * \phi(b).$$

Then ϕ is an **isomorphism** from G to G' .

G and G' are isomorphic

$$\phi : G \cong G'$$

$$(\mathbb{R}, +) \cong (\mathbb{R}^+, *)$$

$$(\mathbb{R}, +) \cong (\mathbb{R}^+, *)$$

$$\phi(x) = e^x$$

Theorem

Suppose that $\phi : G \cong G'$. Let e and e' be identities of G and G' , respectively.

$$(1) \quad \phi(e) = e'$$

$$(2) \quad \phi(a^{-1}) = (\phi(a))^{-1}$$

$$(3) \quad \phi^{-1} : G' \cong G$$

Klein Four-group (四元群; $K_4; V$)

*	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

$$a^2 = b^2 = c^2 = (ab)^2 = e$$

$$ab = c = ba \quad ac = b = ca \quad bc = a = cb$$

Klein Four-group (四元群; $K_4; V$)

*	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

$$a^2 = b^2 = c^2 = (ab)^2 = e$$

$$ab = c = ba \quad ac = b = ca \quad bc = a = cb$$

$$U(8) = \{1, 3, 5, 7\}$$

permutation?

Definition (Order of Elements (元素的阶))

Let G be a group, e be the identity of G .

The **order** of e is the **smallest** positive integer r such that $a^r = e$.

$$\text{ord } a = r$$

Definition (Order of Elements (元素的阶))

Let G be a group, e be the identity of G .

The **order** of e is the **smallest** positive integer r such that $a^r = e$.

$$\text{ord } a = r$$

If such r does not exist, then $\text{ord } a = \infty$.

$$\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}$$

$$\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}$$

$$\mathbb{Z}_5^* = \{1, 2, 3, 4\}$$

$$\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}$$

$$\mathbb{Z}_5^* = \{1, 2, 3, 4\}$$

$$(\mathbb{Z}, +)$$

Theorem

Let G be a group and e be the identity of G .

$$((\text{ord } a = n) \wedge (\exists m \in \mathbb{Z}. a^m = e)) \implies n \mid m.$$

Theorem

Let G be a group and e be the identity of G .

$$((\text{ord } a = n) \wedge (\exists m \in \mathbb{Z}. a^m = e)) \implies n \mid m.$$

$$m = nq + r \quad (0 \leq r < n)$$

Theorem

Let G be a group and e be the identity of G .

$$((\text{ord } a = n) \wedge (\exists m \in \mathbb{Z}. a^m = e)) \implies n \mid m.$$

$$m = nq + r \quad (0 \leq r < n)$$

If $r > 0$,

$$a^r = a^{m-nq} = a^m \cdot (a^n)^{-q} = e \cdot e = e$$

Theorem

Let G be a group and e be the identity of G .

$$((\text{ord } a = n) \wedge (\exists m \in \mathbb{Z}. a^m = e)) \implies n \mid m.$$

$$m = nq + r \quad (0 \leq r < n)$$

If $r > 0$,

$$a^r = a^{m-nq} = a^m \cdot (a^n)^{-q} = e \cdot e = e$$

$$\text{ord } a \neq n$$

Definition (Cyclic Group (循环群))

Let G be a group. If

$$\exists a \in G. G = \langle a \rangle \triangleq \{a^0 = e, a, a^2, a^3, \dots\},$$

then G is a **cyclic group**.

Definition (Cyclic Group (循环群))

Let G be a group. If

$$\exists a \in G. G = \langle a \rangle \triangleq \{a^0 = e, a, a^2, a^3, \dots\},$$

then G is a **cyclic group**.

If $G = \langle a \rangle$, then a is a **generator** (生成元) of G .

$(\mathbb{Z}, +)$ is an **infinite** cyclic group

$(\mathbb{Z}, +)$ is an **infinite** cyclic group

$$(\mathbb{Z}, +) = \langle 1 \rangle = \langle -1 \rangle$$

$(\mathbb{Z}, +)$ is an **infinite** cyclic group

$$(\mathbb{Z}, +) = \langle 1 \rangle = \langle -1 \rangle$$

$(\mathbb{Z}_n, +)$ is a **finite** cyclic group of order n

$(\mathbb{Z}, +)$ is an **infinite** cyclic group

$$(\mathbb{Z}, +) = \langle 1 \rangle = \langle -1 \rangle$$

$(\mathbb{Z}_n, +)$ is a **finite** cyclic group of order n

$$(\mathbb{Z}_m, +) = \langle 1 \rangle$$

$$\mathbb{Z}_5^* = \{1, 2, 3, 4\}$$

$$\mathbb{Z}_5^* = \{1, 2, 3, 4\}$$

$$\mathbb{Z}_5^* = \langle 2 \rangle = \langle 3 \rangle$$

Theorem

(1) Let $G = \{e, a, a^{-1}, a^2, a^{-2}, \dots\}$ be an infinite cyclic group.

$$\forall k, l \in \mathbb{Z}. (a^k = a^l \rightarrow k = l).$$

Theorem

(1) Let $G = \{e, a, a^{-1}, a^2, a^{-2}, \dots\}$ be an infinite cyclic group.

$$\forall k, l \in \mathbb{Z}. (a^k = a^l \rightarrow k = l).$$

(2) Let $G = \{e, a, a^2, \dots, a^{n-1}\}$ be a finite cyclic group of order n .

$$\forall k, l \in \mathbb{Z}. (a^k = a^l \leftrightarrow n \mid (k - l)).$$

Theorem (Structure Theorem of Cyclic Groups (循环群结构定理))

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $G = \langle a \rangle$ is an infinite cyclic group, then $G \cong (\mathbb{Z}, +)$.
- (2) If $G = \langle a \rangle$ is a finite cyclic group of order n , then $G \cong (\mathbb{Z}_n, +)$.

Theorem (Structure Theorem of Cyclic Groups (循环群结构定理))

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $G = \langle a \rangle$ is an infinite cyclic group, then $G \cong (\mathbb{Z}, +)$.
- (2) If $G = \langle a \rangle$ is a finite cyclic group of order n , then $G \cong (\mathbb{Z}_n, +)$.

$$\phi : \mathbb{Z} \rightarrow G$$

$$k \mapsto a^k, \quad \forall k \in \mathbb{Z}$$

Theorem (Structure Theorem of Cyclic Groups (循环群结构定理))

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $G = \langle a \rangle$ is an infinite cyclic group, then $G \cong (\mathbb{Z}, +)$.
- (2) If $G = \langle a \rangle$ is a finite cyclic group of order n , then $G \cong (\mathbb{Z}_n, +)$.

$$\begin{aligned}\phi : \mathbb{Z} &\rightarrow G \\ k &\mapsto a^k, \quad \forall k \in \mathbb{Z}\end{aligned}$$

$$\begin{aligned}\phi : \mathbb{Z}_n &\rightarrow G \\ k &\mapsto a^k, \quad \forall k \in \mathbb{Z}_n\end{aligned}$$

Theorem (Generators of Cyclic Groups)

Let $G = \langle a \rangle$ be a cyclic group.

(1) If $|G| = \infty$, then a and a^{-1} are the only generators of G ;

Theorem (Generators of Cyclic Groups)

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $|G| = \infty$, then a and a^{-1} are the only generators of G ;
- (2) If $|G| = n$, then a^r is a generator of G iff $(r, n) = 1$.

Theorem (Generators of Cyclic Groups)

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $|G| = \infty$, then a and a^{-1} are the only generators of G ;
- (2) If $|G| = n$, then a^r is a generator of G iff $(r, n) = 1$.

Theorem (Generators of Cyclic Groups)

Let $G = \langle a \rangle$ be a cyclic group.

- (1) If $|G| = \infty$, then a and a^{-1} are the only generators of G ;
- (2) If $|G| = n$, then a^r is a generator of G iff $(r, n) = 1$.

$$\text{ord } a^r = \frac{n}{(n, r)}$$


$$(\mathbb{Z}_{12}, +)$$

Generators : 1, 5, 7, 11

Theorem (Subgroups of Cyclic Groups)

Every subgroup of a cyclic group is cyclic.

Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn