(十四) 群论: 子群 (Subgroup)

魏恒峰

hfwei@nju.edu.cn

2021年06月10日



Definition (Subgroup (子群))

Let (G, *) be a group and $\emptyset \neq H \subseteq G$.

If (H, *) is a group, then we call H a subgroup of G, denoted $H \leq G$.

Definition (Subgroup (子群))

Let (G, *) be a group and $\emptyset \neq H \subseteq G$.

If (H, *) is a group, then we call H a subgroup of G, denoted $H \leq G$.

 $H = G, H = \{e\}$ are two trivial (平凡) subgroups.

Definition (Subgroup (子群))

Let (G, *) be a group and $\emptyset \neq H \subseteq G$.

If (H, *) is a group, then we call H a subgroup of G, denoted $H \leq G$.

 $H = G, H = \{e\}$ are two trivial (平凡) subgroups.

If $H \subset G$, then H is a proper subgroup (真子群).

$$(H = \{mz \mid z \in \mathbb{Z}\}, +) \le (\mathbb{Z}, +)$$

$$(H = \{ mz \mid z \in \mathbb{Z} \}, +) \le (\mathbb{Z}, +)$$

$$H = \{1,2,4\} \leq G = \mathbb{Z}_7^* = \{1,2,3,4,5,6\}$$

Suppose that $H \leq G$.

(1) The identity of H is the same with that of G.

$$e_H = e_G$$

(2) The inversion of a in H is the same with that in G.

$$\forall a \in H. \ a_H^{-1} = a_G^{-1}$$

Suppose that $H \leq G$.

(1) The identity of H is the same with that of G.

$$e_H = e_G$$

(2) The inversion of a in H is the same with that in G.

$$\forall a \in H. \ a_H^{-1} = a_G^{-1}$$

$$e_H e_H = e_H = e_H e_G$$

Suppose that $H \leq G$.

(1) The identity of H is the same with that of G.

$$e_H = e_G$$

(2) The inversion of a in H is the same with that in G.

$$\forall a \in H. \ a_H^{-1} = a_G^{-1}$$

$$e_H e_H = e_H = e_H e_G \implies e_H = e_G$$

Suppose that $H \leq G$.

(1) The identity of H is the same with that of G.

$$e_H = e_G$$

(2) The inversion of a in H is the same with that in G.

$$\forall a \in H. \ a_H^{-1} = a_G^{-1}$$

$$e_H e_H = e_H = e_H e_G \implies e_H = e_G$$

$$aa_H^{-1} = e_H = e_G = aa_G^{-1} \implies a_H^{-1} = a^{-1}(G)$$



Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

 $\forall a, b \in H. \ ab^{-1} \in H.$

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. \ ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. \ ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

$$a^{-1} = ea^{-1} \in H$$

Let G be a group and $\emptyset \neq H \subseteq G$. $H \leq G$ iff

$$\forall a, b \in H. \ ab^{-1} \in H.$$

$$e = aa^{-1} \in H$$

$$a^{-1} = ea^{-1} \in H$$

$$ab = a(b^{-1})^{-1} \in H$$

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G$$
.

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G$$
.

$$H_1 = 2\mathbb{Z} \le \mathbb{Z}$$
 $H_2 = 3\mathbb{Z} \le \mathbb{Z}$

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G$$
.

$$H_1 = 2\mathbb{Z} \le \mathbb{Z}$$
 $H_2 = 3\mathbb{Z} \le \mathbb{Z}$

$$H_1 \cap H_2 = 6\mathbb{Z} \leq \mathbb{Z}$$

Suppose that $H_1 \leq G, H_2 \leq G$.

$$H_1 \cap H_2 \leq G$$
.

$$H_1 = 2\mathbb{Z} \le \mathbb{Z}$$
 $H_2 = 3\mathbb{Z} \le \mathbb{Z}$

$$H_1 \cap H_2 = 6\mathbb{Z} \leq \mathbb{Z}$$

$$H_1 \cup H_2$$
?

Definition (Symmetric Group (对称群; Sym(M)))

Let $M \neq \emptyset$ be a set.

All the permutations/bijective functions of M, together with the composition operation, is a group, called the symmetric group of M.

$$M = \{1, 2, \dots, n\}$$
$$S_n \triangleq \operatorname{Sym}(M)$$

 S_3

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} =$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$\tau\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} =$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$\tau \sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$\tau \sigma = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$$

 $\sigma \tau \neq \tau \sigma$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1\ 4)(2\ 3\ 6)(5)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1 \ 4)(2 \ 3 \ 6)(5)$$
$$= (1 \ 4)(2 \ 3 \ 6)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1 \ 4)(2 \ 3 \ 6)(5)$$
$$= (1 \ 4)(2 \ 3 \ 6)$$
$$= (2 \ 3 \ 6)(1 \ 4)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1 \ 4)(2 \ 3 \ 6)(5)$$
$$= (1 \ 4)(2 \ 3 \ 6)$$
$$= (2 \ 3 \ 6)(1 \ 4)$$
$$= (2 \ 3 \ 6)(4 \ 1)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1 \ 4)(2 \ 3 \ 6)(5)$$

$$= (1 \ 4)(2 \ 3 \ 6)$$

$$= (2 \ 3 \ 6)(1 \ 4)$$

$$= (2 \ 3 \ 6)(4 \ 1)$$

$$= (3 \ 6 \ 2)(4 \ 1)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\sigma = (1 \ 4)(2 \ 3 \ 6)(5)$$

$$= (1 \ 4)(2 \ 3 \ 6)$$

$$= (2 \ 3 \ 6)(1 \ 4)$$

$$= (2 \ 3 \ 6)(4 \ 1)$$

$$= (3 \ 6 \ 2)(4 \ 1)$$

$$= (3 \ 6)(6 \ 2)(4 \ 1)$$

 S_3

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

 S_3

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$
$$(1) \quad (12) \quad (13) \quad (23) \quad (123) \quad (132)$$

Definition (Permutation Group (置换群))

Let $M \neq \emptyset$ be a set.

A permutation group of M is a subgroup of Sym(M).

Definition (Permutation Group (置换群))

Let $M \neq \emptyset$ be a set.

A permutation group of M is a subgroup of Sym(M).

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

Definition (Permutation Group (置换群))

Let $M \neq \emptyset$ be a set.

A permutation group of M is a subgroup of Sym(M).

$$S_3 = \{(1), (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$H = \{(1), (1\ 2)\} \le S_3$$

Thank You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn