

(四) 集合: 关系 (Relation)

魏恒峰

hfwei@nju.edu.cn

2021 年 04 月 08 日








Figure 13. A selection of consistency axioms over an execution $(E, \text{repl}, \text{obj}, \text{oper}, \text{rval}, \text{ro}, \text{vis}, \text{ar})$

Auxiliary relations

$\text{sameobj}(e, f) \iff \text{obj}(e) = \text{obj}(f)$

Per-object causality (aka happens-before) order:

$$\text{hbo} = ((\text{ro} \cap \text{sameobj}) \cup \text{vis})^+$$

Causality (aka happens-before) order: $\text{hb} = (\text{ro} \cup \text{vis})^+$

Axioms

EVENTUAL:

$$\forall e \in E. \neg(\exists \text{infinitely many } f \in E. \text{sameobj}(e, f) \wedge \neg(e \xrightarrow{\text{vis}} f))$$

THINAIR: $\text{ro} \cup \text{vis}$ is acyclic

POCV (Per-Object Causal Visibility): $\text{hbo} \subseteq \text{vis}$

POCA (Per-Object Causal Arbitration): $\text{hbo} \subseteq \text{ar}$

COCV (Cross-Object Causal Visibility): $(\text{hb} \cap \text{sameobj}) \subseteq \text{vis}$

COCA (Cross-Object Causal Arbitration): $\text{hb} \cup \text{ar}$ is acyclic

Figure 17. Optimized state-based multi-value register and its simulation

Σ	$= \text{ReplicaID} \times \mathcal{P}(\mathbb{Z} \times (\text{ReplicaID} \rightarrow \mathbb{N}_0))$
\hat{a}_i	$= (r, \hat{a})$
M	$= \mathcal{P}(\mathbb{Z} \times (\text{ReplicaID} \rightarrow \mathbb{N}_0))$
$\text{do}(\text{wr}(a), (r, V), t) =$	$((r, \{(a, \lambda s. \text{if } s \neq r \text{ then } \max\{v(s) (s, v) \in V\} \\ \text{else } \max\{v(s) (s, v) \in V\} + 1\})), \perp)$
$\text{do}(\text{rd}, (r, V), t) =$	$((r, V), \{(a, \lambda s. \perp) \mid s \in V\})$
$\text{send}(\langle r, V \rangle) =$	$((r, V), V)$
$\text{receive}(\langle (r, V), V' \rangle) =$	$(r, \{(a, v) \in V'' \mid$ $v \in \bigcup \{v' \mid \exists a'. (a', v') \in V'' \wedge a \neq a'\}\},$ where $V'' = ((a, \bigcup \{v' \mid (a, v') \in V \cup V'\}) \mid (a, \lambda s. \perp) \in V \cup V'))$
$V[M] \models \langle E, \text{repl}, \text{obj}, \text{oper}, \text{rval}, \text{ro}, \text{vis}, \text{ar} \rangle$	$I \iff (r = a) \wedge (V[M] \models I)$
$V[M] \models \langle E, \text{repl}, \text{obj}, \text{oper}, \text{rval}, \text{ro}, \text{vis}, \text{ar} \rangle, \text{info} \rangle \iff$	$(\forall(a, v), (a', v') \in V. (a = a' \implies v = v')) \wedge$ $(\forall(a, v) \in V. \exists a. v(a) > 0) \wedge$ $(\forall(a, v) \in V. v \in \bigcup \{v' \mid \exists a'. (a', v') \in V' \wedge a \neq a'\}) \wedge$ $\exists \text{distinct } e_{a,k}.$ $\{(e \in E \mid \exists a. \text{oper}^r(e) = \text{wr}(a)) = \{e_{a,k} \mid a \in \text{ReplicaID} \wedge$ $1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V\}\}) \wedge$ $(\forall a, j, k. (\text{repl}^r(e_{a,k}) = a) \wedge (e_{a,j} \xrightarrow{\text{ro}} e_{a,k} \iff j < k)) \wedge$ $(\forall(a, v) \in V. \forall q. \{j \mid \text{oper}^r(e_{a,j}) = \text{wr}(a)\} \cup$ $\{j \mid \exists a, k. e_{a,j} \xrightarrow{\text{vis}} e_{a,k} \wedge \text{oper}^r(e_{a,k}) = \text{wr}(a)\} =$ $\{j \mid 1 \leq j \leq v(q)\}) \wedge$ $(\forall e \in E. (\text{oper}^r(e) = \text{wr}(a)) \wedge$ $\neg \exists f \in E. (\text{oper}^r(f) = \text{wr}(\perp) \wedge e \xrightarrow{\text{ro}} f) \implies (a, \lambda s. \perp) \in V)$

the former. The only non-trivial obligation is to show that if

$$V[M] \models \langle E, \text{repl}, \text{obj}, \text{oper}, \text{rval}, \text{ro}, \text{vis} \rangle, \text{info} \rangle,$$

then

$$\{a \mid (a, \lambda s. \perp) \in V\} \subseteq \{a \mid \exists e \in E. \text{oper}(e) = \text{wr}(a) \wedge$$

 $\neg \exists f \in E. \exists a'. \text{oper}(e) = \text{wr}(a') \wedge e \xrightarrow{\text{ro}} f\} \quad (13)$

(the reverse inclusion is straightforwardly implied by R_{ca}).

Take $(a, \lambda s. \perp) \in V$. We have $(a, v) \in V. \exists a. v(a) > 0$,

$$v \in \bigcup \{v' \mid \exists a'. (a', v') \in V' \wedge a \neq a'\}$$

and

$$\forall(a, v) \in V. \forall q. \{j \mid \text{oper}(e_{a,j}) = \text{wr}(a)\} \cup$$

 $\{j \mid \exists a, k. e_{a,j} \xrightarrow{\text{vis}} e_{a,k} \wedge \text{oper}(e_{a,k}) = \text{wr}(a)\} =$
 $\{j \mid 1 \leq j \leq v(q)\}.$

From this we get that for some $e \in E$

$$\text{oper}(e) = \text{wr}(a) \wedge \neg \exists f \in E. \exists a'. a' \neq a \wedge$$

$$\text{oper}(e) = \text{wr}(a') \wedge e \xrightarrow{\text{ro}} f.$$

Since vis is acyclic, this implies that for some $e' \in E$

$$\text{oper}(e') = \text{wr}(a) \wedge \neg \exists f \in E. \text{oper}(e') = \text{wr}(\perp) \wedge e' \xrightarrow{\text{ro}} f,$$

which establishes (13).

Let us now discharge RECEIVE . Let $\text{receive}(\langle r, V \rangle, V') =$

$$(r, V''), \text{ where } V'' = \{(a, \bigcup \{v' \mid (a, v') \in V \cup V'\}) \mid (a, \lambda s. \perp) \in V \cup V'\};$$

$$V'' = \{(a, v) \in V'' \mid v \in \bigcup \{(a', v') \in V'' \mid a \neq a'\})\}.$$

Assume $(r, V) \models R_{\text{ca}} \mid I, V' \models M \mid J$ and

$$I = ((E, \text{repl}, \text{obj}, \text{oper}, \text{rval}, \text{ro}, \text{vis}, \text{ar}), \text{info});$$

$$J = ((E', \text{repl}', \text{obj}', \text{oper}', \text{rval}', \text{ro}', \text{vis}', \text{ar}'), \text{info}');$$

$$I \cup J = ((E \cup E', \text{repl}^u, \text{obj}^u, \text{oper}^u, \text{rval}^u, \text{ro}^u, \text{vis}^u, \text{ar}^u), \text{info}^u).$$

By agree we have $I \cup J \in \text{IEx}$. Then

$$(\forall(a, v), (a', v') \in V. (a = a' \implies v = v')) \wedge$$

$$(\forall(a, v) \in V. \exists a. v(a) > 0) \wedge$$

$$(\forall(a, v) \in V. v \in \bigcup \{v' \mid \exists a'. (a', v') \in V' \wedge a \neq a'\}) \wedge$$

$$\exists \text{distinct } e_{a,k}.$$

$$\{(e \in E \mid \exists a. \text{oper}^r(e) = \text{wr}(a)) = \{e_{a,k} \mid a \in \text{ReplicaID} \wedge$$

$$1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V\}\}) \wedge$$

$$(\forall a, j, k. (\text{repl}^r(e_{a,k}) = a) \wedge (e_{a,j} \xrightarrow{\text{ro}} e_{a,k} \iff j < k)) \wedge$$

$$(\forall(a, v) \in V. \forall q. \{j \mid \text{oper}^r(e_{a,j}) = \text{wr}(a)\} \cup$$

$$\{j \mid \exists a, k. e_{a,j} \xrightarrow{\text{vis}} e_{a,k} \wedge \text{oper}^r(e_{a,k}) = \text{wr}(a)\} =$$

$$\{j \mid 1 \leq j \leq v(q)\}) \wedge$$

$$(\forall e \in E. (\text{oper}^r(e) = \text{wr}(a)) \wedge$$

$$\neg \exists f \in E. \text{oper}^r(f) = \text{wr}(\perp) \wedge e \xrightarrow{\text{ro}} f) \implies (a, \lambda s. \perp) \in V)$$

and

$$(\forall(a, v), (a', v') \in V'. (a = a' \implies v = v')) \wedge$$

$$(\forall(a, v) \in V'. \exists a. v(a) > 0) \wedge$$

$$(\forall(a, v) \in V'. v \in \bigcup \{v' \mid \exists a'. (a', v') \in V' \wedge a \neq a'\}) \wedge$$

$$\exists \text{distinct } e_{a,k}.$$

$$\{(e \in E' \mid \exists a. \text{oper}^{r'}(e) = \text{wr}(a)) = \{e_{a,k} \mid a \in \text{ReplicaID} \wedge$$

$$1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V'\}\}) \wedge$$

$$(\forall a, j, k. (\text{repl}^{r'}(e_{a,k}) = a) \wedge (e_{a,j} \xrightarrow{\text{ro}'} e_{a,k} \iff j < k)) \wedge$$

$$(\forall(a, v) \in V'. \forall q. \{j \mid \text{oper}^{r'}(e_{a,j}) = \text{wr}(a)\} \cup$$

$$\{j \mid \exists a, k. e_{a,j} \xrightarrow{\text{vis}'} e_{a,k} \wedge \text{oper}^{r'}(e_{a,k}) = \text{wr}(a)\} =$$

$$\{j \mid 1 \leq j \leq v(q)\}) \wedge$$

$$(\forall e \in E'. (\text{oper}^{r'}(e) = \text{wr}(a)) \wedge$$

$$\neg \exists f \in E'. \text{oper}^{r'}(f) = \text{wr}(\perp) \wedge e \xrightarrow{\text{ro}'} f) \implies (a, \lambda s. \perp) \in V').$$

The agree property also implies

$$\forall s, k. 1 \leq k \leq \min \{ \max\{v(s) \mid \exists a. (a, v) \in V\},$$

$$\max\{v(s) \mid \exists a. (a, v) \in V'\} \} \implies e_{a,k} = e'_{a,k}.$$

Hence, there exist distinct

$$e''_{a,k} \text{ for } a \in \text{ReplicaID}, k = 1..(\max\{v(s) \mid \exists a. (a, v) \in V''\}),$$

$$\text{ such that } (\forall s, k. 1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V\} \implies e''_{a,k} = e_{a,k}) \wedge$$

$$(\forall s, k. 1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V'\} \implies e''_{a,k} = e'_{a,k})$$

and

$$(\{(e \in E \cup E' \mid \exists a. \text{oper}^{r,u}(e) = \text{wr}(a)) = \{e''_{a,k} \mid a \in \text{ReplicaID} \wedge 1 \leq k \leq \max\{v(s) \mid \exists a. (a, v) \in V''\}\}) \wedge$$

$$(\forall a, j, k. (\text{repl}^{r,u}(e''_{a,k}) = a) \wedge (e''_{a,j} \xrightarrow{\text{ro}^u} e''_{a,k} \iff j < k)).$$

By the definition of V'' and V'' we have

$$\forall(a, v), (a', v') \in V'', (a = a' \implies v = v').$$

We also straightforwardly get

$$\forall(a, v) \in V'', \exists a. v(a) > 0$$

and

$$(\forall(a, v) \in V'', \forall q. \{j \mid \text{oper}^{r,u}(e''_{a,j}) = \text{wr}(a)\} \cup$$

$$\{j \mid \exists a, k. e''_{a,j} \xrightarrow{\text{vis}^u} e''_{a,k} \wedge \text{oper}^{r,u}(e''_{a,k}) = \text{wr}(a)\} = \{14\})$$

$$\{j \mid 1 \leq j \leq v(q)\}).$$





I'm so excited.



Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Definition (Cartesian Products)

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Definition (Cartesian Products)

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$

Theorem (Ordered Pairs)

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Definition (有序对 (Ordered Pairs))

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Definition (有序对 (Ordered Pairs))

$$(a, b) = (c, d) \iff a = c \wedge b = d$$



Definition (有序对 (Ordered Pairs)) (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$

Definition (Ordered Pairs (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Definition (Ordered Pairs (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Proof.

$$\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$$

Definition (Ordered Pairs (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Proof.

$$\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$$


CASE I : $a = b$

CASE II : $a \neq b$




Definition (Ordered Pairs (Norbert Wiener; 1914))

$$(a, b) \triangleq \left\{ \left\{ \{a\}, \emptyset \right\}, \left\{ \{b\} \right\} \right\}$$



Definition (Ordered Pairs (Norbert Wiener; 1914))

$$(a, b) \triangleq \left\{ \left\{ \{a\}, \emptyset \right\}, \left\{ \{b\} \right\} \right\}$$



Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Theorem

$$(x_1, \dots, x_n) = (y_1, \dots, y_n) \iff x_1 = y_1 \wedge \dots x_n = y_n$$

Definition (笛卡尔积 (Cartesian Products))

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$

Definition (笛卡尔积 (Cartesian Products))

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$


$$X^2 \triangleq X \times X$$

Definition (笛卡尔积 (Cartesian Products))


The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$


$$X^2 \triangleq X \times X$$



$$\text{Ranks} = \{2, \dots, 10, J, Q, K, A\}$$



$$\text{Suits} = \{\}$$



$$\mathbb{Z}^2 \triangleq \mathbb{Z} \times \mathbb{Z}$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$(X \times Y) \times Z \neq X \times (Y \times Z)$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$(X \times Y) \times Z \neq X \times (Y \times Z)$$

$$A = \{1\} \quad (A \times A) \times A \neq A \times (A \times A)$$

Theorem (分配律 (Distributivity))

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$


$$A \times (B \setminus C) = (A \times B) \setminus (A \times C)$$

Theorem (分配律 (Distributivity))

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$A \times (B \setminus C) = (A \times B) \setminus (A \times C)$$



Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

$$X_1 \times X_2 \times \cdots \times X_n \triangleq (X_1 \times X_2 \times \cdots \times X_{n-1}) \times X_n$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

$$X_1 \times X_2 \times \cdots \times X_n \triangleq (X_1 \times X_2 \times \cdots \times X_{n-1}) \times X_n$$

$$X^n \triangleq \underbrace{X \times \cdots \times X}_n$$

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Definition (关系 (Relations))

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

If $A = B$, R is called a relation *on* A .

Definition (关系 (Relations))

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

If $A = B$, R is called a relation *on* A .

Definition (Notations)

$$(a, b) \in R \quad R(a, b) \quad aRb$$

Definition (Relations)

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

Examples

Definition (Relations)

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

Examples

- Both $A \times B$ and \emptyset are relations from A to B .

Definition (Relations)

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

Examples

- ▶ Both $A \times B$ and \emptyset are relations from A to B .
- ▶

$$< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$$

Definition (Relations)

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

Examples

- ▶ Both $A \times B$ and \emptyset are relations from A to B .



$$< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$$



$$D = \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid \exists q \in \mathbb{N} : a \cdot q = b\}$$

Definition (Relations)

A *relation* R from A to B is a *subset* of $A \times B$:

$$R \subseteq A \times B$$

Examples

- ▶ Both $A \times B$ and \emptyset are relations from A to B .



$$< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$$



$$D = \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid \exists q \in \mathbb{N} : a \cdot q = b\}$$

- ▶ P : the set of people

$$M = \{(a, b) \in P \times P \mid a \text{ is the mother of } b\}$$


$$B = \{(a, b) \in P \times P \mid a \text{ is the brother of } b\}$$

Important Relations:

Equivalence Relations

Ordering Relations

Functions (next class)



Outline:

3 Definitions

5 Operations

7 Properties

2 Special Relations

3 Definitions

$\text{dom}(R)$ $\text{ran}(R)$ $\text{fld}(R)$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (值域 (Range))

$$\text{ran}(R) = \{b \mid \exists a : (a, b) \in R\}$$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (值域 (Range))

$$\text{ran}(R) = \{b \mid \exists a : (a, b) \in R\}$$

Definition (域 (Field))

$$\text{fld}(R) = \text{dom}(R) \cup \text{ran}(R)$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$\text{dom}(R) = \mathbb{R} \quad \text{ran}(R) = \mathbb{R} \quad \text{fld}(R) = \mathbb{R}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$\text{dom}(R) = [1, 1] \quad \text{ran}(R) = [-1, 1] \quad \text{fld}(R) = [-1, 1]$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

(6)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

$$\implies \exists b. (a, b) \in R \tag{2}$$

(6)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

$$\implies \exists b. (a, b) \in R \tag{2}$$

(6)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

$$\implies \exists b. (a, b) \in R \tag{2}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{3}$$

(6)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

$$\implies \exists b. (a, b) \in R \tag{2}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{3}$$

$$\implies \exists b. \{a, b\} \in \bigcup R \tag{4}$$

(6)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{1}$$

$$\implies \exists b. (a, b) \in R \tag{2}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{3}$$

$$\implies \exists b. \{a, b\} \in \bigcup R \tag{4}$$

$$\implies \exists b. a \in \bigcup \bigcup R \tag{5}$$

$$\tag{6}$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \quad (1)$$

$$\implies \exists b. (a, b) \in R \quad (2)$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \quad (3)$$

$$\implies \exists b. \{a, b\} \in \bigcup R \quad (4)$$

$$\implies \exists b. a \in \bigcup \bigcup R \quad (5)$$

$$\implies a \in \bigcup \bigcup R \quad (6)$$

5 Operations

$$R^{-1} \quad R|_X \quad R[X] \quad R^{-1}[Y] \quad R \circ S$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

$$\leq = \{(x, y) \mid x \leq y\} \subseteq \mathbb{R} \times \mathbb{R} \quad \leq^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

$$\leq = \{(x, y) \mid x \leq y\} \subseteq \mathbb{R} \times \mathbb{R} \quad \leq^{-1} = \geq \triangleq \{(x, y) \mid x \geq y\}$$

Theorem

$$(R^{-1})^{-1} = R$$

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

(3)

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

$$\iff (b, a) \in R^{-1} \quad (2)$$

$$(3)$$

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

$$\iff (b, a) \in R^{-1} \quad (2)$$

$$\iff (a, b) \in R \quad (3)$$

Theorem (关系的逆)

$$R, S \subseteq A \times B$$

$$(R \cup S)^{-1} = R^{-1} \cup S^{-1}$$

$$(R \cap S)^{-1} = R^{-1} \cap S^{-1}$$

$$(R \setminus S)^{-1} = R^{-1} \setminus S^{-1}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (右限制 (Right-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq Y$. The *right-restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid y \in S\}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (右限制 (Right-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq Y$. The *right-restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid y \in S\}$$

Definition (限制 (Restriction))

Suppose $R \subseteq X \times X$ and $S \subseteq X$. The *restriction* relation of R to S is

$$R|_S = \{(x, y) \in R \mid x \in S \wedge y \in S\}$$

example

Definition (像 (Image))

The *image* of X under R is the set

$$R[X] = \{b \in \text{ran}(R) \mid \exists a \in X. (a, b) \in R\}$$

Definition (像 (Image))

The *image* of X under R is the set

$$R[X] = \{b \in \text{ran}(R) \mid \exists a \in X. (a, b) \in R\}$$

$$R[a] \triangleq R[\{a\}] = \{b \mid (a, b) \in R\}$$

Definition (逆像 (Inverse Image))

The *inverse image* of Y under R is the set

$$R^{-1}[Y] = \{a \in \text{dom}(R) \mid \exists b \in Y : (a, b) \in R\}$$

Definition (逆像 (Inverse Image))

The *inverse image* of Y under R is the set

$$R^{-1}[Y] = \{a \in \text{dom}(R) \mid \exists b \in Y : (a, b) \in R\}$$

$$R^{-1}[b] \triangleq R^{-1}[\{b\}] = \{a \mid (a, b) \in R\}$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R^{-1}[R[X]] \text{ ? } X$$

$$R[R^{-1}[Y]] \text{ ? } Y$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R^{-1}[R[X]] \stackrel{?}{=} X$$

$$R[R^{-1}[Y]] \stackrel{?}{=} Y$$



Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2 : (a, b) \in R$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2 : (a, b) \in R$$

$$\iff \exists a \in X_1 : (a, b) \in R \vee \exists a \in X_2 : (a, b) \in R$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2 : (a, b) \in R$$

$$\iff \exists a \in X_1 : (a, b) \in R \vee \exists a \in X_2 : (a, b) \in R$$

$$\iff b \in R[X_1] \vee b \in R[X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2 : (a, b) \in R$$

$$\iff \exists a \in X_1 : (a, b) \in R \vee \exists a \in X_2 : (a, b) \in R$$

$$\iff b \in R[X_1] \vee b \in R[X_2]$$

$$\iff b \in R[X_1] \cup R[X_2]$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the *relation*

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the *relation*

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the *relation*

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the *relation*

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\}$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\} \quad (S \circ S) \circ S =$$

Definition (复合 (Composition; $R \circ S, R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$


$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$


$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\} \quad (S \circ S) \circ S = \{(2, 2), (2, 3)\}$$



$$|R \circ S| =$$



$$|R \circ S| = 7$$

$$\leq \circ \leq =$$

$$\leq \circ \leq = \leq$$

$$\leq \circ \leq = \leq$$

$$\leq \circ \geq =$$

$$\leq \circ \leq = \leq$$

$$\leq \circ \geq = \mathbb{R} \times \mathbb{R}$$

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, c) \in R \wedge (c, a) \in S \quad (3)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, c) \in R \wedge (c, a) \in S \quad (3)$$

$$\iff \exists c. (c, b) \in R^{-1} \wedge (a, c) \in S^{-1} \quad (4)$$

$$(5)$$

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, c) \in R \wedge (c, a) \in S \quad (3)$$

$$\iff \exists c. (c, b) \in R^{-1} \wedge (a, c) \in S^{-1} \quad (4)$$

$$\iff (a, c) \in S^{-1} \circ R^{-1} \quad (5)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. \left((a, c) \in T \wedge (c, b) \in R \circ S \right) \quad (2)$$

$$\iff \exists c. \left((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R) \right) \quad (3)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

$$\iff \exists d. ((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R) \quad (5)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. \left((a, c) \in T \wedge (c, b) \in R \circ S \right) \quad (2)$$

$$\iff \exists c. \left((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R) \right) \quad (3)$$

$$\iff \exists d. \exists c. \left((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R \right) \quad (4)$$

$$\iff \exists d. \left((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R \right) \quad (5)$$

$$\iff \exists d. \left((a, d) \in S \circ T \wedge (d, b) \in R \right) \quad (6)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

$$\iff \exists d. ((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R) \quad (5)$$

$$\iff \exists d. ((a, d) \in S \circ T \wedge (d, b) \in R) \quad (6)$$



燕小六：“帮我照顾好我七舅姥爷和我外甥女”

“舅姥爷”：姥姥/外婆的兄弟

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

$$G = B \circ (M \circ M) = (B \circ M) \circ M$$

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

$$G = B \circ (M \circ M) = (B \circ M) \circ M$$

“舅姥爷”：妈妈的舅舅

Theorem (关系的复合)

$$(X \cup Y) \circ Z = (X \circ Z) \cup (Y \circ Z)$$


$$(X \cap Y) \circ Z \subseteq (X \circ Z) \cap (Y \circ Z)$$

7 Properties

$$R \subseteq X \times X$$

Definition (Reflexive)


$$\forall a \in X : (a, a) \in R$$



$$R \subseteq X \times X$$

Definition (Reflexive)

$$\forall a \in X : (a, a) \in R$$



Definition (Irreflexive)

$$\forall a \in X : (a, a) \notin R$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 2), (2, 2), (2, 3), (3, 1)\}$$

$$R \subseteq X \times X$$

Definition (Symmetric)

$$\forall a, b \in X : aRb \implies bRa$$



$$R \subseteq X \times X$$

Definition (Symmetric)

$$\forall a, b \in X : aRb \implies bRa$$



Definition (AntiSymmetric)

$$\forall a, b \in X : (aRb \wedge bRa) \implies a = b$$

$$R \subseteq X \times X$$

Definition (Symmetric)

$$\forall a, b \in X : aRb \implies bRa$$



Definition (AntiSymmetric)

$$\forall a, b \in X : (aRb \wedge bRa) \implies a = b$$

>

$$R \subseteq X \times X$$

Definition (Symmetric)

$$\forall a, b \in X : aRb \implies bRa$$



Definition (AntiSymmetric)

$$\forall a, b \in X : (aRb \wedge bRa) \implies a = b$$

> *is* antisymmetric.

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$\{(1, 1), (2, 2), (3, 3)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$


$$\{(1, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 1), (2, 3)\}$$

$$R \subseteq X \times X$$

Definition (Transitive)

$$\forall a, b, c \in X : aRb \wedge bRc \implies aRc$$



$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 3)\}$$

$$A = \{1, 2, 3\}, R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 3)\}$$

$$\emptyset$$

$$R \subseteq X \times X$$

Definition (Connex)

$$\forall a, b \in X : aRb \vee bRa$$

$$R \subseteq X \times X$$

Definition (Connex)

$$\forall a, b \in X : aRb \vee bRa$$

Definition (Trichotomous)

$$\forall a, b \in X : \text{ exactly one of } aRb, bRa, \text{ or } a = b \text{ holds}$$

Theorem

$$R \text{ is reflexive} \iff I \subseteq R$$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is reflexive} \iff I \subseteq R$$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is symmetric} \iff R^{-1} = R$$

Theorem

$$R \text{ is reflexive} \iff I \subseteq R$$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is symmetric} \iff R^{-1} = R$$

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

Theorem

$$R \text{ is reflexive} \iff I \subseteq R$$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is symmetric} \iff R^{-1} = R$$

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

$$(1, 2), (2, 3), (1, 3), (4, 4)$$

Equivalence Relations

Definition (Equivalence Relation)

R is an *equivalence relation* on X iff R is

- ▶ reflexive
- ▶ symmetric
- ▶ transitive

Definition (Equivalence Relation)

R is an *equivalence relation* on X iff R is

- ▶ reflexive
- ▶ symmetric
- ▶ transitive

$$= \in \mathbb{R} \times \mathbb{R}$$

Definition (Equivalence Relation)

R is an *equivalence relation* on X iff R is

- ▶ reflexive
- ▶ symmetric
- ▶ transitive

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\| \in \mathbb{L} \times \mathbb{L}$$

Definition (Equivalence Relation)

R is an *equivalence relation* on X iff R is

- ▶ reflexive
- ▶ symmetric
- ▶ transitive

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

$$a \sim b \iff a \% 12 = b \% 12$$

Definition (Equivalence Relation)

R is an *equivalence relation* on X iff R is

- ▶ reflexive
- ▶ symmetric
- ▶ transitive

$$= \in \mathbb{R} \times \mathbb{R}$$


$$\parallel \in \mathbb{L} \times \mathbb{L}$$

$$a \sim b \iff a \% 12 = b \% 12$$


Why are equivalence relations important?

Equivalence Relations as Abstractions

Equivalence Relations as Abstractions




Equivalence Relations as Abstractions



“全国人民代表大会各省代表团”


Equivalence Relations as Abstractions



“全国人民代表大会各省代表团”

Equivalence Relation \iff Partition

Partition



“不空、不漏、不重”

Definition (Partition)

A family of sets $\{A_\alpha : \alpha \in I\}$ is a *partition* of X if

(i)

$$\forall \alpha \in I : A_\alpha \neq \emptyset$$

(ii)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

(iii)

$$\forall \alpha, \beta \in I : A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (Partition)

A family of sets $\{A_\alpha : \alpha \in I\}$ is a *partition* of X if

(i)

$$\forall \alpha \in I : A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I \exists x \in X : x \in A_\alpha)$$

(ii)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

(iii)

$$\forall \alpha, \beta \in I : A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (Partition)

A family of sets $\{A_\alpha : \alpha \in I\}$ is a *partition* of X if

(i)

$$\forall \alpha \in I : A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I \exists x \in X : x \in A_\alpha)$$

(ii)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

$$(\forall x \in X \exists \alpha \in I : x \in A_\alpha)$$

(iii)

$$\forall \alpha, \beta \in I : A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (Partition)

A family of sets $\{A_\alpha : \alpha \in I\}$ is a *partition* of X if

(i)

$$\forall \alpha \in I : A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I \exists x \in X : x \in A_\alpha)$$

(ii)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

$$(\forall x \in X \exists \alpha \in I : x \in A_\alpha)$$

(iii)

$$\forall \alpha, \beta \in I : A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

$$(\forall \alpha, \beta \in I : A_\alpha \cap A_\beta \neq \emptyset \implies A_\alpha = A_\beta)$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (Equivalence Class)

The *equivalence class* of a modulo R is a **set**:

$$[a]_R = \{b \in X : aRb\}$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (Equivalence Class)

The *equivalence class* of a modulo R is a **set**:

$$[a]_R = \{b \in X : aRb\}$$

Definition (Quotient Set)

The *quotient set* is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X : [a]_R \neq \emptyset$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X : [a]_R \neq \emptyset$$

$$\forall a \in X : \exists b \in X : a \in [b]_R$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X : [a]_R \neq \emptyset$$

$$\forall a \in X : \exists b \in X : a \in [b]_R$$

Theorem

$$\forall a \in X, b \in X : [a]_R \cap [b]_R = \emptyset \vee [a]_R = [b]_R$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X : [a]_R \neq \emptyset$$

$$\forall a \in X : \exists b \in X : a \in [b]_R$$

Theorem

$$\forall a \in X, b \in X : [a]_R \cap [b]_R = \emptyset \vee [a]_R = [b]_R$$

$$\forall a \in X, b \in X : [a]_R \cap [b]_R \neq \emptyset \implies [a]_R = [b]_R$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi : a \in S \wedge b \in S\}$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi : a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X .

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi : a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X .

$$\forall x \in X : xRx$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi : a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X .

$$\forall x \in X : xRx$$

$$\forall x, y \in X : xRy \implies yRx$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi : a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi : a \in S \wedge b \in S\}$$


Theorem

R is an equivalence relation on X .

$$\forall x \in X : xRx$$

$$\forall x, y \in X : xRy \implies yRx$$

$$\forall x, y, z \in X : xRy \wedge yRz \implies xRz$$



Equivalence Relation \iff Partition

Definition

$$\sim \subseteq \mathbb{N}^2 \times \mathbb{N}^2$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_{\mathbb{N}} c$$

Definition

$$\sim \subseteq \mathbb{N}^2 \times \mathbb{N}^2$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_{\mathbb{N}} c$$

Theorem

\sim is an equivalence relation.

Definition

$$\sim \subseteq \mathbb{N}^2 \times \mathbb{N}^2$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_{\mathbb{N}} c$$

Theorem

\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition

$$\sim \subseteq \mathbb{N}^2 \times \mathbb{N}^2$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_{\mathbb{N}} c$$

Theorem

\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition (\mathbb{Z})

$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

Definition

$$\sim \subseteq \mathbb{N}^2 \times \mathbb{N}^2$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_{\mathbb{N}} c$$

Theorem


\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition (\mathbb{Z})

$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

$$[(1, 3)]_{\sim} = \{(0, 2), (1, 3), (2, 4), (3, 5), \dots\} \triangleq -2 \in \mathbb{Z}$$



$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

Definition $(+_{\mathbb{Z}})$

$$[(m_1, n_1)] +_{\mathbb{Z}} [(m_2, n_2)] = [m_1 +_{\mathbb{N}} m_2, n_1 +_{\mathbb{N}} n_2]$$

Definition $(+_{\mathbb{Z}})$

$$[(m_1, n_1)] +_{\mathbb{Z}} [(m_2, n_2)] = [m_1 +_{\mathbb{N}} m_2, n_1 +_{\mathbb{N}} n_2]$$

Definition $(\cdot_{\mathbb{Z}})$

$$\begin{aligned} & [(m_1, n_1)] \cdot_{\mathbb{Z}} [(m_2, n_2)] \\ &= [m_1 \cdot_{\mathbb{N}} m_2 +_{\mathbb{N}} n_1 \cdot_{\mathbb{N}} n_2, m_1 \cdot_{\mathbb{N}} n_2 +_{\mathbb{N}} n_1 \cdot_{\mathbb{N}} m_2] \end{aligned}$$

Definition

$$\sim \subseteq (\mathbb{Z} \times \mathbb{Z} \setminus \{0_{\mathbb{Z}}\})^2$$

$$(a, b) \sim (c, d) \iff a \cdot_{\mathbb{Z}} d = b \cdot_{\mathbb{Z}} c$$


Definition

$$\sim \subseteq (\mathbb{Z} \times \mathbb{Z} \setminus \{0_{\mathbb{Z}}\})^2$$

$$(a, b) \sim (c, d) \iff a \cdot_{\mathbb{Z}} d = b \cdot_{\mathbb{Z}} c$$

Definition (\mathbb{Q})

$$\mathbb{Q} \triangleq \mathbb{Z} \times \mathbb{Z} / \sim$$




$$\mathbb{Q} \triangleq \mathbb{Z} \times \mathbb{Z} / \sim$$

How to define \mathbb{R} as equivalence classes of ordered pairs of \mathbb{Q} ?

How to define \mathbb{R} as equivalence classes of ordered pairs of \mathbb{Q} ?



Thank
You!



Office 926

hfwei@nju.edu.cn