

(十二) 图论: 对偶 (Duality: Matching, Network Flow, and Connectivity)

魏恒峰

hfwei@nju.edu.cn

2021 年 05 月 27 日





let's get
married
today




The Marriage Problem (Philip Hall, 1935)


If there is a finite set of **girls**, each of whom knows several **boys**,
under what conditions can all the girls marry boys in such a way that
each girl marries **a boy** that she knows?


The Marriage Problem (Philip Hall, 1935)

If there is a finite set of **girls**, each of whom knows several **boys**,
under what conditions can all the girls marry boys in such a way that
each girl marries **a boy** that she knows?



Philip Hall (1904 ~ 1982)






Definition (X -Perfect Matching (X -Saturating Matching))


Let $G = (X, Y, E)$ be a bipartite graph.

An X -perfect matching of G is a matching which covers each vertex in X .

$$|X| \leq |Y|$$



$$|X| \leq |Y|$$



$$\forall W \subseteq X. |W| \leq |N(W)|$$

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is an X -perfect matching of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

By induction on the number $|X|$ of vertices in X .

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$.

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$. I am married!

Induction Hypothesis: Suppose that it holds if $|X| < m$.

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$. I am married!

Induction Hypothesis: Suppose that it holds if $|X| < m$.

Induction Step: Consider the case $|X| = m$.

Consider the case $|X| = m$.

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a $(X - \{x\})$ -perfect matching in G' .

Consider the case $|X| = m$.

- CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a $(X - \{x\})$ -perfect matching in G' .

Therefore, there is a $(X - \{x\})$ -perfect matching in G .

- ▶ CASE II: There is a set of $k < m$ girls in X who know k boys in Y .

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is a X -perfect matching of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

Definition (M -alternating Paths)

Definition (M -alternating Paths)

Definition (M -augmenting Paths)

By contradiction.

By contradiction.

Suppose that there is *no* X -perfect matching.

By contradiction.


Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.

By contradiction.

Suppose that there is *no* X -perfect matching.


We show that Hall's Condition is violated for *some* $S \subseteq X$.



By contradiction.

Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

By contradiction.

Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

Let $u \in X$ be a vertex of X not saturated by M .

By contradiction.

Suppose that there is *no* X -perfect matching.


We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

Let $u \in X$ be a vertex of X not saturated by M .

Consider all *M -alternating paths* starting from u .




$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.



$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.

$S \triangleq$ the set of vertices in X reachable from u by M -alternating paths.




$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.

$S \triangleq$ the set of vertices in X reachable from u by M -alternating paths.


We will show that

$$T = N(S) \wedge |T| = |S - \{u\}|$$

$$|T| = |S - \{u\}|$$




$$|T| = |S - \{u\}|$$




We show that there is a bijection from T to $S - \{u\}$.

$$|T| = |S - \{u\}|$$




We show that there is a bijection from T to $S - \{u\}$.
 M matches T with $S - \{u\}$.

$$T = N(S)$$




$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$


$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T$


$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\}$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\} \implies \{s, y\} \notin M$$

$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\} \implies \{s, y\} \notin M \implies y \in T$$

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is a X -perfect matching of G iff


$$\forall W \subseteq X. |W| \leq |N_G(W)|$$


algorithm

Definitions (Vertex Cover (点覆盖))

A **vertex cover** of a graph G is a set $Q \subseteq V(G)$ that **covers** all edges.

$$\forall e \in E(G). e \cap Q \neq \emptyset$$







examples


Theorem (König (1931), Egerváry (1931))

If G is a bipartite graph, then the maximum size of a matching in G equals the minimum size of a vertex cover of G






$$R = Q \cap X \quad T = Q \cap Y$$



$$R = Q \cap X \quad T = Q \cap Y$$

$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G




$$R = Q \cap X \quad T = Q \cap Y$$

$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G

G has no edges from $X - R$ to $Y - T$.



$$R = Q \cap X \quad T = Q \cap Y$$


$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G


G has no edges from $X - R$ to $Y - T$.

H has a R -perfect matching and H' has a T -perfect matching.

H has a R -perfect matching and H' has a T -perfect matching.




H has a R -perfect matching and H' has a T -perfect matching.



By contradiction.


H has a R -perfect matching and H' has a T -perfect matching.



By contradiction.

$$\exists S \subseteq R. |N_H(S)| < |S|$$

H has a R -perfect matching and H' has a T -perfect matching.



By contradiction.

$$\exists S \subseteq R. |N_H(S)| < |S|$$


$T \cup (R - S + N_H(S))$ is a smaller vertex cover than Q

independent sets and covers

Definition (Network (网络))


A **network** is a **digraph** with

- ▶ a distinguished **source vertex** s ,
- ▶ a distinguished **sink vertex** t ,
- ▶ a **capacity** $c(e) \geq 0$ on each edge e




Definition (Flow (流))

A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



Definition (Flow (流))


A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



$$f^+(v) = \sum_{vw \in E} f(vw)$$

Definition (Flow (流))

A **flow** f is a **function** that assigns a value $f(e)$ to each edge e .



$$f^+(v) = \sum_{vw \in E} f(vw) \quad f^-(v) = \sum_{uv \in E} f(uv)$$

Definition (Feasible)


A flow f is **feasible** if it satisfies

Capacity Constraints:

$$\forall e \in E(G). 0 \leq f(e) \leq c(e)$$

Conservation Constraints:


$$\forall v \in V(G) - \{s, t\}. f^+(v) = f^-(v)$$



Definition (Value (值))

The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$



Definition (Value (值))

The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$



Definition (Value (值))



The **value** $\text{val}(f)$ of a **flow** f is



$$\text{val}(f) = f^-(t) = f^+(s).$$





Definition (Maximum Flow (最大流))

A **maximum flow** is a **feasible flow** of maximum **value**.





$$s \rightarrow x \rightarrow v \rightarrow t$$



$$s \rightarrow x \rightarrow v \rightarrow t$$


Definition (f -augmenting Paths (增广路径))


$$\min_{e \in E(P)} \epsilon(e)$$

Definition (Source/Sink Cut (割))

In a network, a **source/sink cut** $[S, T]$ consists of the edges **from** a **source set** S **to** a **sink set** T , where


$$V = S \uplus T \wedge s \in S \wedge t \in T$$





Definition (Capacity of Cut (割的容量))

$$\text{cap}(S, T) = \sum_{u \in S, v \in T, uv \in E} c(uv)$$




Definition (Minimum Cut (最小割))

A **minimum cut** is a **cut** of minimum value.

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.


$$\text{val}(f) \leq \text{cap}(S, T).$$



Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S)$$

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S)$$

Theorem (Weak Duality (弱对偶定理))

Let f be any feasible *flow* and $[S, T]$ be any source/sink *cut*.


$$\text{val}(f) \leq \text{cap}(S, T).$$



$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S) \leq \text{cap}(S, T)$$

Lemma

$$\max_f \text{val}(f) \leq \min_{[S,T]} \text{cap}(S, T)$$



What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?


$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^{-1}(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$

What if $\text{val}(f) = \text{cap}(S, T)$ for some flow f and some cut $[S, T]$?

$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^{-1}(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$



Theorem (Max-flow Min-cut Theorem (Ford and Fulkerson; 1956))

$$\max_f \text{val}(f) = \min_{[S,T]} \text{cap}(S,T)$$


(Strong Duality)




L. R. Ford Jr. (1927 ~ 2017)



D. R. Fulkerson (1924 ~ 1976)



Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn